

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

#### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

#### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/

#### HARVARD UNIVERSITY




LIBRARY OF THE MINERALOGICAL LABORATORY

UNIVERSITY MUSEUM

Transferred to
CABOT SCIENCE LIBRARY
June 2005







Quality is assured as we permit return of specimens at our expense.



#### CATALOG PRICES POST-PAID

§ . 50
1.00
1.50

#### REFUND OF CATALOG PRICE

Will be made against first \$10.00 purchase, or on returning catalog, if not wanted.

QE 287 F6

## Complete Mineral Catalog

COMPILED BY W. M. FOOTE

TWELFTH EDITION, ENTIRELY REVISED AND ENLARGED
WITH THREE HUNDRED FIGURES AND PLATES
COVER ILLUSTRATION IODYRITE

PUBLISHED BY THE

## Foote Mineral Company

107 North 19th Street, Philadelphia, Pa., U. S. A.

ESTABLISHED 1876 BY DR. A. E. FOOTE

#### **CONTENTS**

PART I. Introductory	
Acknowledgment	(
Rare Element Minerals for Manufacturers	(
Quality	7
Terms. Free Transportation	7
The "Traveling Exhibit"	8
Oak Chests	8
Pasteboard Trays	I
Labels	12
Sizes of Specimens	15
Note	18
	19
PART II. Advanced Systematic Collections. Synopsis of Dana's "System of Mineralogy"	
Description of Advanced Collections	23
Remarks on the List and Synopsis	29
List of the Complete Type and Other Advanced Collections, Embraced in a	
Synopsis of Dana's System	32
PART III. INDEX TO COMPLETE TYPE COLLECTION. DANA'S SYSTEM. PRICE LIST OF HAND SIZE SPECIMENS	
Abbreviations, Etc	212
Alphabetical Index and Price List	213
PART IV. ELEMENTARY SYSTEMATIC COLLECTIONS	
Description of Collections	242 245
PART V. Economic Mineralogy. Industrial Minerals and Ores	
Description of Collections	254
Lists of School of Mines and Mining Collections	260
PART VI. CRYSTALLOGRAPHY. LOOSE CRYSTALS FOR MEASUREMENT AND STUDY	
Advanced Collections. Description	276
List of Complete Crystal Collection	278
Price List and Index to the Collection	293
Elementary. School Crystal Collection	296
Lecture Table Crystals	298
PART VII. Physical Mineralogy	
Sets Illustrating Hardness, Structure, Color, Effect of Radium, Etc	300
List of Complete Physical Series	303
PART VIII. CHEMICAL MINERALOGY	
Collections of Minerals for Analysis	308
Laboratory List of Minerals Sold by Weight	311

## PART I

## Introductory

Terms, Labels, Trays, Sizes, Cabinets, Etc.

#### Acknowledgment.

The complete revision and expansion of most of the lists throughout the present edition is exemplified in the very useful combining of the Complete Type Collection List with the oft published Synopsis of Dana's "System of Mineralogy." The different types and varieties thus follow each species heading with fuller descriptions of the form and physical characters of each than was practicable formerly. A feature which will especially commend itself to mineralogists, is the insertion in proper position in the systematic arrangement, of all recently discovered minerals. This includes the minerals described in the new "Second Appendix" to the "System," access to the manuscript being kindly given by Prof. Dana. The compiler is especially indebted to Dr. F. Ward for the selection of data from the Supplement and two Appendixes, and locating the new minerals; to Prof. Wm. E. Ford for advice concerning species of doubtful relationship; to Prof. Amos P. Brown for suggesting the presentation of the hardness by group; and finally to Prof. Edward S. Dana and Messrs. Wiley & Sons for permission to reproduce from the "System" the numerous figures which add so much to the value of the catalog.

#### Rare Minerals

#### In Commercial Lots for Manufacturers

Progress in metallurgy, lighting and other special industries in the past decade has been marked by a rapidly increasing One of the obstacles which the techuse of the rare metals. nologist faces is the limited and irregular supply of raw material. Our facilities for acquiring these ores are exceptional, as we employ traveling experts and are constantly receiving samples from correspondents throughout the world. The results of thirtythree years of wholesale collecting affords a choice of over two thousand different varieties of minerals, as listed in the Complete Type Collection, Part II. Specimen prices of several hundred useful minerals are given in Part V, Economic Mineralogy. The more important are mentioned in the Laboratory List, Part VIII, with prices by weight for samples. Correspondence is solicited with consumers or experimenters desiring ton lots.



#### Highest Awards

AT THE EXPOSITIONS OF

PHILADELPHIA, 1876 CINCINNATI, 1881 NEW ORLEANS, 1884-85 NEW ORLEANS, 1885-86 LOUISVILLE, 1886 LONDON, 1887 PARIS, 1889 PARIS, 1900

# Assurance of Quality We Pay Transportation To Any Address In The World.

SPECIFIC GUARANTEE. If you do not like the specimens return them at our expense. We take responsibility of loss or breakage in transit. The risk of double transportation we assume, gives assurance that your requirements will be met.

WE REFER TO curators and teachers of mineralogy in all countries.

PRICES ARE UNIFORM. The "one-price" system simplifies buying, and wins universal favor.

PAYMENTS. For convenience, we accept the following rate of exchange: \$1.00 = 4/- = M. 4. = Fcs. 5. = L. 5.

## The "Traveling Exhibit.' An Oak Mineral Chest

Is presented, if requested, with each purchase of Hand size or larger specimens, totaling \$20.00 or over. If chest is not requested,

#### we deduct instead 10 per cent.

from total price. These offers apply only when all specimens are kept.



PLATE I. OAK CHESTS NOS. I AND 2.

OUR DISTRIBUTION of specimens (transportation paid) takes two forms:—

FIRST.—Shipments to those who prefer to select from the specimens themselves. Our knowledge of the needs of the buyer are often indefinite and the selection we make may not always accord with his individual taste. The result is that some specimens may be returned at large expense to the seller and trouble to the buyer.

DISCRIMINATING BUYERS are requested to give us full advice as to their requirements, thus permitting us to choose specimens which may win entire approval.

SECOND.—Many are able to order from a catalog, and to such our material gives universal satisfaction. Those ordering in this direct way benefit by the chest offer, or equivalent 10 per cent. allowance, having always the right of prompt rejection.

THE CHESTS are iron-bound and made in the best and strongest manner, of  $\frac{7}{8}$  in. (22 mm.) first quality oak, antique wax finish. There are three sizes:—

- No. 1. Flat Mineral Chest. Measures inside  $23\frac{1}{2} \times 17\frac{7}{8} \times 2\frac{3}{4}$  in. (60 x 45 x 7 cm.). Given with not less than \$20.00 worth of minerals. See upper chest in Plate I.
- No. 2. Two-tray Mineral Chest, with handles. Measures inside 245% x 19 x 634 in. (63 x 48½ x 17 cm.). Given with not less than \$35.00 worth of minerals. See Plate II, also lower chest in Plate I.
- No. 3. Four-tray Mineral Chest, with handles. Measures inside  $24\frac{5}{8} \times 19 \times 12\frac{3}{4}$  in.  $(63 \times 48\frac{1}{2} \times 32\frac{1}{2} \text{ cm.})$ . Given with not less than \$50.00 worth of minerals.

The Oak Trays are of uniform size, measuring inside  $23\frac{1}{2}$  x  $17\frac{7}{8}$  x  $2\frac{3}{4}$  in. (60 x 45 x 7 cm.). They are made of  $\frac{1}{2}$  inch (13 mm.) solid oak, with hand-holes at ends, being more convenient to handle than drawers. Each oak tray is fitted with 25 white pasteboard trays measuring  $4\frac{3}{4}$  x  $3\frac{9}{10}$  in. (12 x 9 cm.). Two of these 12 x 9 cm. pasteboard trays are interchangeable with one 18 x 12 cm. tray, or one is interchangeable with two 9 x 6 cm. trays. Thus each oak tray holds from 15 to 50 fine pasteboard trays. These are indispensable in keeping orderly arrangement.

THE COST (WHEN NOT GIVEN WITH SPECIMENS) delivered to any address is, for No. 1, \$4.00; No. 2, \$7.00; No. 3, \$12.00. If made singly by a good cabinet maker they would cost double these figures. Chest prices do not include pasteboard trays.

If a chest is not desired, then deduct 10 per cent. from your remittance (of \$20.00 or over.) This is the average per cent. saved us by avoiding return transportation and handling, when you retain the entire lot. This allowance is only on hand size or larger specimens and only when no specimens are returned.



PLATE II. OAK CHEST NO. 2, HOLDING ABOUT 60 HAND SIZE SPECIMENS OR 30 MUSEUM SIZE SPECIMENS




PLATE III

12 LABELS



PLATE IV. EXAMPLES OF SMALL LABLES USED.



PLATE V.

## Sample of Attached Label On Hand Size Specimen in Pasteboard Tray

LABELING is one of the features of our business in which our clients rely upon the careful and conscientious work of the trained mineralogists in our employ. The name of the mineral, both species and varietal, and the composition in chemical formula, are essential on a reference specimen. The correct locality is of importance and is given especial attention. The reference number in Dana's "System of Mineralogy" is useful in arranging a collection.

The larger size of the universal loose label makes it more easily read and permits mention of the crystalline system. A great disadvantage is the frequent misplacement of the loose labels. Hence our use of the small attached label. Every specimen leaving our establishment has pasted on the back one of these miniature labels. The museum size specimens are also accompanied by the large Exhibition Label when requested.



PLATE VI. SAMPLE MUSEUM SIZE SPECIMEN ON BLOCK-MOUNT. (In practice the small label is attached to back of specimen.)

#### Museum Size Specimens

Averaging 12 x 9 cm.  $(4\% \times 3\% \text{ in.})$ 

Weight averages about one kilogram  $(2\frac{1}{5} \text{ lbs. avd.})$ 

The illustration opposite shows the average size of specimens listed by us for private or public museums, for the school or college-class room, or for office or laboratory display, where large examples of showy appearance are desired. While intended for glass cases, as shown in Plate VII., they may be held in a drawer cabinet fitted with pasteboard trays, the drawer being 7 cm. (2¾in.) deep. In preparing collections in the museum size, especial attention is paid to the shapeliness of each specimen, and to the selection of bright colors and striking crystallizations, wherever this can be done without impairing the representative character of the collection.

Where specimens are desired for the lecture table or passing among the class, a generously proportioned specimen tells the various properties and characteristics of the mineral, much better than a smaller piece requiring longer study.

A PASTEBOARD TRAY ACCOMPANIES EACH SPECIMEN. See Plate III.

OUR IMPROVED PASTEBOARD BLOCK-MOUNT SUBSTITUTED FOR TRAY, if requested, with museum size specimens. See opposite page. It is made of extra heavy pasteboard, covered with fine white paper, glazed to resist dust. This neat and light paper block, affording a simple white background, displays the average specimen much more effectively than the old-fashioned and sombre wooden one. The depth of the mount is: Top, 9 cm. (3½ in.); base, 13 cm. (5 in.). The slanting front measures 4¼ cm. (15% in.). The length is 16 cm. (6¼ in.) and the height 2¼ cm. (½ in.). Price for extra blocks, express paid, \$8.00 per 100.

LABELING is illustrated in Plates IV., V. and VI.

Prices for museum size are double the prices for hand size, following mineral names in this catalog.



PLATE VII. COLLECTION OF MUSEUM SIZE SPECIMENS WITH EXHIBITION LABELS.

#### Hand Size Specimens Averaging 10 x 7 cm. (4 x 2% in.)

#### Average weight about 450 grams (approximately 1 lb. avd.)

"Size is a secondary factor in the utility of a specimen, but it is one about which opinions vary greatly. What have you found to be the most desirable size for mineral specimens, intended for use of students, and for practical purposes of reference?"

In 1907 the above inquiry was addressed to teachers of mineralogy throughout the world. In nearly 100 replies received, the average or composite size preferred was 10 x 7 cm.  $(4 \times 2\frac{3}{4})$  in.). Few preferred a smaller size and some preferred a size even larger. The result of our canvass of the opinion of experts led us to adopt this as our principal stock size, as illustrated in Plate V.

THE CHARACTERISTICS of the mineral and its associations are displayed far better in this size than is possible in a fragment. The representative character of the specimens in illustrating physical properties and crystallization, is considered of first importance, but incidentally many of the specimens are of attractive appearance.

PRICES given after mineral names in all collection lists, save the crystal lists, are for standard Hand Size Specimens averaging 10 x 7 cm.  $(4 \times 2\frac{3}{4})$  in.).

A PASTEBOARD TRAY ACCOMPANIES EACH SPECIMEN. See Plates III and V.

LABELS are illustrated in Plates IV and V.

#### Small Size Specimens Averaging 7 x 5 cm. (2\% x 2 in.)

Weight averages about 225 grams (1/2 lb. avd.)

If purchased in lots of less than 10 of one kind, the prices are the same as for the hand size specimens. When purchased for classes, in lots of 10 or more pieces of one kind, the cost per specimen is one-half the catalogued prices for hand size specimens. This includes delivery to any address, but does not include chests.

#### Drawer Cabinets

Prices include crating and transportation to any address. Made after our own designs, based on long experience in handling specimens, being like cases recently manufactured for our own equipment. The work of a conscientious cabinet maker is here combined with the practical knowledge of the mineralogist's requirements.

FINE SOLID MAHOGANY (7/8 in. or 12 mm.) is used for the drawer fronts and for all parts except the back of the case, sides, back and bottom of drawers, which are made of 5/8 in. (16 mm.) cherry. Quartered oak, at the same cost, will be used if specified in order.

FINISH is in three coats of shellac, rubbed to a dull light natural finish. (Stained dark if specified in order).

SOLID BRASS KNOBS, firmly secured on inside.

THE DRAWERS measure inside 28!4 in. wide x 17% in. deep x 2¾ in. high (72 cm. wide x 46 cm. deep x 7 cm. high). Each drawer will hold (1) 30 hand size trays, or (2) 24 hand size and 12 small size trays, or (3) 12 museum size and 6 hand size trays. A groove is cut in the sides of the drawer, which slides smoothly on runners. Corners are hand-dove-tailed.

- Size A. Eighteen drawers in two tiers. Measures, over all, 66 in. wide x 20½ in. deep x 36 in. high (168 cm. wide x 52 cm. deep x 92 cm. high). Holds about 600 to 700 specimens, averaging  $4 \times 2\frac{3}{4}$  in. (10 x 7 cm.). Price, \$72.00.
- Size B. Ten drawers in one tier. Measures, over all, 35 in. wide x 20½ in. deep x 38 in. high (85 cm. wide x 52 cm. deep x 89 cm. high). Holds about 350 to 400 specimens averaging 4 x 2¾ in. (10 x 7 cm.). See Plate IX. Price, \$45.00.
- Size C. Six drawers. Measures, over all,  $34\frac{1}{2}$  in. wide x  $20\frac{1}{2}$  in. deep x  $24\frac{1}{2}$  in. high (88 cm. wide x 52 cm. deep x 61 cm. high). To hold Collections Nos. 14, 27, or 111, about 200 specimens averaging  $4 \times 2\frac{3}{4}$  in. (10 x 7 cm.). Price, \$30.00.

Prices do not include duty on cabinets or chests delivered in the few countries where Customs' duty is charged on same.

#### Note

Since the appearance of our former complete catalog, an improvement in the form of label used by us and increasing care devoted to the preparation of material, has resulted in a general raising of the already high standard for which our collections are noted. As in the past, our aim is always to select the most typical representatives of each mineral obtainable in the mines and quarries of the world.

A price is given opposite each specimen in the collection lists, the figure quoted being for the popular hand size. One who does not desire an entire set exactly as listed by us, may thus make, without correspondence, an immediate selection of any of the specimens comprising the collection, with a knowledge of the cost of each item. Selections may also be made from the Alphabetical Price List in Part III.

All but the smallest collections may be purchased in the catalogued parts or installments, without proportionately increasing the cost. This permits the gradual purchase of the more expensive collections, each part filling important gaps in the growing nucleus and not being merely a detached section. The "collection price" for each set is less than the sum of the individual values of the specimens comprising it. Our collections are prepared a number at a time, thus effecting a material saving in labor-cost.

All collections listed, except Nos. 1A, 1 and 3A, are ready for shipment immediately on receipt of order.




PLATE VIII. ELEMENTARY SCHOOL CRYSTAL SET IN CABINET. REDUCED TO % DIAMETERS.

## PART II

Advanced Systematic Collections

Synopsis of Dana's "System of Mineralogy"



PLATE IX. TEN-DRAWER CABINET, HOLDING ABOUT 400 HAND SIZE SPECIMENS.

#### Advanced Systematic Collections

These are arranged according to the generally accepted classification in Dana's "System of Mineralogy," or in the "Text-book of Mineralogy" by the same author. They can be rearranged to accord with other classifications, as the purchaser may desire. The specimens have attached, labels and numbers to correspond to numbered lists.

Despite our exceptional facilities for securing minerals through our travelers or numerous correspondents, and by the purchase of old collections replete with historic rarities, it requires several years to assemble a Complete Type Collection, such as is here catalogued. Hence we have always some of these advanced collections in course of preparation.

The descriptions in the list, made from collections in stock, are moderate in their indication of perfection, definiteness, luster and the characteristics which go to make up good crystallizations. No attempt is made to indicate the beauty or attractiveness of the specimens beyond the bare statement of form and color, etc. The average standard of crystallization and general excellence, in any collection delivered by us, will be as high as is here indicated.

It should be kept in mind that wide variation in types is encountered by all endeavoring to recognize the innumerable forms of the mineral kingdom. The student who has mastered a few hundred specimens is often warned that they represent but the commoner types, and in the field new and unknown varieties confuse and puzzle him at every turn.

Practice in the examination of widely varying types means a fuller acquaintance with minerals and increased power of observation. Advanced courses in mineralogy include constant drill in the identification of a large series of minerals by sight and by the quick tests applicable in the field. Prominent teachers who are regular and large buyers of minerals, credit their success largely to these practical methods.

Therefore as complete a collection as possible should be selected. The rarer specimens, even if not carefully studied, will prove invaluable for reference and comparison.

#### No. 1A. Complete Type Collection

Twenty-five hundred museum size specimens, averaging 12 x 9 cm.  $(4\frac{3}{4} \times 3\frac{1}{2} \text{ in.})$ . Selected by us from the 2640 numbered and priced items in the list.

Intended for individuals or institutions desiring a collection, which for study or comparison, is fairly complete in the light of present knowledge. Over six hundred distinct species are represented, embracing the most important in Dana's "System," including the Supplement and two Appendixes.

The principal known varieties and types of common and rare species, the crystal forms described in the Complete Crystal List and the features of the Complete Physical Series, together with the specimens comprising the large Economic, Chemical and Rock-forming series catalogued elsewhere, all find a place in this complete general collection, duplication always being avoided.

The multiplication of local examples is avoided, unless a variation of type is thereby gained. The occurrence of the commercial minerals is especially considered worthy of illustration by as many examples as their variations demand.

Our last published catalog included a list completed to 1500 specimens. In its present enlarged form, reaching over 2500 specimens, there are comparatively few public or private collections which excel it in point of comprehensiveness and general value for reference purposes.

PRICES OF PARTS, delivered to any address, with pasteboard trays, in cabinets. Without cabinets, 10 per cent. less.

- PART I. Six hundred specimens, marked + or \* in list, (University Collection No. 5A) totaling over \$1000.00 Price with trays, in two 18-drawer cabinets . . . . . . . . \$900.00
- PART II. Nine hundred additional and generally rarer specimens, marked o, totaling \$2400.00. Price with trays, in three 18-drawer cabinets......\$2100.00

PART III. One thousand specimens selected from the remaining numbered items in list, including most of the very rarest species and varieties, totaling \$3400.00. With trays, in three 18-drawer cabinets..\$3000.00

#### No. 1. Specialist's Complete Type Collection

Twenty-five hundred hand size specimens, averaging 10 x 7 cm.  $(4 \times 2\frac{3}{4})$  in.). Like the preceding, but smaller in size.

PRICES OF PARTS, delivered to any address, with paste-board trays, in mahogany 18-drawer cabinets. Without cabinets, 10 per cent. less.

- PART I. Six hundred specimens, marked + or \*
  (Specialist's University Collection No.
  5), with trays, in 18-drawer cabinet.. \$450.00
- PART II. Nine hundred additional specimens, marked o, totaling \$1200.00. Price with trays, in two 18-drawer cabinets . . . . \$1050.00

#### No. 3A. Varietal Collection

Fifteen hundred museum size specimens, averaging  $12 \times 9$  cm.  $(4\frac{3}{4} \times 3\frac{1}{2})$  in.). The list comprises the minerals marked with +, \* or ° in the Complete Type Collection List. In general the specimens omitted from this collection, but which appear in No. 1A, are exceedingly rare, or they exhibit minor variations. The collection as a whole has been very carefully planned to include the most important varieties of more than 500 distinct species.

PRICES OF PARTS, delivered to any address, with pasteboard trays, in mahogany 18-drawer cabinets. Without cabinets, 10 per cent. less.

- PART I. Six hundred specimens, marked + or \* (University Collection No. 5A), with trays, in two 18-drawer cabinets.....\$900.00
- PART II. Nine hundred remaining specimens, marked o, totaling \$2400.00. With trays, in three 18-drawer cabinets.....\$2100.00

#### No. 3. Specialist's Varietal Collection

Fifteen hundred hand size specimens, averaging 10 x 7 cm. (4 x 23/4 in.). Same list as the preceding, but in smaller specimens. This collection is in stock, ready for immediate delivery.

Price, including delivery to any address, with pasteboard trays, in three mahogany 18-drawer cabinets, \$1500.00. Without cabinets, 10 per cent. less.

PURCHASE IN PARTS. Delivered to any address.

PART I. Six hundred specimens, marked + or \*
(Specialist's University Collection
No. 5), totaling over \$500.00. With
trays, in 18-drawer cabinet......\$450.00

PART II. Nine hundred remaining specimens, marked o, totaling \$1200.00. With trays, in two 18-drawer cabinets......\$1050.00

#### No. 5A. University Collection.

Six hundred museum size specimens, averaging 12 x 9 cm.  $(4\frac{3}{4} \times 3\frac{1}{2} \text{ in.})$ .

The University List, comprising the names marked with + or \*, aims to include such minerals as are taken up in most university work. The 300 or more distinct species emphasized by heavy type in Dana's "Text Book of Mineralogy," are all represented. Some of them are quite rare, but are chemically important and essential in the illustration of a comprehensive and thorough course in pure mineralogy. Examples are shown of most of the economic minerals which the student or expert may wish to recognize, because of their commercial value. Numerous well known varieties which are found with the ores are also worthy of representation, although not in themselves valuable.

Individual museum size specimens of most kinds may be purchased at double the hand size prices given after each type. The sum of such individual values, in the museum size, exceeds \$1000.00. The "collection price," including delivery to any address, with trays and two 18-drawer cabinets, is \$900.00. Without cabinets, 10 per cent. less.

Purchase in Parts. Free delivery, with pasteboard trays and cabinets. Without cabinets, 10 per cent. less.

- PART II. Two hundred and forty remaining specimens marked \*, with trays and 18-drawer cabinet.....\$420.00

#### No. 5. Specialist's University Collection

Six hundred hand size specimens, averaging 10 x 7 cm.  $(4 \times 2\%)$  in.). Same as the preceding, but smaller size.

Individual hand size specimens of most varieties are sold at the listed prices. These exceed \$500.00. The "collection price" for all the specimens is \$450.00, delivered to any address, with pasteboard trays and mahogany 18-drawer cabinet. Without cabinet, 10 per cent. less.

Purchase in Parts. Free delivery, with pasteboard trays and cabinet.

- PART I. Three hundred and sixty specimens marked + (Specialist's College Collection No. 9), with trays and 18-drawer cabinet....\$261.00

#### No. 9A. College Collection

Three hundred and sixty museum size specimens, averaging 12 x 9 cm.  $(4\frac{3}{4} \times 3\frac{1}{2} \text{ in.})$ .

In the list which follows, the names marked + comprise the College Collection. No effort is spared in the work of abridgment, to make this as useful an advanced collection as the limited number of specimens will permit. The College List includes the most important minerals in Dana's "Text-book of Mineralogy," as well as all of the minerals contained in the Mining List and High School List. About two-thirds of the specimens are distinct species. As in the larger collections, every care is exercised that the College Collection may be thoroughly illustrative.

and serve as a useful adjunct to private study or class work. With its many pieces of striking form and beautiful color, it makes a most attractive exhibit in the lecture-hall or college museum.

Individual museum size specimens of most kinds may be purchased at double the hand size prices given after each name. The sum of such individual values, in the museum size, exceeds \$550.00. The "collection price," including delivery to any address, with trays and mahogany 18-drawer cabinet, is \$480.00. Without cabinet, 10 per cent. less.

Purchase in Parts. Free delivery, with pasteboard trays.

- PART I. One hundred and eighty specimens (Collection No. 14A), with trays and 18-drawer cabinet......\$210.00 (Part I without cabinet, \$162).
- PART II. One hundred and eighty remaining specimens, with trays, but without cabinet. \$270.00

#### No. 9. Specialist's College Collection

Three hundred and sixty hand size specimens, averaging 10 x 7 cm.  $(4 \times 2\frac{3}{4})$  in.). Same as the preceding, but smaller size.

Individual hand size specimens of most varieties are sold at the listed prices. These exceed \$275.00. The "collection price" for all the specimens is \$240.00, delivered to any address, with pasteboard trays and mahogany 10-drawer cabinet. Without cabinet, 10 per cent. less.

Purchase in Parts. Free delivery.

- PART II. One hundred and eighty remaining specimens, with trays, but without cabinet. \$135.00

#### Complete Type Collection

#### Embracing College, University and Varietal Collections

#### Remarks on the Synopsis of

"The System of Mineralogy"

#### Sixth Edition with Appendixes by Edward Salisbury Dana

In this synopsis are inserted in proper position, new species and distinct varieties or types mentioned in the "Supplement" and "First Appendix" to the "System" and likewise the new unpublished "Second Appendix" (1909), to the manuscript of which access was kindly given. These minerals are marked "S.," "I." or "II." in the Species No. column. Many of them, like some of the "related compounds" in the "System," are of doubtful position, owing to the incomplete knowledge concerning them.

THE FIRST SYNOPSIS of Dana's "System of Mineralogy," was published in the "Naturalist's Agency Catalogue," issued by Dr. A. E. Foote in 1876. The original "Table of Species," as it was called, gave in a condensed form, the physical and chemical characters of species, but did not mention varieties or subspecies. In its present form it is a combination of the last published "Synopsis," new matter from the Appendixes, and our "Complete Type Collection List."

THE NUMBERS OF THE COMPLETE TYPE COLLECTION LIST, I to 2640, are given in the first column.

COLLECTION LISTS are indicated in the second column by the following signs:—

Cross (+) indicates the commonest or most important minerals comprising the "College List" of 360 specimens.

Asterisk (\*) indicates 240 additional common or important minerals, which with the preceding, comprise the "University List" of 600 specimens.

Circle (°) indicates 900 rarer or less important minerals, largely varieties, which with the preceding, comprise the "Varietal List" of 1500 specimens.

DANA'S SPECIES NUMBERS, I to 824, are given in the third column in heavy type.

THE SPECIES NAME of each mineral is given in heavy type.

THE CHEMICAL COMPOSITION is generally expressed by the dualistic formula, which in the case of complex compounds, often presents the chemical constitution more clearly than does the empirical formula.

THE SYSTEM OF CRYSTALLIZATION follows.

THE CRYSTAL FORMS are indicated in the commoner or the best defined cases with frequent references to the figures.

THE STRUCTURE is mentioned where of importance.

THE COLOR is generally referred to under each species.

HARDNESS is broadly indicated under each group heading by giving the range from the softest to the hardest species.

VARIETAL NAMES, in ordinary type, are indented.

SUBSPECIES or "related compounds," also in ordinary type, are in alignment with the species names.

ALTERATIONS (pseudomorphs) are given only in the commoner types.

CRYSTAL FORMS are indicated by the letters used by Dana. Miller's symbols are shown in the Complete Crystal List. The figures, reproduced from the "System," are idealized to show the form clearly. In the majority of cases the actual crystals do not equal the figures in the matter of completeness nor symmetry; moreover the actual specimens generally consist of crystals on the matrix or grouped, unless marked "loose."

SIZE OF CRYSTALS is given in approximate terms, as follows: "Microscopic," usually under I mm. (about  $^{1}/_{25}$  in.); "minute," usually under 3 mm. (about  $^{1}/_{8}$  in.); "small," usually under I cm. (about  $^{2}/_{5}$  in.); "large," usually over 3 cm. (about  $^{13}/_{16}$  in.); "very large," usually over 9 cm. (about  $^{3}/_{2}$  in.).

SIZE OF SPECIMENS listed at \$0.20 to \$2.50 each, is generally hand size, averaging 10 x 7 cm.  $(4 \times 2\frac{3}{4})$  in.). Those priced higher are mostly of this size, but in very many instances they are smaller and even insignificant. They are, however, always of typical character.

NAMES OMITTED in this synopsis, but mentioned in Dana's "System," "Supplement" and two "Appendixes," are (1). Names rejected by Dana. (2). Names given to supposed minerals which have later proved to be rocks, mixtures or highly impure substances. (3). Some obscure varietal names not prominently mentioned by Dana. Generally these have only local significance. (4). Many names of unimportant compounds of such doubtful character, that their rank even as varieties, is questioned. (5). The less frequently used synonyms.

RELATIVE RARITY of good typical specimens of each kind, is indicated by the price following the description. Items not priced, are, as a class, rarely obtainable, although there are individual exceptions to the rule.

INDEX. The position of any mineral in the Synopsis, may be found by referring to the Index and Price-List in Part III.

#### Single Specimens

From the following list, comprising all the priced items, we have on hand (outside of prepared collections), most of those marked with +, \* or °, and can fill orders for individual specimens.

Of the priced items not so marked, many are not on hand in duplicate. They are generally quite rare and sometimes are historical rarities no longer found. Much of this "floating stock," comes to us in one or two specimens at a time, through the purchase of old collections. They are often immediately reserved for our large advanced collections, in course of preparation, or they are sent to clients who place advance orders.

YOUR DESIDERATA LIST should be filed with us if you want rare minerals. From time to time, we will submit for your inspection, specimens which may "fill in the gaps."

"The System of Mineralogy" of James Dwight Dana. Sixth Edition by Edward Salisbury Dana. Entirely rewritten and much enlarged. Illustrated with 1425 figures. Over 1200 pages with Appendix I. It may be purchased for \$12.50 of the publishers, Messrs. John Wiley & Sons, New York, or of Foote Mineral Company, (or of Chapman & Hall, London, £2-12-6).

## The General Classification

Complete Type Collection Accords With

The System of Mineralogy

of James Dwight Dana

#### Sixth Edition (See Note)

#### By Edward Salisbury Dana

- I. Native Elements.
- II. Sulphides, Selenides, Tellurides, Arsenides, Antimonides.
- III. Sulpho-Salts—Sulpharsenites, Sulphantimonites, Sulphobismuthites.
- IV. Haloids—Chlorides, Bromides, Iodides; Fluorides.
- V. Oxides.
- VI. Oxygen-Salts.
  - 1. Carbonates.
  - 2. Silicates, Titanates.
  - 3. Niobates, Tantalates.
  - 4. Phosphates, Arsenates, Vanadates; Antimonates. Nitrates.
  - 5. Borates. Uranates.
  - 6. Sulphates, Chromates, Tellurates.
  - 7. Tungstates, Molybdates.
  - 8. Iodates.
- VII. Salts of Organic Acids—Oxalates, Mellates, Etc.
- VIII. Hydrocarbon Compounds.

Note—New Minerals described in the Supplement, First Appendix and New Second Appendix to the System, are inserted in their proper position in the following arrangement.

## I. Native Elements

#### I. Non-Metals

1. Carbon Group. Hardness 10 and 1-2

1. Diamond. C. Isometric, tetrahedral crystals small, bright and translucent:-

striated octahedron (fig.), white, (in tube). Accompanied I+ by specimen of peridotite or "blue earth." 2.50

cube. 3.00 29

30 dodecahedron, slightly rounded, gray 2.00

tetrahexahedron f, nearly spherical. 3.00 4

triangular, modified. 3.00 5

6 twins, tw. pl. octahedron o, (fig.). 6.00

octahedral cleavage or "splint." 1.00 7

Bort, rough, gray. 1.00 8\*

Carbonado, granular, black, small. 5.00 99 z. Diamond

Cliftonite (meteoric). Minute cubes, black.

10\* 2. Graphite, Plumbago, Black Lead. C. Rhombohedral, thin hexagonal tables, black. .30

radio-foliate globular concretions. .75 II

12+ foliated mass. .40

fine granular. .40 13

16

earthy, impure. .30 149

Graphitoid, Schungite. C. Massive,

combustible.

## 2. Sulphur Group. Range of Hardness 1.5-2.5

3. Sulphur. S. Orthorhombic. Perfect crystals. translucent fine yellow:-

pyramids  $\phi$ , s, brachydome n and base c 15+ (fig.), very brilliant. .75

crystal, doubly terminated, sharp. .75



34 Type Species No. No.	COMPLETE TYPE COLLECTION. DANA'S SYSTEM Sulphur—Continued
170	acute pyramid p, prominent (fig.), bright.
18 19°	obtuse pyramidal, well defined. 1.50 tabular, base predominating, transparent,
20º 2I	large. 2.00 sphenoidal (fig.), distinct. 1.50 elongated in parallel growth, transparent, very 17. Sulphur
22	brilliant, loose. 1.00 drusy crystalline50
23 24	dark brownish-yellow75 massive35
	encrusting lava35 Arsensulfurite.
	uisqueite, chiefly C and S. Amorphous, black.
	bic (?), crust, reddish.
	rsenschwefel. As <sub>3</sub> S <sub>3</sub> +H <sub>2</sub> O. Tetragonal (?). Blue-gray.
5. Se	elenium. Occurrence in nature very doubtful.

#### II. Semi-Metals

- 3. Tellurium-Arsenic Group. Rhombohedral
  Range of Hardness 2—3.5
- 6. Selen-Tellurium. Te containing Se. Massive, indistinct-ly columnar, blackish-gray.
- 7. Tellurium. Te. Rhombohedral, minute hexagonal prisms.
- prismatic cleavage, tin-white. 1.00 granular, tin-white. 1.00
- 28 8. Arsenic. As. Rhombohedral, spherical aggregates of small rhombs. .75
- 29+ fine granular, tin-white, tarnishing. .75
- 30° reniform. 1.00

Arsenolamprite. Nearly pure As. Cleavable, brilliant, lead-gray.

- 31\* 9. Allemontite. As containing Sb. Rhombohedral, crystalline granular reniform, tin-white, tarnishing. 3.00 Antimonial arsenic. 17As+1Sb. Crystalline.
  - 10. Antimony. Sb. Containing sometimes Ag, Fe or As. Rhombohedral, crystals.

Туре	TELLURIUM—ARSENIC AND GOLD GROUPS 35 Species Antimony—Continued No.
No. 32+	No. crystalline granular, tin-white. 2.50
ა≃ : 33°	cleavage. 3.00
	compactly fibrous. 2.50
34 35	11. Bismuth. Bi with occasional traces of As, etc. Rhombo-
<b>3</b> 3	hedral, twinned arborescent, tarnished. 5.00
36°	foliated cleavage reddish-silver-white, coated with molyb-
Ju-	denite. 2.00
37+	crystalline disseminated. 1.00
37 · 38	crystalline granular. 1.00
30	12. Zinc. Zn. Rhombohedral. Existence in nature requires
	confirmation.
III	. Metals
4	. Gold Group. Isometric. Range of Hardness 2.5—3.5
•	(Lead 1.5)
	(Lead 1.5)
39	13. Gold. Au usually alloyed with Ag. Isometric, minute
	cubes, gold-yellow. 3.00
40º	octahedron $o$ , minute, hollowed faces (fig.). 3.00
4 I	arborescent. 7.00
42 <sup>9</sup>	dodecahedron d, microscopic. 2.50
43	trisoctahedron, minute. 6.00
44	filiform. 2.00
45	spongiform, crystalline. 3.00
46*	quartz, disseminated masses. 2.50
47°	quartz, disseminated plates. 2.00
48+	quartz, disseminated grains. 2.00
49+	nugget. I.50
50+	nattened grains or "dust." 1.50
51+	Electrum (argentiferous), elongated crystals, pale gold-
	yellow. 2.00
5 <b>2</b> º	Electrum, "leaf gold," flattened parallel to octahedron o,
	with triangular markings. 2.00
53	Porpezite (palladium gold). 5.00
	Rhodite (rhodium gold), doubtful.

54° 14. Silver. Ag with some Au, Cu, etc. Isometric, small elongated octahedrons, silver-white, tarnishing. 2.00

Bismuth gold, "Black Gold" (tarnished).

Gold Amalgam, 57 to 61 p.c.Hg.

36 Type	COMPLETE TYPE COLLECTION. DANA'S SYSTEM Species Silver—Continued No.
55	dodecahedrons, small. 3.00
56	arborescent grouping. 6.00
57	cruciform parallel grouping of elongated crystals, coated with smaltite. 7.00
58	filiform, wire silver, coarse. 2.00
59+	filiform, fine, matted. 1.50
60*	disseminated grains. 1.00
610	disseminated scales. 2.00
62+	leaf silver, plates. 1.50
63	filmy coating. I.00
64	massive. 2.00
•	Küstelite (auriferous).
	cupriferous, associated with native copper.
650	antimonial with smaltite. 3.00
66	15. Copper. Cu often containing Ag Bi, d
	etc. Isometric, dodecahedron
	prominent (fig.), small, copper-
_	red75
67	tetrahexahedrons, small, distinct.
<b>.</b>	1.00
680	complex twins75
69	paraner groupings. 1.00
70+	crystallized, arborescent (fig.)50
71 72	elongated dodecahedron. 1.00 filiform (wire)75
72 73°	plates or "leaf copper." .50
74+	massive75
75 <sup>+</sup>	disseminated in conglomerate20
76°	disseminated in limpid calcite. 1.00
77°	altering to cuprite. 1.00
78+	16. Mercury, Quicksilver. Hg with sometimes
	a little Ag. Liquid, minute globules,
	brilliant tin-white, in shale. 1.00
79°	ditto, in cinnadar. 1.00
8 <b>0</b> *	17. Amalgam. Ag containing Hg. Isometric, small dodeca- hedron, silver-white. 4.00


hedron, silver-white. 4.00
81 Arquerite, small octahedron. 9.00

Arquerite, massive. 3.00 Kongsbergite, crystallized.

82

Type Species

- 18. Lead. Nearly pure Pb. Isometric, minute crystals.
- 83+ plate, lead-gray. 1.00 filmy coating on polyadelphite. dendritic.
  - 19. Tin. Nearly pure Sn. Crystalline grains, grayish-white.
  - 5. Platinum-Iron Group. Range of Hardness 4-7
- 84+ 20. Platinum. Pt alloyed with Fe, Ir, etc. Isometric, Non-magnetic, minute grains and scales, steel-gray, with gold. 1.50
- 85 Magnetic, grains. 1.50
- 86e Magnetic, showing polarity, nugget. 4.00
- 87 black, grains. 2.00
- 88 21. Iridium. Ir with Pt. Isometric, minute cubes, yellowish tin-white, loose. 2.50
- 89° angular grains. 2.00
- 90 22. Iridosmine. Ir with Os. Rhombohedral. Nevyanskite. Over 40 p.c. Ir. Minute, hexagonal prisms a with pyramid x and base c (fig.). 3.00



- 91+ irregular flattened grains, tin-white. 2.00 Siserskite. Not over 30 p.c. Ir. Steel-gray.
  - 23. Palladium. Pd alloyed with Pt and Ir. Isometric, minute octahedrons, whitish steel-gray.
  - 24. Allopalladium. Pd. Rhombohedral, minute six-sided tables, grayish silver-white.
  - 25. Iron. Normally about 82—92 p.c. Fe with Ni, Co, etc. Isometric.
- 92+ I. Terrestrial, nearly pure, dark steel-gray, rusting. 1.00
- 93 Awaruite, nickeliferous, grains, steel-gray. 3.00
- Josephinite, Fe<sub>2</sub>Ni<sub>5</sub>, massive pebbles, gray. 1.00 Siderazot, Fe<sub>5</sub>N<sub>2</sub>, coating on lava.
- 95 II. Meteoric, Siderite (iron), diamondiferous mass. 3.00
- 96+ Meteoric, Siderite (iron) crystalline, etched plate, showing Widmanstätten figures. 2.00
- 97º Meteoric, Siderolite (iron and stone). 2.50
- 98º Meteoric, Aerolite (stone), gray, with black crust. 1.50

38 Type Sp No. 1	COMPLETE TYPE COLLECTION. DANA'S SYS	TEM
No. 1	Meteoric, altered to limonite-magnetite shale	. 1.00
	Iron Compounds from Meteoric Iron	18
100	Edmonsonite. Fe—Ni alloy. Chalypite. Fe with 7 to 11 p.c. C. Cohenite. (Fe,Ni,Co),C. Isometric (?), d tals, tin-white becoming bronze-yellow Schreibersite. (Fe,Ni),P. Graphic steel-gra meteoric iron, etched plate. 6.00 Rhabdite. Fe, Ni phosphide. Tetragonal, n	y crystals, in
II.	Sulphides, Selenides, Tell Arsenides, Antimonide	•
I Su	alphides, Selenides, Tellurides of the Se	emi-Metals
1.	Realgar Group. RS. Monoclinic. Hardn	ess 1·5—2
1019 2	26. Realgar. AsS. Monoclinic, small crystal, dolomite75	light red, on
102	group of large prismatic crystals. 2.00	
1039	drusy crystals in crystalline mass, dark i	ed. 1.25
104	microscopic crystals incrusting lava75	
105+	compact, light red. 1.00	
2.	Stibnite Group. R <sub>2</sub> S <sub>3</sub> . Orthorhombic. H (Guanajuatite 2·5—3·5)	ardness 2
106	27. I. Orpiment. As <sub>2</sub> S <sub>3</sub> . Monoclinic, crystals,	vellow. 3.00
1079	foliated cleavage, canary-yellow. 1.00	^
108+	crystalline mass, lemon-yellow. 1.00	
1090	reniform, greenish-yellow. 1.25	$\int p/p$
110	globular, radiated structure, with realgar.  1.50	4
	28. Stibnite. Antimony Glance. Sb <sub>2</sub> S <sub>3</sub> . Ortho- rhombic, large prism, deeply furrowed, splendent lead-gray. 2.00	m m è
112	slender prism, acutely terminated50	1 1 11
113*	acicular crystals (fig.), radiating group.	II3. Stibuite
	1.00	and presente

Type Species

1330

No. No.
114 crystal, bent50
115 globular, radiated structure75
116+ crystalline, columnar bladed35
117° crystalline, granular35
1180 partially oxidized crystals, yellow. 1.00
Metastibnite. Sb <sub>2</sub> S <sub>3</sub> . Amorphous, red.
119 29. Bismuthinite. Bismuth Glance. Bi <sub>2</sub> S <sub>3</sub> . Orthorhombic,
acicular, lead-gray. 3.00
120º foliated cleavage. 1.00
121+ bladed cleavage. 1.00
S. seleniferous, 4Bi <sub>2</sub> S <sub>3</sub> , Bi <sub>2</sub> Se <sub>3</sub> , prisms.
122+ 30. Guanajuatite. Bi <sub>2</sub> Se <sub>3</sub> . Orthorhombic, acicular prisms.
2.00
123 compact, 2.00
Range of Hardness 1.5-2
124* 31. Tetradymite. Bi and Te. Rhombohedral, foliated, pale
steel-gray. 1.50
125° sulphurous, small acute rhombs, striated. 2.00
126 sulphurous, crystalline granular. 1.50
I. Grünlingite. Bi <sub>4</sub> TeS <sub>3</sub> . Rhombohedral (?), massive, gray,
tarnishing.
32. Josëite. Bi and Te with some S and Se. Laminated,
perfect cleavage, grayish.
33. Wehrlite. Bi and Te with some S and Ag. Foliated, per-
fect cleavage, grayish-white.
3. Molybdenite Group. RS <sub>2</sub> . Hardness 1—1.5
127 34. Molybdenite. MoS <sub>2</sub> . Hexagonal, tapering prism, bluish-
lead-gray. 1.00
128+ tabular hexagon40
129 <sup>®</sup> hexagonal cleavages, loose40
130 large foliated cleavage75
131 disseminated scales40
leaves in prism of rock crystal. 1.00
The second secon

granular. .40
II. Patronite. VS<sub>4</sub>(?). Amorphous, black.

# II. Sulphides, Selenides, Tellurides, Arsenides, Antimonides of the Metals

## A. Basic Division. Dyscrasite Group. Range of Hardness 3—4.5

Type Species No. No. 134 35. Dyscrasite. Ag<sub>3</sub>Sb. Orthorhombic, prisms (altering). 8.00 crystalline, coarse foliated, silver-white, tarnishing. 2.50 crystalline, fine granular.

36. Horsfordite. Cu<sub>6</sub>Sb.(?). Massive, silver-white, tarnishing.

136 II. Keweenawite. (Cu,Ni,Co)<sub>2</sub>As. Massive, pale pinkishbrown. 5.00

37. Domeykite. Cu, As. Reniform.

137+ compact, gray, tarnishing iridescent-bronze. 1.25

138e argentiferous, granular. 2.50

II. Stibiodomeykite, contains some Sb.

Orileyite. (Cu<sub>2</sub>Fe)<sub>3</sub>(AsSb)<sub>2</sub>(?). Massive, purplish steel-gray.

II. Ledouxite. Cu<sub>4</sub>As. Massive, silver-white.

II. Mohawkite. (Cu, Ni, Co), As. Massive. 6.00

139° 38. Algodonite. Cu<sub>6</sub>As. Massive granular, silver-white, tarnishing bronze. 3.00

140 39. Whitneyite. Cu<sub>9</sub>As. Massive, reddish-white, tarnishing. 5.00

- 40. Chilenite. Perhaps Ag<sub>6</sub>Bi. Amorphous, silver-white, tarnishing.
- 41. STÜTZITE. Perhaps Ag<sub>4</sub>Te. Hexagonal (?), highly modified, reddish lead-gray.

#### B. Monosulphides, Selenides, Tellurides, Etc.

- 1. Galena Group. RS. Isometric, holohedral Range of Hardness 2—3
- 141• 42. Argentite, Silver Glance. Ag<sub>2</sub>S. Isometric, small distinct cube, blackish-lead-gray. 2.00

octahedron o. 2.50

143 dodecahedron d, small, well defined. 2.00

144 arborescent. 2.50

145\* distorted crystal, small. 1.50

146• massive. 2.00

Argentite-Continued disseminated. 1.25 147 +coating. 1.25 148 Jalpaite. 3Ag<sub>2</sub>S.Cu<sub>2</sub>S. 149+ 43. Hessite. Ag<sub>2</sub>Te. Isometric, small highly modified crystal, dark gray. 2.50 150 massive. 2.50 151+ 44. Petzite. (Ag, Au), Te. Massive, iron-gray, tarnishing. 3.00 152+ 45. Galena, Galenite. PbS. Isometric, cube, large, well defined, lead-gray. .75 1539 cubo-octahedrons, ideal symmetry. .75 octahedron, large, perfect. 1.00 154\* dodecahedron d modifying octahedron o and cube a, 1550 large. 1.25 1560 contact-twins. 1.25 1579 penetration-twins (fig.), definite. 1.50 reticulated. 1.00 158 hollow crystals. 1.00 159 1600 deeply eroded. .75 elongated crystals. .75 1619 thin tabular crystals, minute. .50 162 163 minute cubes on lava. 1.00 157. Galena cleavage, broad. .40 164+ fibrous. .75 165 argentiferous, coarse granular. .75 166+ 1679 fine granular. .40 cryptocrystalline. .60 168 Steinmannite, contains As and Sb, crystallized. 1.50 169 altered to cerussite. 1.00 170 171 altered to leadhillite. 1.50 Huascolite. Nearly PbS.1½ZnS. Granular. 172 Cuproplumbite. Cu<sub>2</sub>S.2PbS. Massive. 2.50 173º 46. Altaite. PbTe. Isometric, disseminated, vellowish tinwhite, tarnishing. 2.00 174º 47. Clausthalite. PbSe. Isometric, crystalline disseminated, lead-gray. 2.00 Tilkerodite (cobaltiferous).

Isometric, skeleton dodecahe-175 S. I. Aguilarite. Ag<sub>2</sub>S.Ag<sub>2</sub>Se. drons, iron-black. 6.00

48. Naumannite. (Ag<sub>2</sub>Pb)Se. Isometric, cubes.

42	COMPLETE	TYPE COLLECTION.	DANA'S	SYSTEM
Type No.	Species No.	Naumannite—Continued	i	

1760 massive, iron-black. 4.00

177° 49. Berzelianite. Cu<sub>2</sub>Se. Finely disseminated, silver-white, tarnishing. 1.50

S. Umangite. CuSe.Cu<sub>2</sub>Se. Massive, tarnishes violet-blue.

178° 50. Lehrbachite. PbSe with HgSe. Massive granular, dark gray. 3.00

179 51. Eucairite. Cu<sub>2</sub>Se. Ag<sub>2</sub>Se. Isometric, massive, silvery leadgray. 9.00

180° 52. Zorgite. Cu and Pb selenide (varying). Massive granular, lead-gray, tarnishing. 3.00

181° 53. Crookesite. (Cu,Tl,Ag)<sub>2</sub>Se. Finely disseminated, lead-gray. 7.00

#### 2. Chalcocite Group. RS. Orthorhombic.

Range of Hardness 1.5—3

182 54. Chalcocite, Copper Glance. Cu₂S. Orthorhombic, Redruthite, prismatic, blackish-lead-gray, tarnishing. 2.00

183° Redruthite, twins, pseudo-hexagonal (fig.), perfect, sharp. 1.50

twins, cruciform. 3.00

185° massive, granular. 1.00

186+ compact. 1.00

187 II. Chalmersite. Cu<sub>2</sub>S.Fe<sub>4</sub>S<sub>3</sub>. Orthorhombic 183. Chalcoctte prisms, bronze-yellow. 6.00

188 55. Stromeyerite. (Ag,Cu)<sub>2</sub>S. Orthorhombic, crystallized. 8.00 189° massive, dark steel-gray. 2.50


190° 56. Sternbergite. AgFe<sub>2</sub>S<sub>3</sub>. Orthorhombic, twins, thin tabular pseudo-hexagonal, dark brown. 3.00 Frieseite, Ag<sub>2</sub>Fe<sub>5</sub>S<sub>8</sub>, twins, thick tabular.

Argentopyrite. Ag, Fe sulphide. Orthorhombic, pseudohexagonal prismatic twin, bronze-yellow, tarnished.

Hardness 2—2.5

191° 57. Acanthite. Ag<sub>2</sub>S. Orthorhombic, acicular, iron-black. 2.00 Daleminzite. Ag<sub>2</sub>S. Orthorhombic, short prisms (pseudomorphous after stephanite?).

II. Rickardite. Cu<sub>4</sub>Te<sub>3</sub>. Massive, purple.



# 3. Sphalerite Group. RS. Isometric, tetrahedral. Range of Hardness 2:5-2:5

Type Species	Range of Hardness 2.5—3.5
	phalerite, Zinc Blende. ZnS. Iso-
	metric, tetrahedron modified
	by cube (fig.). 1.50
1930	bright cube with + and — tetra-
_	hedrons prominent, alternately
	splendent and dull black, sharp
	ideal symmetry. 1.00
194*	trisoctahedron m and dodecahedron 192. Sphalerite
	d, rounded into obtuse cone
	(fig.), transparent brown,
	brilliant50
1950	brownish crystals on chert50
196+	Ruby Blende, bright transparent.
1970	·75 greenish-yellow penetration-twins,
1970	· adamantine, transparent. 1.00
198+	distorted crystals, grouped, "Black 194. Sphalerite
-,-	Jack," glistening50
1990	hemitrope twins, contact    to octa-
	hedron o (fig.), splendent black,
	sharp. 1.00
200	twins, contact 1 to octahedron o, clear
	brown75
2010	thin tabular transparent twin, on
	granular dolomite. 1.00
202	indescent crystais. 1.25
203*	cleavage dodecahedron, opaque75
204	ditto dodecahedron, clear greenish. 1.50
205	ditto, perfect, splendent clear yellowish. 1.50
206+	coarse granular, cleavable, resinous brown20
207	fine granular, gray50
208	fibrous40
209	Cleiophane, pure white. 1.00
2100	Schalenblende, compact, reniform, interstratified with galena, grayish60
011	Marmatite, 16 p.c. Fe, twins, splendent black. 1.50
211	maimante, 10 p.c. re, twins, spiendent biack. 1.50

Type Species Sphalerite—Continued No. No.
Christophite (18 p.c. Fe), mixture of fine to coarse gran-
ular, black40
213 cadmiferous, Pribramite75
mercurial.
stanniferous.
214 indiferous. 1.00
215 59. Metacinnabarite. HgS. Isometric, tetrahedral, small
twins, tw. pl. octahedron o, iron-black. 2.50
216º minute globular druses. 1.00
217° massive disseminated. 1.00
Guadalcazarite, zinc-metacinnabarite. Rhombohedral(?).
218 60. Tiemannite. HgSe. Isometric, tetrahedral, small, highly
modified crystals. 6.00
219• massive, dark gray. 2.50
61. Onofrite. Hg(S,Se). Massive granular, blackish gray.
220 62. Coloradoite. HgTe. Massive granular, iron-black. 9.00
221 63. Alabandite. MnS. Isometric, tetrahedral, small crystals.
3.00
cleavable-granular. 1.00
massive, iron-black, tarnishing. 1.00
Hardness 4
64. Oldhamite. Meteoric. CaS. Isometric, small spherules,
clear pale brown.
224+ 65. Pentlandite. (FeNi)S. Isometric, octahedral cleavage.
light bronze-yellow. 2.00
I. Gunnarite. Fe <sub>3</sub> Ni <sub>2</sub> S <sub>8</sub> (?). Yellowish tin-white tarnishing.
* <u> </u>
4. Cinnabar—Wurtzite—Millerite Group.
Rhombohedral or Hexagonal.
·
Cinnabar Series. Hardness 2
225 66. Cinnabar. HgS. Rhombohedral, trapezohedral, small
rhombic, adamantine, fine red. 2.00
226° tabular, bright. 1.50
penetration-twins, complete ideal symmetry, small, loose.
(6)75
penetration-twins, tw. axis c, dull. 9.00
229+ acicular prisms. 1.25
230 radiating crystalline. 1.50

....

-

. ...

Type S <sub>I</sub>	CINNABAR—WURTZITE—MILLERITE GROUP 45 Cinnabar—Continued
No. 231	no. drusy incrustation. 1.00
2329	fine granular-massive. 2.50
233*	coarse granular disseminated, cochineal-red75
234	earthy coating75
2350	hepatic (containing Idrialite), dark brown. 1.50
	67. Covellite, Indigo Copper. CuS. Hexagonal, or rhombo-
•	hedral, small deeply striated thin hexagonal tables,
	ideal symmetry, indigo-blue. 3.00
237	very thin flexible leaves, disseminated. 2.00
238+	foliated, crystalline, fine indigo-blue. 2.00
239	compact. 2.50
240	platiniferous, enclosing sperrylite, porous. 2.00
<b>24</b> I	coating. I.00
	Wurtzite Series. Range of Hardness 2-3
2420	68. Greenockite. CdS. Hexagonal, hemimorphic, prism with several pyramids, minute, distinct, clear resinous
	yellow. 4.00
243+	coating on sphalerite. 1.50
244	coloring smithsonite. 1.50
245	69. Wurtzite. ZnS. Hexagonal, hemimorphic minute pyramidal. 2.00
246	Schalenblende, fibrous, brownish-black. 1.25
247+	Schalenblende, massive. 1.00
	Erythrozincite. A manganese-wurtzite(?). Thin plates, translucent red, in lapis-lazuli.
	Millerite Series. Range of Hardness 3.5-5.5
248°	70. Millerite. NiS. Rhombohedral, acicular, brass to
	bronze-yellow. 2.00
249	radiating capillary tufts. 1.50
250	capillary matted coating. 1.00
251	fine capillary, in clear calcite. 1.00
252+	fibrous plates, semi-globular. 1.00
253°	71. Niccolite, Arsenical Nickel. NiAs. Hexagonal, prism and
054	pyramids, reddish-gray, tarnishing. 3.00
254	reniform, columnar. 2.00 reticulated.

46 COMPLETE TYPE COLLECTION. DANA'S SYSTEM  Type Species Niccolite—Continued No. No.
255+ massive. 1.00
256 Antimonial, massive. 1.00
257° 72. Breithauptite. NiSb. Hexagonal, thin tabular. 2.50
258 arborescent, massive, violet copper-red. 1.50
259 73. Troilite. FeS (meteoric). Massive, brown. 1.50
260° 74. Pyrrhotite, Magnetic Pyrites. Fe <sub>11</sub> S <sub>12</sub> , containing sometimes
5 p.c.Ni. Hexagonal, small tabular, sharp. 1.50
261 thick tabular. 1.50
acute pyramidal.
262° compact, bronze-yellow, tarnishing20
263+ granular-massive, nickeliferous20
C. Intermediate Division
Group 1. Range of Hardness 3—4.5
264° 75. Polydymite. Ni <sub>4</sub> S <sub>5</sub> (?). Isometric, imperfect cubic cleave
age, steel-gray, tarnishing. 3.00
I. Hauchecornite. (Ni,Co), (S,Bi,Sb) <sub>8</sub> . Tetragonal, bronze
yellow.
S. Sychnodymite. (Co, Cu) S <sub>5</sub> . Isometric, octahedral, steel-gray
76. Beyrichite. Ni <sub>3</sub> S <sub>4</sub> (?). Prismatic, lead-gray.
265 77. Melonite. Ni <sub>2</sub> Te <sub>3</sub> (?). Hexagonal, basal cleavages, dis
seminated, reddish-white. 9.00
0 0 0 0 0 0
Group 2. Range of Hardness 3—5.5
266° 78. II. Bornite, Erubescite. 5Cu <sub>2</sub> S.Fe <sub>2</sub> S <sub>3</sub> . Isometric, penetra-
tion-twins, rounded. 2.50
fine granular, argentiferous, coppery bluish-brown, tar-
nishing75
268 compact, iridescent75
269+ 79. Linnæite. Co <sub>3</sub> S <sub>4</sub> . Isometric, octahedron o, small, idea
symmetry, splendent. 2.00
270° massive, pale steel-gray, tarnishing. 1.50
Siegenite (niccoliferous), octahedral. 2.00
272 80. Daubreelite. FeS.Cr <sub>2</sub> S <sub>3</sub> . Massive, in meteoric iron, bril
liant black. 8.00
273 81. Cubanite. CuFe <sub>2</sub> S <sub>4</sub> . Isometric, massive, bronze-yellow. 6.00
Chalcopyrrhotite. Fe <sub>4</sub> CuS <sub>6</sub> . Massive.
82. Carrollite. CuS.Co <sub>2</sub> S <sub>3</sub> . Isometric, steel-gray, faintly red

MILLERITE SERIES AND PYRITE GROUP Type Species No. No. 274+ 83. Chalcopyrite, Copper Pyrites. CuFeS, varving. Tetragonal. noidal, small ideal sphenoids (fig.), brass-yellow, tarnishing, on pearl-spar. .50 ditto, iridescent, parallel grouping 275 on sphalerite. 1.50 sphenoids, elongated. 1.00 274. Chalcopyrite 276 octahedroid, + and - sphenoids, 2779 complete. 2.00 scalenohedron and sphenoid (fig.). 278 contact-twin || to sphenoid, perfect (fig.). 279\* 1.00 280° penetration-twin. 2819 hollow sphenoids. 2.00 compact. .35 282+ 283 fine granular. .35 2849 reniform. 1.25 278. Chalcopyrite

285° 84. II. Stannite, Tin Pyrites. Cu<sub>2</sub>S.FeS.SnS<sub>2</sub>. Scalenohedral, tetragonal symmetry, twins (pseudo-tetrahedral), brilliant,

perfect, minute. 3.00 ·


286 twins, complex, dull. 3.00

288

massive, fine granular, olive-steel-287+ gray, tarnishing. .75

massive, coarse, greenish-iron-black, tarnishing.


II. Teallite. PbS.SnS<sub>2</sub>. Orthorhombic, foliated, blackish gray.



#### D. Disulphides, Diarsenides, etc.

- 1. Pyrite Group. RS<sub>2</sub>, RAs<sub>2</sub>, RSb<sub>2</sub>. Isometric, pyritohedral. Range of Hardness 4-6.5 (Laurite 7.5)
  - 85. Pyrite, Iron Pyrites. FeS<sub>2</sub>. Isometric, pyritohedral, crystals of ideal symmetry, splendent pale brassvellow:-

pyritohedron (fig.), large, loose. 289+ cube a, striated. (fig.). .50 290+



45 Type Species No. No.	COMPLETE TYPE COLLECTION. DANA'S SYSTEM  Pyrite—Continued
291	cube on coal shale50
292+	octahedron o, sharp50
293°	diploids, small. 1.00
294*	pyritohedron e modifying octahedron o
	(fig.), large. 1.00
<b>2</b> 95°	cube modifying pyritohedron (fig.)75 290. Pyrite
296	cube a modifying octahedron o50
297	octahedron o modifying pyritohedron e (fig.). 1.00
2989	octahedron modifying cube50
299●	diploid modifying cube75
300	diploid modifying octahedron, rounded.
301●	trapezohedron n modifying octahedron.
-	I.00 294. Pyrite
	dodecahedron $d$ modifying cube $a$ .
302	highly modified, rounded, striated and
	pitted. 1.00
303●	penetration-twins, pyritohedrons. tw.pl.  normal to dodecahedron d (fig.), loose. (3)75
304*	oscillatory combination (striated) of cube and pyritohedron, group, metallicadamantime. 1.00
305	elongated octahedron, loose. 5.00
306	acicular elongated cube small. 2.00
307°	distorted, saddle-shaped cube, loose30
308•	flat disk, radiated crystallized. 1.00
309°	nodule of crystals30
310	globular50
311	stalactitic, radiated75
312	drusy, iridescent30
313	fine granular20
314+	compact20
315°	auriferous (2½ oz. Au. to ton)
II.	coarse50 Bravoite, highly nickeliferous.
316+	altered to limonite, cube, large, dull brown, loose50
317	ditto, pyritohedron, complete50 303. Pyrite

<b>.</b>	PYRITE GROUP	49
Type S <sub>l</sub> No.	pecies Pyrite—Continued No.	
318	diploid and octahedron, complete (2)75	
3190	86. Hauerite. MnS <sub>2</sub> . Isometric, octahedron, complete i	deal
	symmetry, loose, brownish-black. 1.00	
320*	cubo-octahedron, loose. 1.00	
321	globular drusy, in clay. 1.00	
3229	87. Smaltite, Arsenical Cobalt. CoAs <sub>2</sub> . Isometric, cubo-oct	ahe-
	dral, small, steel-gray, tarnishing. 1.50	
323	twins, complex. 2.00	
324 <del>9</del>	reticulated. 2.50	
325+	massive. 1.00	
326	argentiferous. 2.00	
1	NOTE.—Smaltite and Chloanthite graduate chemically into each other.	
327	88. Chloanthite. NiAs <sub>2</sub> . Isometric, small cubo-octahedr	ons,
•	steel-gray. 2.50	
328*	massive. I.25	
	89. Cobaltite, Cobalt Glance. CoS2.CoAs2. Isometric, o	om-
	plete, highly symmetrical bright, pale reddish-	gray
	crystals loose:—	
329*	pyritohedron e (3). 1.00	
330	cube $a$ , perfect (3). 1.25	
3319	octahedron o, sharp. 1.25	
3320	pyritohedron e and octahedron o	۸.
	(fig.), modifying cube. 1.00	1
333	cubo-octahedron, sharp (3). 1.00	u
334+	crystalline, fine granular, altering to	ا م
	erythrite60	- /
335	massive60	V
	Ferrocobaltite (28 p.c. Fe), plumose.	//
336	90. Gersdorffite, Nickel Glance. NiS <sub>2</sub> . 332. Cobaltite	_
	NiAs <sub>2</sub> . Isometric, small, perfect	
	cubo-octahedrons, bright steel-	
	gray, tarnishing. 4.00	
337+	massive, granular. 1.50	
338	91. Corynite. Ni(As,Sb)S. Isometric, octahedron, silv	ery.
	gray. 9.00	
339	92. Ullmannite. NiS <sub>2</sub> . NiSb <sub>2</sub> . Isometric, cube, steel-gray.	3.00
340 <del>0</del>	massive granular, steel-gray. 1.00	
<b>341</b>	I. Willyamite. CoS <sub>2</sub> . NiS <sub>2</sub> . CoSb <sub>4</sub> . NiSb <sub>2</sub> . Isometric, c	ubic
	cleavage, steel-gray. 8.00	
	S. Kallilite. NiS <sub>2</sub> . NiBi <sub>2</sub> . Massive, bluish-gray.	

50 Type S No.	
342+	93. Sperrylite. PtAs <sub>2</sub> . Isometric, microscopic distinct cubes,
	brilliant tin-white, loose (lot). 3.00
	94. Laurite. RuS <sub>2</sub> . (and osmium 3.03 p. c.). Isometric, octa- hedrons, minute, dark iron-black.
343	95. Skutterudite. CoAs <sub>3</sub> . Isometric, modified octahedron,
	pale lead-gray. 8.00
	Nickel-skutterudite, RAs, with R=Ni:Co:Fe=4:2:1,
	granular, gray.
	I. Bismutosmaltite, contains Bi.
	2. Marcasite Group. RS <sub>2</sub> , RAs <sub>2</sub> , Etc. Orthorhombic. Range of Hardness 4.5—6.5
Th	ese species closely parallel those of the preceding Pyrite Group.
	· · · · · · · · · · · · · · · · · · ·
344°	96. Marcasite. FeS <sub>2</sub> . Orthorhombic,
	prism $m$ and base $c$ predominating (fig.), pale bronze-
	vellow 100
	yellow. 1.00 344. Marcasite
2454	2 244 Marie Commen
345*	octahedroid, macro- and brachy-
345*	octahedroid, macro- and brachy- domes e and l prominent
	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. 1.00
345* 346 347+	octahedroid, macro- and brachy- domes e and l prominent
346	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00
346	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00 Cockscomb Pyrites, (fig.), aggrega-
346 347+	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. 1.00
346 347 <sup>+</sup> 348° 349	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. 1.00 capillary in calcite50
346 347+ 348°	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. 1.00 capillary in calcite50 hepatic, massive, brown40
346 347 <sup>+</sup> 348° 349	octahedroid, macro- and brachydomes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. 1.00 capillary in calcite50 hepatic, massive, brown40 cellular by disappearance of incrusted minerals50
346 347+ 348° 349 350°	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. 1.00 capillary in calcite50 hepatic, massive, brown40 cellular by disappearance of incrusted minerals50 stalactitic, radial structure, exterior
346 347+ 348° 349 350° 351	octahedroid, macro- and brachydomes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. 1.00 capillary in calcite50 hepatic, massive, brown40 cellular by disappearance of incrusted minerals50
346 347+ 348° 349 350° 351	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. 1.00 capillary in calcite50 hepatic, massive, brown40 cellular by disappearance of incrusted minerals50 stalactitic, radial structure, exterior
346 347+ 348° 349 350° 351 352° 353 354	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. I.00 radiated. I.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. I.00 capillary in calcite50 hepatic, massive, brown40 cellular by disappearance of incrusted minerals50 stalactitic, radial structure, exterior well crystallized. I.00 arborescent, crystallized75 globular50
346 347+ 348° 349 350° 351 352° 353 354	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. I.00 radiated. I.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. I.00 capillary in calcite50 hepatic, massive, brown40 cellular by disappearance of incrusted minerals50 stalactitic, radial structure, exterior well crystallized. I.00 arborescent, crystallized75 globular50  97. Löllingite. FeAs <sub>2</sub> . Orthorhom-
346 347+ 348° 349 350° 351 352° 353 354	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. 1.00 radiated. 1.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. 1.00 capillary in calcite50 hepatic, massive, brown40 cellular by disappearance of incrusted minerals50 stalactitic, radial structure, exterior well crystallized. 1.00 arborescent, crystallized75 globular50  97. Löllingite. FeAs <sub>2</sub> . Orthorhombic, small, sharply de-
346 347+ 348° 349 350° 351 352° 353 354	octahedroid, macro- and brachy- domes e and l prominent (fig.), small, distinct. I.00 radiated. I.00 Cockscomb Pyrites, (fig.), aggregations of flattened twins75 Spear Pyrites, twins, symmetrical. I.00 capillary in calcite50 hepatic, massive, brown40 cellular by disappearance of incrusted minerals50 stalactitic, radial structure, exterior well crystallized. I.00 arborescent, crystallized75 globular50  97. Löllingite. FeAs <sub>2</sub> . Orthorhom-

twins, cruciform. 1.00

massive, silver-gray. .35

356

357\*

Type Species	Löllingite—Continued
358e	Leucopyrite, Fe <sub>3</sub> As <sub>4</sub> , massive35
	Geyerite, 6.73 p.c. S.
359	Glaucopyrite, cobaltiferous75
	Pacite. Fe sulph-arsenide. Massive.
	Arsenopyrite, Arsenical Pyrites or Mispickel. FeAsS.
-	Orthorhombic, unit prism m and brachydome u, well
	defined (fig.), silver-gray. 1.00
361 <b>0</b>	penetration-twins, tw. pl.    to macro-
	dome e, sharp75
362*	repeated-twins, brilliant. 1.25
	cruciform-twins.
363+	granular massive, silver-gray25 360. Areenopyrite
364	compact35
	Danaite (cobaltiferous), brilliant crystals.
	nickeliferous.
	Safflorite. CoAs <sub>2</sub> . Orthorhombic, prism and macrodome.
365°	massive, tin-white, tarnishing. 2.00
	Badenite. (Co,Ni,Fe) <sub>2</sub> (As,Bi) <sub>3</sub> . Massive, steel-gray.
	Rammelsbergite. NiAs <sub>2</sub> . Orthorhombic, crystals.
366°	massive, reddish tin-white. 1.00
367 <b>101</b> . 0	Glaucodot. (CoFe)AsS. Orthorhombic, prism and two
	brachydomes, perfect, grayish-tin-white, loose. 1.25
368◎	prism and brachydome, minute. 1.00
369*	twins, loose, sharp, bright. 1.25
370	crystalline, disseminated. 1.00
371 <b>0102.</b> A	Moclasite. Co(As, Bi)S. Orthorhombic, columnar aggre-
	gates, steel-gray. 2.50
103. <b>T</b>	Wolfachite. Ni(As,Sb)S(?). Orthorhombic, prism and brac-

#### 3. Sylvanite Group. Range of Hardness 1-2

372+104. I., II. Sylvanite, Goldschmidtite. Au<sub>2</sub>AgTe<sub>6</sub>. Monoclinic, small prisms, brilliant silver-white. 3.00

373 skeleton crystals, steel-gray. 2.50

374° "Graphic Tellurium," arborescent twinning. 2.00

hydome, minute, tin-white.

II. Von Diestite. Ag, Bi telluride. Massive.

375 105. I. Krennerite. AuTe<sub>2</sub>. Orthorhombic, small prisms, brilliant pale brass-yellow. 5.00

376+ I. Calaverite. AuTe<sub>2</sub>. Triclinic(?), pale bronze-yellow. 4.00

52 COMPLETE TYPE COLLECTION. DANA'S SYSTEM

Type Species

377 106. Nagyagite. Au<sub>2</sub>Pb<sub>14</sub>Sb<sub>3</sub>Te<sub>7</sub>S<sub>17</sub>. Orthorhombic, small tables, blackish-lead-gray. 4.00

378+ Foliated Tellurium, crystalline foliæ, embedded. 3.00 I. Kalgoorlite. HgAu<sub>2</sub>Ag<sub>6</sub>Te<sub>6</sub>. Massive, iron-black.

#### Oxysulphides. Hardness 1-1.5 and 4-4.5

379 107. Kermesite. 2Sb<sub>2</sub>S<sub>3</sub>.Sb<sub>2</sub>O<sub>3</sub>. Monoclinic, capillary tufts, cherry-red. 3.00

380° radio-fibrous, crystalline. 2.50

108. Voltzite. 4ZnS.ZnO. Globules, curved lamellar structure.

#### Appendix to Sulphides, Etc.

Bolivianite. Antimonial Ag sulphide. Orthorhombic. acicular rhombic prisms, lead-gray.

Plumbostannite. Antimonial Sn, Pb, Fe sulphide. Granular, gray.

## III. Sulpho-Salts

The metals chiefly present as bases are copper, silver and lead, also iron, zinc and mercury, rarely nickel, cobalt, etc.

#### I. Sulpharsenites, Sulphantimonites, Etc.

#### A. Acidic Division. Range of Hardness 2-3.5

109. Livingstonite. HgS.2Sb<sub>2</sub>S<sub>3</sub>. Prismatic crystals.

columnar massive, blackish-lead-gray. 2.50

- 382 110. Guejarite. Cu<sub>2</sub>S.2Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, prisms flattened parallel to brachypinacoid b, bluish-steel-gray.
  - 111. Chiviatite. 2PbS.3Bi<sub>2</sub>S<sub>3</sub>. Foliated massive, lead-gray.
  - 112. Cuprobismutite. 3Cu<sub>2</sub>S.4Bi<sub>2</sub>S<sub>3</sub>. Slender prisms, bluishblack.

Dognacskaite. Bi, Cu sulphide. Cleavages, gray.

113. Rezbanyite. 4PbS.5Bi<sub>2</sub>S<sub>3</sub>. Massive, lead-gray, darkening.

#### B. Meta Division

Zinkenite Group. RS.(As,Sb,Bi)<sub>2</sub>S<sub>3</sub>. Orthorhombic. Range of Hardness 2—4

114. Zinkenite. PbS.Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, crystals. 383 capillary, matted, steel-gray. 2.50

Type Species No. No.

Zinkenite-Continued

384+

fibrous, massive. 2.00

- I. Andorite. 2PbS.Ag<sub>2</sub>S.<sub>3</sub>Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, highly modi-385 fied flat prisms, metallic-adamantine. 7.00
- 386° massive, dark steel-gray. 4.00
- 387 115. Sartorite. PbS.As<sub>2</sub>S<sub>3</sub>. Orthorhombic, slender prisms, dark lead-gray. 5.00
- II. Hutchinsonite. (Tl,Ag,Cu)<sub>2</sub>S.As<sub>2</sub>S<sub>3</sub>+PbS.As<sub>2</sub>S<sub>3</sub>(?). Ortho-388 rhombic, flattened prisms, adamantine, red. 9.00
- 3890116. Emplectite. Cu<sub>2</sub>S.Bi<sub>2</sub>S<sub>3</sub>. Orthorhombic, thin prisms, grayish. 1.25
  - II. Histrixite. 7Bi<sub>2</sub>S<sub>3</sub>.2Sb<sub>2</sub>S<sub>3</sub>.5CuFeS<sub>2</sub>. Orthorhombic, prismatic, steel-gray.
- 390 117. I. Chalcostibite. Cu<sub>2</sub>S.Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, small furrowed flat prism, perfect, iron-gray. 9.00
- 391°118. Galenobismutite. PbS.Bi<sub>2</sub>S<sub>3</sub>. Columnar crystalline, light lead-gray. 2.50
- argentiferous, Alaskaite, compact. 3.00 392 seleniferous.
  - 119. Berthierite. FeS.Sb<sub>2</sub>S<sub>3</sub>. Elongated prisms.
- fibrous massive, dark steel-gray. 1.00 393+ granular massive.
  - II. Trechmannite. Ag<sub>2</sub>S.As<sub>2</sub>S<sub>3</sub>. Rhombohedral, small prisms, bright red.
  - II. Smithite. Ag<sub>2</sub>S.As<sub>2</sub>S<sub>3</sub>. Monoclinic, pyramidal, adamantine, light red.
  - 120. Matildite. Ag<sub>2</sub>S.Bi<sub>2</sub>S<sub>3</sub>. Slender prisms, gray. Plenargyrite. Ag<sub>2</sub>S.Bi<sub>2</sub>S<sub>3</sub>(?). Crystalline, black.
- 394° I. Lorandite. Tl<sub>2</sub>S.As<sub>2</sub>S<sub>3</sub>. Monoclinic, highly modified prism, dark red, on realgar. 2.50
- 395 121. Miargyrite. Ag<sub>2</sub>S.Sb<sub>2</sub>S<sub>3</sub>. Monoclinic, thick tabular, brilliant, dark steel-gray. 6.00

#### C. Intermediate Division. Hardness 2.5

- 396 122. Plagionite. 5PbS.4Sb<sub>2</sub>S<sub>3</sub> (?). Monoclinic, thick tabular, drusy. 3.00
- 397€ massive, fine granular, blackish lead-gray. 2.00
- 398 II. Baumhauerite. 4PbS.3As<sub>2</sub>S<sub>3</sub>. Monoclinic, adamantine, lead-gray. 4.00

- II. Liveingite. 5PbS.4As<sub>2</sub>S<sub>3</sub>. Monoclinic.
- 123. I. Binnite. Formerly regarded as a distinct species. Now classed as a variety of tennantite, No. 149.
- 124. Klaprotholite. 3Cu<sub>2</sub>S.2Bi<sub>2</sub>S<sub>3</sub>. Orthorhombic, furrowed prisms, steel-gray, tarnishing.
- 125. Schirmerite. 3(Ag<sub>2</sub>, Pb)S.2Bi<sub>2</sub>S<sub>3</sub>. Massive, lead-gray.
- 126. Warrenite. 3PbS.2Sb<sub>2</sub>S<sub>3</sub>. Acicular, matted, grayish-black.

#### Jamesonite Group. 2RS.(As,Sb,Bi)<sub>2</sub>S<sub>3</sub>.

#### Range of Hardness 2-3.5

- 399°127. II. Dufrenoysite. 2PbS.As<sub>2</sub>S<sub>3</sub>. Monoclinic, flat prism, blackish lead-gray. 2.00
- 400 I. Rathite. S23.72, As 17.24, Sb 4.53, Pb 52.98. Orthorhombic, prismatic, black. 7.00
  - 128. Cosalite. 2PbS.Bi<sub>2</sub>S<sub>3</sub>. Orthorhombic, prismatic || macrodome e.
- 401° crystalline radio-fibrous, steel-gray. 2.00
  - 129. Schapbachite. PbS.Ag<sub>2</sub>S.Bi<sub>2</sub>S<sub>3</sub>. Orthorhombic (?), minute needles, lead-gray.

massive, fine granular.

- 402 130. Jamesonite. 2PbS.Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, acicular. 1.50
- 403° capillary, matted. 1.25
- 404+ crystalline granular, steel-gray. 1.00
- fibrous, lead-gray. 1.00
- 406° plumose, "Feather Ore." 1.00
- 407 compact. 1.25
- 408 131. Kobellite. 2PbS.(Bi,Sb)<sub>2</sub>S<sub>3</sub>. Massive, steel-gray. 8.00
  - 132. Brongniardite. PbS.Ag<sub>2</sub>S.Sb<sub>2</sub>S<sub>3</sub>. Isometric, octahedron o truncated by dodecahedron d, grayish-black.

#### -Hardness 2.5

- 409 133. Semseyite. 7PbS.3Sb<sub>2</sub>S<sub>3</sub>(?). Monoclinic, small distinct tables in rosette-like aggregates, dark gray. 9.00
- 410°134. Diaphorite. 5(Pb,Ag<sub>2</sub>)S.2Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, small prisms, splendent dark steel-gray. 2.50
- 411\*135. Freieslebenite. 5(Pb,Ag<sub>2</sub>)S.2Sb<sub>2</sub>S<sub>3</sub>. Monoclinic, prismatic, blackish lead-gray. 2.50

#### D. Ortho Division

Bournonite Group. 3RS. (As,Sb,Bi)<sub>2</sub>S<sub>3</sub>. Orthorhombic.

Range of Hardness 2.5-4

Type Species No. No. 4120136. Bournonite. 3(Pb,Cu<sub>2</sub>)S.Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, tabular (fig.), splendent blackishgray. 2.00 Bournonite 412. prismatic, perfect. 1.50 413 prismatic, parallel aggregate. 1.25 4140 repeated twins, "Wheel Ore" 415+ (fig.). 1.50 massive, fine granular. 1.25 416 II. Seligmannite. Cu<sub>2</sub>S.2PbS.As<sub>2</sub>S<sub>3</sub>(?). Orthorhombic, small crys-415. Bournonite tals, lead-gray.

137. Wittichenite. 3Cu<sub>2</sub>S.Bi<sub>2</sub>S<sub>3</sub>. Orthorhombic, crystals like bournonite.

417 massive disseminated, tin-white, tarnishing. .300
418°138. Aikinite. 3(Pb,Cu<sub>2</sub>)S.Bi<sub>2</sub>S<sub>3</sub>. Orthorhombic, acicular in quartz, blackish lead-gray, tarnishing reddish. 3.00

139. I. Boulangerite. 5PbS.2Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, prisms.

crystalline plumose, lead-gray. 1.00 compact, with stibnite. .75

140. Lillianite. 3PbS.Bi<sub>2</sub>S<sub>3</sub>. Massive, crystalline, steel-gray.

141. Stylotypite. 3(Cu<sub>2</sub>,Ag<sub>2</sub>,Fe)S.Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, columnar prisms, iron-black.

Dürfeldtite. Chiefly Pb,Ag,Mn sulphantimonite. Indistinctly fibrous, light gray.

S. Falkenhaynite. 3Cu<sub>2</sub>S.Sb<sub>2</sub>S<sub>3</sub>. Massive, gray-black.

421-142. Guitermanite. 10PbS.3As<sub>2</sub>S<sub>3</sub>. Massive compact, bluishgray. 1.00

422 II. Lengenbachite. 6PbS(Ag,Cu)<sub>2</sub>S.2As<sub>2</sub>S<sub>3</sub>(?). Triclinic(?), bladed, steel-gray. 4.00

143. Tapalpite.  $_3Ag_2(S,Te).Bi_2(S,Te)_3$  (?). Massive granular, steel-gray, tarnishing.

#### Pyrargyrite Group. 3Ag<sub>2</sub>S.(As,Sb)<sub>2</sub>S<sub>3</sub>.

Rhombohedral, hemimorphic. Range of Hardness 2-2.5 Species Type

423°144. Pyrargyrite, Antimonial or Dark Ruby Silver. 3Ag<sub>2</sub>S.Sb<sub>2</sub>S<sub>3</sub>. Rhombohedral, six-sided prism (fig.), reddish-black, deep red by transmitted light. 2.00

rhombohedron prominent, perfect. 2.00

425 twins. 2.50

426+ compact massive. 1.00

427 disseminated. 1.00

428°145. Proustite, Arsenical or Light Ruby Silver. 3Ag<sub>2</sub>S.As<sub>2</sub>S<sub>3</sub>. Rhombohedral, small acute rhomb, splendent vermillion. 3.00

429 scalenohedron, translucent, small, but distinct. 2.50

423. Pyrargyrite •

430° twins, perfect. 3.00

431+ massive, compact, dark red. 1.25

fine granular, disseminated. 1.25

S. Sanuginite. Ag sulpharsenite. Hexagonal, fine glittering scales, black, dark red by transmitted light.

433 146. Pyrostilpnite. 3Ag<sub>2</sub>S.Sb<sub>2</sub>S<sub>3</sub>. Monoclinic, minute prisms, hyacinth-red. 4.00


147. Rittingerite. As, Se, and 57.7 p.c. Ag. Monoclinic, minute tables, iron-black.

#### E. Basic Division

Tetrahedrite Group. 4RS.(Sb,As)<sub>2</sub>S<sub>3</sub>. Isometric, tetrahedral. Hardness 3—4.5

148. Tetrahedrite, Gray Copper or Fahlerz. 4Cu<sub>2</sub>S.Sb<sub>2</sub>S<sub>3</sub>. Isometric, small crystals of ideal symmetry, splendent iron-black:—

434+ tetrahedrons modified by trigonal tristetrahedron n, and dodecahedron d (fig.), on quartz crystals. 1.00



-Hardness 3 and 2.5

445°150. S. Jordanite. 4PbS.As<sub>2</sub>S<sub>3</sub>. Monoclinic, tetrahedral, sixsided, base c predominating. 2.50

446°151. Meneghinite. 4PbS.Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, tetrahedral, acicular, splendent blackish-lead-gray, loose. 1.00

-Hardness 2.5

447°152. Geocronite, Kilbrickenite. 5PbS.Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic. tetrahedral, massive. 1.50

448°153. Stephanite, Brittle Silver. 5Ag<sub>2</sub>S.Sb<sub>2</sub>S<sub>3</sub>. Orthorhombic, tetrahedral, hemimorphic, small tables, base predominating (fig.). 3.00



twins, hexagonal tables, splendent 449+ iron-black. 2.00

450 twins, hexagonal prismatic, bright. 2.50

massive, compact. 2.00 45I

disseminated. 2.00 4529

Type Species No. No.

435°

436

437°

438

439<sup>®</sup> 440+

44 I

443

4449

–Hardness 2—3

154. II. Kilbrickenite. Formerly regarded as a distinct species. It is identical with geocronite, No. 152.

COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type Species No. No.
155. Beegerite. 6PbS.Bi <sub>2</sub> S <sub>3</sub> . Isometric(?), brilliant indistinct
crystals, gray.
Richmondite. 6RS.Sb <sub>2</sub> S <sub>3</sub> with R=Cu <sub>2</sub> ,Fe,Ag <sub>2</sub> Zn(?).
156 I. Polybasite. 9Ag <sub>2</sub> S.Sb <sub>2</sub> S <sub>3</sub> . Monoclinic, ideal pseudo-
hexagonal crystals, splendent iron-black (in thin
splinters, cherry-red):—
repeated twins, tabular, tw.pl. unit prism m. 2.00
ditto, prisms. 2.50
455° massive, disseminated. 2.00
I. Pearceite. 9Ag <sub>2</sub> S.As <sub>2</sub> S <sub>3</sub> . Monoclinic, pseudo-rhombohe-
dral tables, black.
Hardness 2·5
157. Polyargyrite. 12Ag <sub>2</sub> S.Sb <sub>2</sub> S <sub>3</sub> . Isometric, distorted cubo-
octahedrons, indistinct, iron-black.
II C 1.1
II. Sulpharsenates, Sulphantimonates, Etc.
Enargite Group. Hardness 3 and 3.5
456°158. Enargite. 3Cu <sub>2</sub> S.As <sub>2</sub> S <sub>5</sub> . Orthorhombic, unit prism m,
macropinacoid $a$ , and base $c$ , symmetrical. 2.00
twins, iron-black.
457° bladed-columnar cleavage. 1.50
458+ granular-cleavable, grayish-black. 1.00
459 massive, grayish-black. 1.00
I. Lautite. CuAsS.
Clarite. 3Cu <sub>2</sub> S.As <sub>2</sub> S <sub>3</sub> . Monoclinic, dark lead-gray.
NOTE—Enargite and Famatinite graduate chemically toward each other.
159. Famatinite. 3Cu <sub>2</sub> S.Sb <sub>2</sub> S <sub>5</sub> . Orthorhombic, isomorphous
with enargite, gray with tinge of copper-red.
460° massive. 2.50
Hardness 2
4610 II. Sulvanite. 3Cu <sub>2</sub> S.V <sub>2</sub> S <sub>5</sub> . Massive, bronze-yellow, tarnish-
ing. 1.00
462 160. I. Xanthoconite. Ag <sub>3</sub> AsS <sub>3</sub> . Monoclinic, thin tabular,
reddish-yellow. 2.50
reniform mass, granular structure.
463 161. Epiboulangerite. 3PbS.Sb <sub>2</sub> S <sub>5</sub> . Orthorhombic (?), prismetic readles bluich block.
matic needles, bluish-black. 2.00
Hardness 3.5
162. Epigenite. 4Cu <sub>2</sub> S.3FeS.As <sub>2</sub> S <sub>5</sub> (?). Orthorhombic, short prisms with macro- and brachydome, steel-gray.

#### -Hardness 2.5

- I. Canfieldite.  $4Ag_2S(SnGe)S_2$ . Isometric, tetrahedral(?), octahedrons o, with dodecahedrons d, bluish-black. Regnolite.  $5CuS.FeS.ZnS.As_2S_3$ . Isometric, tetrahedral.
- 464° I. Franckeite. 5PbS.Sb<sub>2</sub>S<sub>3</sub>.2SnS<sub>2</sub>. Massive, imperfectly radiofoliate, blackish-gray. 1.50
- 465° I. Cylindrite, Kylindrite. 6PbS.Sb<sub>2</sub>S<sub>3</sub>.6SnS<sub>2</sub>. Massive, cylindrical-foliated structure, blackish-lead-gray. 1.25
- 466 163. I. Argyrodite. 4Ag<sub>2</sub>S.GeS<sub>2</sub>. Isometric, tetrahedral, dodecahedron d and tetrahedron o, loose. 9.00
- penetration-twins, small distinct. 6.00
- 4689 drusy crystals, reniform grouping. 4.00
- 469\* compact massive, dark steel-gray. 2.50

# IV. Haloids.—Chlorides, Bromides, Iodides; Fluorides

## I Anhydrous Chlorides, Bromides, Iodides; Fluorides.

Calomel Group. R<sub>2</sub>Cl<sub>2</sub>. Range of Hardness 1—2

- 470\*164. Calomel. Hg<sub>2</sub>Cl<sub>2</sub>. Tetragonal, highly complex. 4.00 471 twins, gray. 4.00
  - II. Kleinite. Hg, NH<sub>4</sub> chloride(?). Hexagonal, short prisms, orange-yellow.

Mercuric chloride, HgCl<sub>2</sub>(?).

- 472°165. Nantokite. Cu<sub>2</sub>Cl<sub>2</sub>. Isometric, granular, white. 2.00
- 473 I. Marshite. Cu<sub>2</sub>I<sub>2</sub>. Isometric, tetrahedral, minute truncated tetrahedrons, ideal symmetry, adamantine, reddish-brown. 6.00
  - I. Miersite. Ag<sub>2</sub>I<sub>2</sub>. Isometric, tetrahedral, cubes with tetrahedral faces, adamantine, yellow.


## Halite Group. RCl, etc. Isometric.

Chlorides, etc., of sodium, potassium, ammonium and silver.

- 474°166. Halite, Rock Salt. NaCl. Isometric, cubes, perfect, white. .75
- cubo-octahedrons, symmetrical. .50
- 476° octahedrons, clear, loose, (3). .50
- 477+ elongated cubo-octahedrons, clear, loose (3). .50

ล
Ш
Ш
Ш
W
<b>/</b>
al,
,
th
00
ıs,
<b>W</b> -
a-

HALITE AND FLUORITE GROUPS Type Species 505 173. Iodyrite. Agl. Kraus and Cook, A.J.S. 27, 210, 1909. Only four of the ten small but distinctly formed types now in stock and described by these authors, are here cata-Hexagonal, hemimorphic, diametral prism a and base c,c'predominating, truncated by unit prism m and unit pyramid u,u', apparently holohedral, minute, symmetrical, lemon-yellow, on psilomelane. 8.00 hemimorphic, unit prism m, base c, unit 506 pyramid i' all prominent with several truncating unit pyramids, loose (12). 1.00 parallel grouping, barrel-shaped, of sev-507\* eral similar individuals (c,i,m,i) (fig.), loose (12).



507. Iodyrite Kraus and Cook


5080 contact-twins, tw.pl. pyramid e (fig.), loose (12). 1.00

crystalline granular, sulphur-yel-509+ low. 2.00

thin plates with lamellar struct-510 ure. 2.00

massive. 2.00 511

Tocornalite. Ag, Hg iodide. Granular, yellow.



Kraus and Cook

I. Cuproiodargyrite. CuI.AgI. Incrustation, sulphur-yellow.

#### Fluorite Group. $R(Cl,F)_2$ . Isometric.

Range of Hardness, Fluorite 4 Chloromagnesite series very soft, except Sellaite 5, Tysonite 4.5-5, Cryolite series 2.5-3.5

174. Hydrophilite. CaCl<sub>2</sub>. Isometric, cubic, encrusting.

175. Fluorite, Fluor Spar. CaF<sub>2</sub>. Isometric, bright crystals of sharp ideal symmetry, transparent to translucent:-



512. Fluorite

512+ cubic, blue (fig.). .50

62 COMPLETE TYPE COLLECTION. DANA'S SYSTE  Type Species Fluorite—Continued No. No.	M
513° cubic, large, yellow. 1.00	$\wedge$
514 cubic, splendent iridescent, brown. 1.50	
515° cubic, dark green75	
516 cubic, elongated, complete, opaque grayish-	0/0/
violet. 1.00	
517* octahedron o (fig.), pale green. 1.00	517. Fluorite
octahedron, rose-pink, loose. 1.50	
octahedron, small, colorless, on lava. 1.00	
520° dodecahedron, small. 1.50	
hexoctahedron t, and cube a (fig.), dull translucent. 1.25	a a
522° octahedron modifying cube, splendent75	(V)
523* tetrahexahedron e modifying cube a, "fluoroid" (fig.), small. 1.00	711/
dodecahedron $d$ modifying cube $a$ ,	Fluorite
525 trigonal trisoctahedron modifying octahedron. 1.25	
526* penetration-twins, tw.pl. octahedron o (fig.), vicinal striations. 1.00	
527° pseudo-octahedron, formed by parallel grouping of dodecahedrons, capped by dodecahedron. 1.00	523. Fluorite
528 ditto, grouping of modified cubes capped by modified cube, on white altered barite, sea-green. 1.25	
529 ditto, emerald-green on pink rhodo- chrosite. 2.00	
530 capping of blue flat cube on green modified octahedron. 2.00	
531+ cleavage octahedron, emerald- green50	
532 cleavage, tetrahedral symmetry, pink50 526.	Fluorite
533° banded crystal, loose. 1.00	
534 columnar-granular banded, polished, transluc John." 1.00	ent "Blue-
535+ cleavable-granular, coarse, greenish-white2	<b>:</b> 0
536 crystalline granular, fine30	

#### FLUORITE GROUP

Tuna Saad	FLUORITE GROUP  Fluorite—Continued	63	
Type Speci- No. No			
537	massive, compact50		
	Antozonite, odor attributed to free Fl.		
538°	Chlorophane, fluorescent. 1.50		
539 <sup>e</sup>	encrusted with milky quartz75		
540		.00	
5 <b>4</b> I	Capped Fluor, primary growth (cube), secondary cr		
	of quartz, with capping of fluor and siderite. I	.50	
542 <sup>9</sup>	altered to blue chalcedony. 1.25		
176	. Chloromagnesite. MgCl <sub>2</sub> . A deliquescence on lava.		
	. Sellaite. MgF <sub>2</sub> . Tetragonal, prismatic, clear colorless. 6		
178	. Lawrencite. FeCl <sub>2</sub> . Solid, on exposure exuding from i	ron	
	as minute drops, finally oxidizing, brown. 2.00		
179	. Scacchite. MnCl <sub>2</sub> . A deliquescent mass.		
	Chloralluminite. AlCl <sub>3</sub> + $x$ H <sub>2</sub> O. In lava.		
5440180	. Cotunnite. PbCl <sub>2</sub> . Orthorhombic, acicular. 1.50		
Pseudocotunnite. PbCl <sub>2</sub> .KCl (?). Acicular, yellow.			
181	. Molysite. FeCl <sub>3</sub> . Incrusting.		
182	. Tysonite. (Ce,La,Di)F <sub>3</sub> . Hexagonal, thick prisms.		
545	massive, yellowish. 3.00		
5460183	. Cryolite. 3NaF.AIF <sub>3</sub> . Monoclinic,		
	short modified square prisms	_	
	with cubic aspect (fig.),	<u>d</u>	
	grouped parallel, transpar-	"	
	ent colorless, sharp. 1.50	1.5	
547+	massive, translucent, white30 $\left  \left  \left  \left  \left  \right  \right ^{a} \right  \right $	<i>//</i>	
548	massive, brownish40		
549	massive, with galena, chalcopy-	/	

Elpasolite. K, Na, Al fluoride. II. Cryolithionite. 3LiF.3NaF.2AlF<sub>3</sub>. Isometric, dodecahedrons, transparent colorless.

546. Cryolite

550 184. Chiolite. 5NaF.3AlF<sub>3</sub>. Tetragonal, pyramidal, white. 2.00 185. Hieratite. 2KF.SiF<sub>4</sub>. Isometric. Hydrofluorite. HF. Gas (volcanic).

Cryptohalite. 2NH<sub>4</sub>F.SiF<sub>4</sub>(?). Volcanic. Proidonite. SiF<sub>4</sub>. Volcanic.

rite and siderite. 1.00

#### II. Oxychlorides, Oxyfluorides

#### A. Oxychlorides.

Type Species Range of Hardness 2.5—3.5

551°186. Matlockite. PbCl<sub>2</sub>. PbO. Tetragonal, tabular || to c. 2.00 552 rosette-like group, yellowish. 3.00

- 553°187. Mendipite. PbCl<sub>2</sub>.2PbO. Orthorhombic, columnar mass, white. 3.00
- 554 188. Schwartzembergite. Pb(I,Cl)<sub>2</sub>,2PbO(?). Rhombohedral, incrusting, yellow. 4.00
  - I. Penfieldite. PbO.2PbCl<sub>2</sub>. Hexagonal, prismatic, white.
- 555 II. Terlinguaite. Hg<sub>2</sub>ClO. Monoclinic, small crystals, adamantine sulphur-yellow. 9.00
- 556 II. Egglestonite. Hg<sub>2</sub>O.2HgCl. Isometric, minute dodecahedrons, brownish-yellow. 8.00
- 557°189. Laurionite. PbCl<sub>2</sub>.Pb(OH<sub>2</sub>). Orthorhombic, small flat prisms, adamantine, clear colorless. Formed by action of sea-water on ancient slag. 1.00
  - I., II. Paralaurionite. PbCl<sub>2</sub>. Pb(OH)<sub>2</sub>. Monoclinic, prisms, white.
    - 190. Daviesite. Lead oxychloride. Orthorhombic, minute prisms, colorless.
    - 191. I. Fiedlerite. Contains Pb and Cl. Monoclinic, minute tables, colorless, transparent.
- 558 192. I. Percylite. Pb(OH)Cl.Cu(OH)Cl. Isometric, cubes, minute, sky-blue. 4.00
- 559°S. I. Cumengéite. Pb(OH)Cl.Cu(OH)Cl. Tetragonal(?), small sharp octahedroids, indigo-blue, loose. 1.00
- 560\*S. I. Boléite. Pb(OH)Cl.Cu(OH)Cl.+½AgCl. Isometric(?), cubic habit, ideal symmetry, indigo-blue, loose. .75
- 561+193. Atacamite. CuCl<sub>2</sub>.3Cu(OH)<sub>2</sub>. Orthorhombic, acicular, transparent emerald-green. 1.50
- short thick prisms (fig.), sharp, brilliant. 2.50 octahedroid, well defined, bright. 2.50
- 564 crystal aggregate, bright emerald-green. 2.00
- 565° granular massive. 1.00

  II. Paratacamite. CuCl<sub>2</sub>.3Cu(OH)<sub>2</sub>. Rhombohe-<sup>562. Atacamite</sup>

  dral, bright green.



Tallingite. Cu<sub>5</sub>(OH)<sub>8</sub>Cl<sub>2</sub>+4H<sub>2</sub>O. Thin crusts of minute globules, greenish-blue.

S. Footeite. 8Cu(OH)<sub>2</sub>.CuCl<sub>2</sub>+4H<sub>2</sub>O. Monoclinic, minute prisms, deep blue.

Melanothallite. CuCl<sub>2</sub>.CuO.2H<sub>2</sub>O (?). Volcanic.

566 Erythrocalcite. CuCl<sub>2</sub>(H<sub>2</sub>O undetermined). 5.00 Atelite. 2CuO.CuCl<sub>2</sub>.3H<sub>2</sub>O. An altered tenorite. Green, volcanic.

194. Daubréeite. 2Bi<sub>2</sub>O<sub>3</sub>.BiCl<sub>3</sub>.3H<sub>2</sub>O (?). Amorphous, whitish.

II. Koenenite. Al, Mg oxychloride. Rhombohedral, crusts, red.

#### B. Oxyfluorides. Hardness of Fluocerite 4

567°195. Nocerite. 2(Ca,Mg)F<sub>2</sub>.(Ca,Mg)O(?). Hexagonal, acicular, white. 1.00

568•196. Fluocerite. R<sub>2</sub>O<sub>3</sub>.4RF<sub>3</sub>, where R=Ce metals chiefly. Massive, reddish-yellow. 1.50

## III. Hydrous Chlorides, etc.

#### A. Hydrous Chlorides. Very Soft

569°197. Bischofite. MgCl<sub>2</sub>+6H<sub>2</sub>O. Crystalline, clear colorless. .50 198. Kremersite. KCl.NH<sub>4</sub>Cl.FeCl<sub>3</sub>+H<sub>2</sub>O. Isometric, octa-

hedrons, ruby-red.

199. Erythrosiderite. 2KCl.FeCl<sub>3</sub>.H<sub>2</sub>O. Orthorhombic, red.

200. Douglasite. 2KCl.FeCl<sub>2</sub>.2H<sub>2</sub>O(?).

201. Carnallite. KCl.MgCl<sub>2</sub>+6H<sub>2</sub>O. Orthorhombic, pseudohexagonal pyramids.

570 massive granular, white. .30

571+ massive granular, reddish. .30

202. Tachhydrite. CaCl<sub>2</sub>.2MgCl<sub>2</sub>+12H<sub>2</sub>O. Rhombohedral, cleavages.

572\* massive, yellowish. .30

#### B. Hydrous Fluorides

203. Fluellite. AIF<sub>3</sub>+H<sub>2</sub>O. Orthorhombic, white.

573 204. Prosopite. CaF<sub>2</sub>.2Al(F,OH)<sub>3</sub>. Monoclinic (or triclinic), embedded crystals, whitish. 5.00

574°205. Pachnolite. NaF.CaF<sub>2</sub>.AlF<sub>3</sub>.H<sub>2</sub>O. Monoclinic, prism and acute pyramid, distinct, minute, clear colorless. 1.00

575°206. Thomsenolite. NaF.CaF<sub>2</sub>.AlF<sub>3</sub>.H<sub>2</sub>O. Monoclinic, prism and base, cubic symmetry. 1.50

66	COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type No.	Species Thomsenolite—Continued No.
576	prismatic, transparent colorless. 1.50
577	massive, white. 1.00
578	Hagemannite. Impure thomsenolite?, jaspery, yellow50
579°	207. Gearksutite. CaF <sub>2</sub> .Al(F,OH) <sub>3</sub> H <sub>2</sub> O. White earthy masses
	of minute colorless needles. 1.00
580°	208. Ralstonite. (Na <sub>2</sub> ,Mg)F <sub>2</sub> .3Al(F,OH) <sub>3</sub> 2H <sub>2</sub> O. Isometric octahedrons, whitish. 2.50
	209. Yttrocerite. $2(2RF_3.9CaF_2).+3H_2O$ , with R=Ce(La,Di): $Y(Er)=1:2$ . Massive, white, in quartz.
5819	violet-blue, in mica75

## V. Oxides

## I. Oxides of Silicon. Hardness 7 (Opal 5.5—6.5)

210. Quartz. SiO<sub>2</sub>. Rhombohedral. Rare and unimportant types are omitted.

Note—The term "pyramid" (six-sided), is here used interchangeably with "two rhombohedrons r and s."

#### A. PHENOCRYSTALLINE OR VITREOUS VARIETIES

Crystals sharply defined, brilliant and transparent.

Rock Crystal colorless forms and types follow:—

	Rock Crystal, colorless forms and types follow:—
582+	prism $m$ and pyramid (two rhombohedrons $r$ and $z$ ),
	large, symmetrical, grouped50
583	ditto, very large, loose50
584+	ditto, but shorter (fig.), complete and quite
•	limpid, (so-called "diamonds"), loose   m m
	(6)50
585	ditto, on fine white marble75
586	ditto, acicular, group. 1.00 584. Quarts
587°	one rhombohedron r, and prism (fig.), loose50
588	one pyramidal plane only developed (rhombehodron r)
•	terminating very large slender prism, limpid. 1.00
589+	"quartzoid," double six-sided pyramid
0 2	(fig.), symmetrical, on hematite75
590	ditto, loose (6)50
5910	cuboid, r prominent (fig.), large, loose. $m \mid_{m} \mid_{m}$
J)-	1.00
59 <b>2</b>	chisel-edge termination, opposite rhombo-
U ) -	hedrons abnormally developed50 587. Quarts

_	
•	٠
o	ч

589. Quarts

#### Quartz-Continued Type Species trigonal pyramid s, modifying prism m, 593° rhombohedrons r and z (fig.), loose. .50 594 acute rhombohedron M, prism m, rhombohedrons r and s (fig.), tapering crystal. trigonal trapezohedron x, pyramid s, 595\* prism m, rhombohedrons r, z, large "right-handed crystal" (fig.). 1.00 ditto, large "left-handed crystal" (fig.). 596° highly modified, triangular etching, 597 (fig.), large. .75 598\* penetration-twin, tw. axis c (tw.pl. m), individuals right-handed 591. Quarts (shown in fig. by x), irregular dull and bright areas adjacent on r loose, large. 1.50 penetration-twin, irregular, Brazil law, 599 tw. pl. a (fig.). 3.00 6000 contact-twin, tw.pl. $\xi$ (1122), axes ccrossing at 84°33', a plane m coincident in both individuals (fig.), loose, large. 3.00 601 grouping simulating twin. 6029 distorted crystal (fig.), very large. 603 flattened || prism m, large. .75 bent crystal, large. .50 604 cavernous, angular cavities in faces, 6050 large, loose. .50 606 capped, large. 1.25 607° etching of trigonal pyramid s and only rhombohedron, pyramidal planes being alternately bright and dull. .50 drusy, large geode (two halves). .50 608\* drusy, three small unbroken geodes con-609

taining loose microscopic? crystals.

.50

**OXIDES OF SILICON** 



593. Quartz

68 Ty	COMI	PLETE TYPE COLLECTION. DANA'S SYST	EM
_		cleavage    to rhombohedron. 1.00	<b>^</b> -
		water-worn pebble, conchoidal fracture.	7
61	12 R	adiated crystalline, translucent	2
61		brous crystalline, translucent75	m m
		ar-quartz (asteriated), cut "en cabo- chon," clear colorless. 1.00	m / 2
61	15+ A1	nethyst, prisms, light violet50	A
61	16	pyramids in agate geode, dark purple precious. 3.00	595. Quartz
61	7	pyramids, druse on petrified wood. 1.00	$\wedge$
61	<b>189</b>	pyramids with surficial ferruginous in- clusions, surface red, translucent. 1.50	
61	19	dark rhombic "phantom" in light prisms terminated by one rhombohedron.  .75	m m
62	20*	dark pyramids terminating both ends	7
		of colorless prisms, loose75	596. Quarts
		parallel growth on smoky quartz. 1.00	1:1
62		"sceptre," purple quartzoid tipping ster of rock crystal, loose. 2.00	m-like prism
62	230	crystalline, banded with milky quartz, translucent50	
62	24	twinning shown by alternate rhom- bohedrons of amethyst and rock crystal in one large cross-section of prism, polished. 3.00	
62	25+ Ro	ose, translucent pink mass30	
62	26 Ro	ose, rich pink, iridescent internal re- flections (fractures), polished. 1.50	597. Quartz
62	27° Ro	ose, asteriated, transparent pink with purplish tinge, polished ball. 2.00	
62	28* Ye	ellow, Citrine, loose (6)50	KI TIL
	99 Sn	noky, light, Cairngorm Stone, loose, large50	2 m m
63		noky, dark translucent, very large, taper- ing (fig. 594)25	
63	31 Sn	noky, light, twisted flat parallel growth.	
		<b>4.</b> W	598. Quarts

Type Species No. No.	Quarts—Continued 69
No. No. 632	Greasy, sub-transparent mass30
633+	Milky, prism and pyramids, ideal sym-
-33	metry (fig.584), dull opaque, loose(3).
	.50
634	Milky, pyramids, translucent50 m
6359	Milky, translucent mass20
636	Siderite, Sapphire - quartz, translucent
	blue mass75
637*	Sagenitic (net-like), inclosing Rutile,
	acicular, "Flêches d'amour," pol-
640	ished. 2.50
638	Inclosing Rutile, capillary. 1.50 Inclusions of other acicular or capillary minerals:—
639	Black Tourmaline, capillary, in smoky
039	crystal, large50
640	ditto, with projecting "stem" or "core."
040	2.00
641* -	ditto, polished cross-section. 1.25
6429	Göthite, acicular, "Onegite." 1.50
643	Stibnite, acicular, in crystal. 2.00
6449	Asbestus, capillary, in crystal. 1.00
645	Actinolite acicular 1 00
646	Black Hornblende, acicular75
040	Epidote, acicular.
6479	Cat's-Eye, fibrous, gray, chatoy-
-47	ant50
	Tiger-Eye, see Crocidolite, al-
	tered
648°	Aventurine, Sunstone, red. 1.00
649	Aventurine, Sunstone, green,
.,	polished. 1.25 602. Quarts
	Impure from presence of other minerals densely distrib-
	uted:—
650*	Ferruginous, dull terra-cotta-red, pyramids75
651	Ferruginous, splendent brick-red druse. 1.50
6520	Ferruginous, dull ochre-yellow, mass of crystals75
653*	Chloritic, "phantom" crystal. 1.00
654	Chloritic, mossy, crystal75
	Actinolitic. Micaceous.
	Arenaceous.
655	containing albite crystals, in "phantom" form. 1.00
~ ~ ~ ~	

70 CC Type Species No. No.	OMPLETE TYPE COLLECTION. DANA'S SYSTEM  Quartz—Continued
656	containing anthracite, rock crystal50
657*	containing water and moving bubble, prism75
658	ditto, with moving anthracite, limpid crystal. 3.00
-	containing liquid CO <sub>2</sub> and moving bubble, which dis-
	appears on vaporizing liquid by warmth of hand,
	amethyst crystal.
٠.	B. CRYPTOCRYSTALLINE VARIETIES
659	Chalcedony, mammillary translucent white. 1.00
660+	mammillary40
66 I	botryoidal, brownish75
6620	stalactitic, tendon-color. 1.00
663	gray, polished75
664	geode, drusy lining50
665°	Enhydros, translucent geode containing water. 1.50
666°	Carnelian, translucent red, cut50
667	Sard, subtranslucent brownish-red, cut50
668*	Chrysoprase, translucent apple-green, precious. 1.00
669	Chrysoprase, translucent bluish-green. 1.00
6709	Chrysoprase, translucent turquoise-blue, precious. 1.50
6710	Prase, translucent dull leek-green75
672	Plasma, subtranslucent leek-green35
673*	Blood-stone, Heliotrope, subtranslucent leek-green with
	red spots50
	Agate:—
674+	Banded-agate, gray and white, polished75
675°	Banded-agate, red, in limestone75
676	Fortification-agate, brownish, polished. 1.25
6779	Eye-agate, concentric, polished. 1.50
678	Clouded-agate, polished75
679	Artificially colored agate. Long banded strip, cut into
	six cross-pieces, each colored differently and
	after joining, polished. 4.00
680°	Moss-agate, leek-green sea moss, polished75
681	ditto, with patches of chalcedony outlined by carne-
6000	lian. 1.25
6829	Moss-agate, Mocha-stone, large delicate branching, in clear chalcedony, cut. 1.50
683*	Dendritic-agate, black moss in gray ground. 1.00
684	Agatized-wood, brown, polished. 1.00
685	Onyx, straight banding, gray, polished. 1.50

72 Type Specie No. No.	COMPLETE TYPE COLLECTION. DANA'S SYSTEM Tridymite—Continued
719	polysynthetic twins in spherical rosettes, minute. 1.25
	Cristobalite. SiO <sub>2</sub> . Minute octahedrons, white.
720	Granuline. Identical with tridymite (?). Pulverulent, on lava, white. 1.00
721*	Melanophlogite. Contains SiO <sub>2</sub> , SO <sub>3</sub> , H <sub>2</sub> O. Pseudomor-
•	phous (?). Minute cubes75
	Sulfuricin. Contains SiO <sub>2</sub> , SO <sub>3</sub> , S, H <sub>2</sub> O.
	Porous.
212.	Opal. SiO <sub>2</sub> .nH <sub>2</sub> O. Amorphous.
722+	Precious Opal (i.e. with play of
	colors), milky. 1.00
723 <sup>9</sup>	in porphyry. 2.00
7249	in layers or zones. 1.00
725	bluish. 1.50
726*	greenish, in limonite. 1.50
727	reddish ground, "Harlequin
	opal." 2.00
728	in wood-opal. 1.00 718. Tridymite
729	replacing shell. 1.50
730+	Fire-opal, red, slightly irised75
731	Girasol, translucent bluish-white, reddish reflections. 1.00
	Common Opal, in part translucent:—
732+	Milk-opal, translucent40
733	Resin-opal (Wax-opal), opaque yellowish60
734°	Green-opal, translucent olive75
735°	Brick-red, Semiopal, opaque75
736	Hydrophane, translucency increases in water, adheres
	to the tongue, whitish. 1.50
737	Forcherite, orange-yellow. 1.50
7380	Cacholong, opaque whitish, adheres to the tongue. 1.00
739°	Opal-agate, variegated bands. 1.50
740°	Menilite, opaque concretion, dull grayish40
741*	Jasp-opal, opaque brownish-yellow40
742+	Wood-opal, petrified cellular, radial and concentric structure well marked, yellowish-brown40
743*	Hyalite, clear glassy, botryoidal60
744	Hyalite, translucent, whitish75
	Fiorite, Siliceous Sinter, includes:—
745 <b>°</b>	Pearl-sinter, stalactitic, pearly white. 1.50 Michaelite, pearly, capillary.

	GROUPS
Type Specie No. No.	s Opal—Continued
746°	Geyserite, porous concretions75
747	Geyserite, porous cauliflower-like. 1.25
748	Geyserite, massive75
749°	Float-stone, spongy structure, very light. 1.00 Tripolite includes:—
750+	Infusorial or Diatomaceous Earth (microscopic shells of Diatoms, etc.), fine chalky clay, snow-white30
75 I	ditto, gray30 Randannite, loose, mealy, white.
752	Tripoli Slate, laminated, impure30 Alumocalcite, very soft, milky.
753	Lussatite. Anhydrous (?) silica. Crystalline, translucent chalcedony-like globular crusts, whitish. 1.50
754	Tabasheer. Opaline silica deposited in joints of bam- boo, milk-white. 2.00

## II. Oxides of the Semi-Metals; also Mo and W

## 1. Arsenolite Group. R<sub>2</sub>O<sub>3</sub>. Isometric

Hardness 1.5 and 2.5

- 755 213. Arsenolite. As<sub>2</sub>O<sub>3</sub>. Isometric, octahedrons, white. 3.00 minute capillary, crusts.
- 756 214. Senarmontite. Sb<sub>2</sub>O<sub>3</sub>. Isometric, octahedrons, small, ideal symmetry, grayish. 2.50
- 757\* ditto, complete, loose (12). .75 granular massive.

## 2. Valentinite Group. R<sub>2</sub>O<sub>3</sub>. Hardness 2.5

- 758 215. Claudetite. As<sub>2</sub>O<sub>3</sub>. Monoclinic, very thin tabular || to b, small, flexible, pearly white. 6.00
- 759 216. Valentinite. Sb<sub>2</sub>O<sub>3</sub>. Orthorhombic, minute prisms, stellated druse, gray. 3.00
- 7600 tabular || to b, small fan-shaped aggregate. 4.00
- 761° crystalline stellated, yellow. 2.00 massive granular, white.
- 762°217. Bismite. Bi<sub>2</sub>O<sub>3</sub>. Orthorhombic (not crystallized in nature), pulverulent, straw-yellow. 2.00
  763 disseminated, greenish-yellow. 1.50

#### 3. Tellurite Group

Type Species RO<sub>2</sub>. Orthorhombic. Hardness, 2

764 218. Tellurite. TeO<sub>2</sub>. Orthorhombic, tufts of slender prisms, yellowish. 6.00

spherical masses, radiated structure.

## 4. Molybdite Group. Soft

- 765 219. II. Molybdite, Molybdic Ocher. Fe<sub>2</sub>O<sub>3</sub>.3 MoO<sub>3</sub>. 7½ H<sub>2</sub>O. Orthorhombic, minute capillary tufts, yellow. 2.00
- 766\* subfibrous disseminated, yellow. 1.00
- 767 pulverulent, yellowish-white. 1.00
- 768° Ilsemannite. MoO<sub>2</sub>.4MoO<sub>3</sub>. Crypto-crystalline, blackish, blue on exposure. 2.00
  - 220. Tungstite. WO<sub>3</sub>. Orthorhombic, earthy, yellow.
- 769 Meymacite. WO<sub>3.2</sub>H<sub>2</sub>O. Lamellar, yellow. 4.00

—Hardness 4—5

- 221. Cervantite. Antimony Ocher. Sb<sub>2</sub>O<sub>3</sub>.Sb<sub>2</sub>O<sub>5</sub>. Orthorhombic, acicular.
- 770+ massive, yellowish-white. .40
- 7710 pulverulent, sulphur-yellow, with stibnite. .40
  - 222. Stibiconite. Sb<sub>2</sub>O<sub>4</sub>.H<sub>2</sub>O(?). Massive.

Volgerite. Sb<sub>2</sub>O<sub>5</sub>.4H<sub>2</sub>O(?). Amorphous, white.

Rivotite. Sb,Cu oxide and carbonate. Amorphous, green. Stibianite. Sb<sub>2</sub>O<sub>5</sub>.H<sub>2</sub>O. Massive, reddish-yellow.

Stibioferrite. Chiefly Sb<sub>2</sub>O<sub>5</sub>. Amorphous, yellow.

772° Partzite. Contains Sb<sub>2</sub>O<sub>3</sub>, Cu<sub>2</sub>O, Ag<sub>2</sub>O and H<sub>2</sub>O. Massive, greenish-black. 2.00

Stetefeldtite. Chiefly Sb<sub>2</sub>O<sub>5</sub>,Ag,Cu and H<sub>2</sub>O. Massive, brownish-black.

## III. Oxides of the Metals

## A. Anhydrous Oxides

- I. Protoxides. R<sub>2</sub>O and RO. Hardness 1.5 and 3.5
- 223. Water. H<sub>2</sub>O. Exists in three states:—
  - 1. Solid, Ice (hexagonal), massive.

Snow, delicate six-rayed stellate crystals of ideal symmetry and of very great variety and complexity.

Type Species No. No.

Frost, arborescent and other crystalline to crystallized forms.

Hail, often crystalline, rarely in distinct quartzoids.

- 2. Liquid, Water.
- 3. Gas, Steam and Aqueous Vapor.
- 773°224. Cuprite, Ruby Copper. Cu<sub>2</sub>O. Isometric, ideal cubes, small, translucent red. 2.50
- 774\* octahedrons, minute, perfect. 2.00
- 775° dodecahedrons, minute, sharp. 2.00
- 776 dodecahedron and octahedron modifying cube, symmetrical, small. 3.00
- 777 highly modified, definite, small. 2.00
- 778+ Chalcotrichite, capillary (cubes elongated in the direction of octahedral axis), adamantine, ruby-red. .75
- 779 Tile Ore, earthy, impure. .50
- 780+ massive, fine granular, dark red. 1.50
- 781 compact massive. 1.50
- 7820 altering to malachite, ideal dodecahedron, loose. 1.50
- 783 ditto, hollowed octahedron, loose, definite. 1.50
- 784 Hydrocuprite. Hydrated cuprite (?). Coating, yellow. .50

#### Periclase Group. RO. Isometric. Range of Hardness 5.5—6

225. Periclase. MgO. Isometric, minute cubes, clear colorless.

785° grains altering to brucite, disseminated in hausmannite, translu-

cent gray. 1.25

786 grains in volcanic limestone, green.
2.00

226. Manganosite. MnO. Isometric, minute octahedrons.


787° disseminated cleavages, lustrous dark emerald green, dull black

on exposure. 2.00

227. Bunsenite. NiO. Isometric, minute octahedrons.

———Hardness 4—4.5

788 228. Zincite. ZnO. Hexagonal, hemimorphic, minute hexagonal pyramid p, with short prism m and base c, (fig.), well defined, bright red. 9.00



76 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Zincite—Continued No. DANA'S SYSTEM	1
789° foliated cleavage, deep red. 1.50	
790 crystalline granular, large irregular nodules	in calcite
cleavage. 1.00	
791+ crystalline, disseminated in franklinite, deep	red75
792° massive with willemite. 1.00	
793 finely disseminated, orange-yellow75	
Hardness 2	
229. Massicot, Plumbic ocher. PbO. Massive, scaly c	rvstalline.
794+ earthy, orpiment-yellow. 2.00	.,
795 II. Montroydite. HgO. Orthorhombic, small slend	er prisms.
clear orange-red. 8.00	
Hardness 3—4	
796°230. Tenorite. CuO. Monoclinic, minute very thin	long flexi-
ble scales, glistening gray, on lava. 2.50	
	1.00
798 Melaconite, pulverulent, dull black. 1.00	
S. Paramelaconite. Essentially CuO+Fe <sub>2</sub> O <sub>3</sub> . To	etragonal.
pyramidal, brilliant black.	
Lime. CaO. In lava.	
II. Melanochalcite. Chiefly CuO with some SiO <sub>2</sub> , Massive, black.	CO <sub>2</sub> ,H <sub>2</sub> O.
II. Sesquioxides. R <sub>2</sub> O <sub>3</sub>	
Hematite Group. Rhombohedral. Range of Hardne (Corundum 9)	ss 5—6·5
231. Corundum. Al <sub>2</sub> O <sub>3</sub> . Rhombohedral. Transpare	ent vari-
eties are precious.	
799+ Sapphire, clear blue, bipyramidal, pyramids adamantine, loose (3). 1.00	$\mathbf{z}$ and $V$ ,
800° Sapphire, light blue, water-worn60	
801° Star Sapphire (asteriated), subtranslucent d	ark blue.
water-worn, loose (6)40	
802 ditto, polished crystal. 1.50	
803* Oriental Ruby, clear dark red, rolled grains. 1.	.00

Oriental Ruby, subtranslucent light red, prism and two

pyramids, in graphitic limestone. 4.00

Oriental Ruby, light red, cleavage. 1.25

804

805

	HEMATITE GROUP 77
Type Species No. No.	Corundum—Continued
806	Oriental Topaz, clear yellow. 2.00
807	Oriental Emerald, clear green. 3.00
808	Oriental Amethyst, clear purple. 3.00
809+	grayish prisms in feldspar50
810	ditto, barrel-shaped, very large, stout. 3.00
8110	ditto, large, slender, loose (3)50
812	twins, polysynthetic. 2.50
8130	gray, cleavage50
8140	gray, parting75
815	crystalline, coarse granular. 1.00
8160	white, cleavage. 1.00
817+	Emery, granular, black20
8180	altered crystals; rough, loose (12)50
232. H	Iematite. Fe <sub>2</sub> O <sub>3</sub> . Rhombohedral.
	<ol> <li>Specular Iron (splendent black) types are sharp and perfect:—</li> </ol>
819	cuboid rhombohedron r, modified by rhombohedron e and base c. 1.50
820	thin tabular (fig.), with rutile. 1.00
821+	thin tabular, small, with smoky
	quartzoids60 820. Hematite
822+	curved rhombohedron u, rhombo-
	hedron $r$ , pyramid $n$ (fig.).
	$\begin{array}{c c} .60 \\  \end{array}$
823	short prism $m$ , base $c$ , modified by
0	rhombohedron d, loose. 1.00
8240	ditto, minute long prisms. 1.00
8250	modified tables, minute, in porous
	14 va20
826	drusy on lava, microscopic, indigo-
0	blue75
8279	Basanomelan, "Eisenrosen,"
0.0	rosette-like group (fig.). 1.50
828	twin, comp. face $\bot$ to base $c$ ,
8290	tabular. 1.50 repeated twin, tw.pl. prism m. 2.00 827. Hematite
830	repeated twin, tw.pr. prisin w. 2.00
8310	parting $   r$ , twinning striæ on c75 parting $   c$ , thick lamellar50
832	thin lamellar, bent40
833*	micaceous, foliated40
~33**	1111-110-10-10-10-10-10-10-10-10-10-10-1

78 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species Hematite—Continued
No. No.
834 granular massive30
835+ compact massive, dark red20
836* 2. Compact columnar, Pencil Ore, fine long divergent60
837° short fibrous reniform, "Kidney Ore." .60
838° 3. Red Ocher, earthy30
839 Reddle (red chalk), clayey30
840° 4. Clay Iron-stone, Argillaceous Hematite. impure
brownish20
841 Jaspery Clay Iron-stone, reddish20
842+ Lenticular Iron Ore, Fossil Ore, oölitic ocherous, red20
843° Martite. Fe <sub>2</sub> O <sub>3</sub> . Isometric. Probably pseudomorph after
either pyrite or magnetite, or both. Small sharp
octahedrons, loose (lot)35
844 octahedrons, minute in chlorite35
octahedrons, bright, perfect75
846 dodecahedrons, symmetrical. 1.25
847+ dodecahedrons and octahedron75
Raphisiderite. Fe <sub>2</sub> O <sub>3</sub> . Orthorhombic(?), minute needles
233. Ilmenite, Titaniferous Iron. Generally FeTiO <sub>3</sub> . Rhom-
bohedral, tetartohedral, iron-black:-Varieties fol-
low in order of Ti p.c.
Kibdelophane. About 30 p.c. Ti.
848° Crichtonite, about 30 p.c. Ti, small tables40
849º Ilmenite, 26—30 p.c. Ti, loose crystal75
850* Ilmenite, lamellar massive25
Menaccanite, about 25 p.c. Ti. Large crystal. 3.00
852+ Menaccanite, granular massive25
853 Menaccanite, sand40
Hystatite, 15—20 p.c. Ti.
854 Washingtonite, 15—20 p.c. Ti, tabular75
Uddevallite, about 10 p.c. Ti.
Kragerö hematite. Less than 3 p.c. Ti.
Magnesian Menaccanite, Picrotitanite (FeMg)TiO <sub>3</sub> .
S. Pyrophanite. MnTiO <sub>3</sub> . Rhombohedral, scale-like crys-
tals, deep blood-red.
I. Senaite. (Fe,Pb)O.2(Ti,Mn)O <sub>2</sub> . Tri-rhombohedral, black
855 Iserine. Titanic iron. Isometric or rhombohedral, minute
octahedroids in sand 40

## III. Intermediate Oxides

8710

Chemically considered, these species are properly aluminates, ferrates, manganates, etc. and in a strict classification would be placed in section 5 of the Oxygen-Salts.

# Spinel Group. RO.R<sub>2</sub>O<sub>3</sub>. Isometric.

Type Species No. No.	Range of Hardness 6.5—8
	pinel. MgO.Al <sub>2</sub> O <sub>3</sub> . Isometric, symmetrical crystals:—
856	bluish-gray, rough indistinct cube. 1.50
857+	gray, octahedron75
858	Ruby-Spinel, Magnesia Spinel, small octahedrons, clear
	deep red, brilliant, loose (12). 1.00
8590	ditto, hemitrope or "spinel twins," tw.pl.
	and comp. face octahedron o (fig.),
	loose, (3). 1.00
860+	ditto, clear octahedrons, slightly water-
	worn (lot)40
	Balas-Ruby, clear rose-red.
861	bluish-gray octahedron. 1.00
862	Ceylonite, Pleonaste, Iron Magnesia 859. Spinel
	Spinel, octahedron, large, black. 3.00
863°	ditto, sharp, minute, on lava. 1.00
864	ditto, dodecahedron d modifying octahedron o, sharp
06-0	splendent. 1.50
865°	ditto, trapezohedron m modifying
866*	octahedron o (fig.) large. 3.00 ditto, rolled pebbles (lot)40
867°	Chlorospinel, Magnesia-Iron Spinel,
007	grass-green (due to presence of
	Cu). 2.00
	Picotite, Chrome-Spinel, brownish.
868 <b>©235.</b> I	Hercynite. FeAl <sub>2</sub> O <sub>4</sub> . Isometric, granular, 865. Spinel
	black50
<b>236.</b> C	Gahnite, Zinc Spinel. ZnAl <sub>2</sub> O <sub>4</sub> . Isometric.
869	Automolite, octahedron in talcose schist, green. 1.50
870*	ditto, sharp, splendent octahedrons with fowlerite. 2.50

Dysluite. (Zn,Fe,Mn)O.(Al,Fe)2O3, octahedron, sharp,

Kreittonnite. (Zn,Fe, Mg)O.(Al,Fe)<sub>2</sub>O<sub>3</sub>, granular.

bright. 2.00

80 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species No. No.
No. No. 237. Magnetite, Magnetic Iron Ore. FeO.Fe <sub>2</sub> O <sub>3</sub> . Isometric,
crystals of ideal symmetry, iron-black:—
872 octahedrons, imperfect, loose (12)40
873+ octahedrons in chlorite, small but sharp40
874° octahedrons with adularia, splendent. 1.25
875° octahedrons curved75
876* dodecahedrons, striated deeply
(fig.), splendent. 1.00
877 dodecahedrons, dull75
878° trapezohedron m modifying octa- hedron o, loose, (3). 1.00
879° twins, tw.pl. o, small, perfect. 1.25
880 twins, polysynthetic, splendent.
1.00
881° parting, octahedral50
882+ granular massive, coarse20 876. Magnetite
granular massive, fine20
884 sand30
885+ Lodestone, with polarity, compact50
886 dendritic in muscovite20
magnesian. (Fe,Mg)O.Fe <sub>2</sub> O <sub>3</sub> .
nickeliferous, 1.76 p.c. NiO.
titaniferous.
Manganmagnetite, 3.80 to 6.27 p.c. Mn.
ocherous, earthy40
Nickel Oxide. NiO.Ni <sub>2</sub> O <sub>3</sub> (?). Sand.
888 238. Magnesioferrite. MgO.Fe <sub>2</sub> O <sub>3</sub> . Isometric, minute octa-
hedrons in limestone, black. 2.00
889*239. Franklinite. (Fe,Zn,Mn)O.(Fe,Mn) <sub>2</sub> O <sub>3</sub> . Isometric, octa-
hedron, well defined, bright. 1.25
890 octahedron, rounded, iron-black75
891° octahedron $o$ modified by dodecahedron $d$ , distinct. 2.00
892 trapezohedron m, dodecahedron d modifying octahedron
o, bright. 3.00
893+ disseminated grains with zincite in willemite40
894 massive granular, coarse40
895° massive compact, iron-black40
896°240. Jacobsite. (Mn, Mg)O. (Fe, Mn) <sub>2</sub> O <sub>3</sub> . Isometric, minute
octahedrons, sharp and bright. 1.50
897 fine granular, deep black. 1.00

```
Type Species No. No.
898°241. Chromite, Chromic Iron.
                                         FeO.Cr<sub>2</sub>O<sub>3</sub>. Isometric, minute
                  octahedrons, loose sand. .50
             massive granular, iron-black.
899+
900
             massive compact. .20
             Mitchellite, 2MgAl,O4.MgCr,O4.FeCr,O4.
             Chrompicotite and Magnochromite are Mg chromites.
           Plumboferrite. 2FeO.Fe<sub>2</sub>O<sub>3</sub>.PbO.Fe<sub>2</sub>O<sub>3</sub>(?).
                                            -Hardness 8.5
901 242. Chrysoberyl. BeO.Al<sub>2</sub>O<sub>3</sub>. Orthorhombic, prismatic, clear
                  pale green, precious, loose. 3.00
9029
             ditto, water-worn (lot).
             tabular, pale green, well-defined. 2.00
903
             repeated twin, pseudo-hexagonal (fig.),
904+
                  sharp. 1.00
             Alexandrite, repeated twin, re-entrant
9050
                  angles (fig.), subtransparent em-
                  erald-green, columbine-red by
                                                           904. Chrysoberyl
                  artificial light. 2.50
906
             Cat's-Eye, chatoyant, greenish, cut. 5.00
                           -Range of Hardness 4.5—6 (Minium 2—3)
907°243. Hausmannite.
                              MnO.Mn_2O_3.
                                              Tetragonal, octahedroids,
                  sharp, bright. 2.00
             repeated twin (fiveling). 3.00
908
             massive granular, brownish-black. .75
909+
      II. Coronadite. MnO<sub>2</sub>.PbO.Mn<sub>3</sub>O<sub>4</sub>.
                                                  Mas-
                  sive, black.
910°244. Minium. 2PbO.PbO<sub>2</sub>. Earthy, red. 4.00
     245. Crednerite. 3CuO.2Mn<sub>2</sub>O<sub>3</sub>. Monoclinic,
                  foliated, black.
                                                           905. Chrysoberyl
911°246. Pseudobrookite. 2Fe<sub>2</sub>O<sub>3</sub>.3TiO<sub>2</sub>(?). Or-
                  thorhombic, minute tables, sharp, adamantine,
                  blackish. 2.00
912*247. Braunite.
                       3Mn<sub>2</sub>O<sub>3</sub>.MnSiO<sub>3</sub>.
                                              Tetragonal, octahedrons,
                  bright, sharp. 1.50
9130
             massive, blackish.
        I. Bixbyite. FeO.MnO<sub>2</sub>. Isometric, cube a with trapezo-
914
                  hedron n, black. 2.00
```

## IV. Dioxides. RO2.

## Rutile Group. Tetragonal.

	zianio oronti romagonam
Type Species	Hardness 6—6.5 (Plattnerite 5—5.5)
	Cassiterite, Tin Stone. Sn O <sub>2</sub> . Tetragonal, prism and pyra-
	mid, symmetrical, splendent brown. 1.50
9169	acicular, "needle ore," modified. 2.00
917•	twin, contact, tw.pl. diametral pyra-
<i>y-1</i>	mid e, adamantine, sharp. 2.50
918•	ditto, penetration, bright black,
<b>9.</b> 0	loose75
919*	repeated twin, well defined (fig.)
919+	1.50
920	reniform with fibrous structure. 2.00
921+	massive, fine granular, grayish. 1.00
-	massive, coarse granular, brownish.
922	I.25
923	massive, compact, yellowish. 1.50
924	disseminated finely in gray quartz30
925+	disseminated coarsely in greisen, brown30
9260	Wood Tin, concentric, radiated. 1.50
927+	Stream Tin, much rounded, fine sand50
928	Stream Tin, coarse angular grains50
929®	Stream Tin, rounded boulder. 1.00
930	Stream Tin, ferruginously cemented (a variegated con-
	glomerate). 1.50
931 A	Ainalite. A cassiterite containing 8.78 p.c. Ta <sub>2</sub> O <sub>5</sub> . Pyr-
	amidal, adamantine, blackish. 4.00
932 249. I	Polianite. MnO <sub>2</sub> . Tetragonal, minute prisms m and h, pyra-
	mids $s$ and $e$ , composite parallel groupings. 2.00
933+	crystalline, radio-fibrous, iron-gray. 1.00
934+250. I	Rutile. TiO2. Tetragonal, unit and
	diametral prisms $m$ and $a$ , diam-
	etral pyramid e (fig.), sharp and
	symmetrical, metallic-adaman-
	tine, red, loose50
935	ditto, with additional ditetragonal
	prism l and unit pyramid s50
9369	pseudo-rhombic modified, highly  934. Ruttle
	splendent, sharp. 2.00

Twne Species	RUTILE GROUP Rutile—Continued	8 <u>'</u> 3
Type Species No. No.		
937°	acicular, sharp, translucent red, loose (6). 1.00	
9380	twin, tw.pl. diametral pyramid e,	
930-	geniculated, perfect, loose75	1
939	twin, modified, splendent. 2.00	1
94 <b>0*</b>	repeated twin, tw.pl. e, deeply striated,	7
74-	brownish-red, loose75	/
941	twin reticulated. 1.25	
	twinning lamellæ    e. 947. Rutile	
942	capillary75	
•	capillary (enclosure), "Flêches d'amour." See quartz.	
943	water-worn crystals, brownish, loose (lot)50	
944	pebbles reddish-black, grayish exterior (lot)50	
945°	cleavage, dark red. 1.00	
946	disseminated, garnet-red50	
947°	ferriferous, twin, tw.pl. e, eightling	
	(similar to fig.), black, loose50	
948+	ferriferous, crystallized aggregate50 $p \mid p \mid$	
	ferriferous, Ilmenorutile, black.	4
7	chromiferous, grass-green.	/
	serite. FeTi <sub>2</sub> O <sub>3</sub> . Brown grains.  Davidite. TiO <sub>2</sub> with Fe,U,V,Cr, and rare	
11. L	earths. Cuboids, black.	
251 D	Mattacrita PhO Tatraconal priems	
231. F	iron-black.	<b>:</b>
949	massive. 7.00	
777	Hardness 5.5—6	
252 (	Octahedrite, Anatase. TiO <sub>2</sub> . Tetragonal, small adamantin	
232.	crystals of ideal symmetry:—	ıc
950+	acute octahedroid habit, unit pyramid $p$ , (fig.), black	le
930	1.50	
951	ditto, yellowish-brown. 2.00	
95 <b>2</b>	diametral prism $a$ , unit pyramid $\pi$ . 2.00	
953 <b>°</b>	diametral prism a, highly modi-	
- 00	fied termination, translucent	
	yellowish brown. 1.25	\
954	tabular,    to base $c$ , with unit	)
	pyramid p and diametral	•
	prism a, symmetrical, dull	
	black, loose75 955. Octahedrite	

84 Type Specie No. No	COMPLETE TYPE COLLECTION. DANA'S SYSTEM Octahedrite—Continued
955°	obtuse pyramid z and diametral prism a, modified by unit prism m, pyramids p, v and diametral pyramid e (fig.), transparent brown. 2.00
956 253	Brookite. TiO <sub>2</sub> . Orthorhombic, small tabular with quartz and chalcopyrite. 4.00
957+	very thin tabular, highly modified, perfect, transparent hair-brown. 1.50
958+	Arkansite, unit prism m, and pyramid e truncated by pyramid z, small, sharp, symmetrical, splendent black (similar to fig.)75
959	Arkansite, ditto, with brachydome t. m m
960*	Arkansite, paramorphosed to rutile, unit prism m, unit pyramid z (fig.), symmetrical, dull black, loose50  Brookite
961	ditto, m with pyramid e50  Hardness 2-2.5
962 <b>254</b>	. Pyrolusite. MnO <sub>2</sub> . Orthorhombic, pseudomorphous (?), small distinct prisms, bright iron-black. 1.00
963	acicular. 1.00
964*	thick tabular, small, perfect. 1.25
965°	columnar crystalline, bright. 1.00
966°	radio-fibrous crystalline50
967+	fine granular crystalline20
968	granular massive, dull20
969	reniform coating50
Ι	. Baddeleyite, Brazilite. ZrO <sub>2</sub> . Monoclinic, twins, tabular    a.
970°	reniform, concentric, greenish-gray. 4.00
	B. Hydrous Oxides
	Hardness of Turgite 5—6
971 <b>°2</b> 55	. Turgite. 2Fe <sub>2</sub> O <sub>3</sub> .H <sub>2</sub> O. Compact fibrous50
972	massive botryoidal, reddish-black50
073*	earthy, red20

# Diaspore Group. $R_2O_3.H_2O.$

Type Species	Hardness 7, 5 and 4
Type Species No. No.	riaspore. Al <sub>2</sub> O <sub>3</sub> .H <sub>2</sub> O. Orthorhombic, rounded prisms
974+236. D	flattened $  b $ , brilliant clear violet, on emery. 2.00
075	ditto, acicular, grayish-white. 1.50
975 976	stout prisms, whitish, with margarite. 3.00
970 977®	foliated columnar, reddish-gray. 1.50
	Southite. $Fe_2O_3$ . $H_2O$ . Orthorhombic, very thin small
970-237.	tables, Rubinglimmer, red. 1.00
979	long thin tabular, blackish-brown. 1.25
9800	acicular, in radial aggregates. 1.00
981*	Sammetblende, velvety globular crusts of minute radiat-
	ing capillary crystals, yellowish-brown. 1.25
	Onegite, acicular (enclosures). See quartz.
982	columnar, dark brown. 1.00
983+	fibrous, concentric radiated, reniform. 1.00
984	scaly-fibrous, Lepidocrocite. 2.00
	compact massive, conchoidal fracture.
	disseminated microscopic crystals afford some varieties
	of aventurine (feldspars etc.).
985 <b>°258</b> . N	Langanite. Mn <sub>2</sub> O <sub>3</sub> .H <sub>2</sub> O. Orthorhombic, small flat prisms
	terminated by base c, distinct, bright, iron-black.
	1.50
986	long prisms terminated by rough zone of macropyramids
_	$\rho$ ,s, $p$ etc., large, splendent. 3.00
987	acicular prisms. 1.50
988+	fibro-columnar, radiated. 1.00
	Hardness 5—5.5
989 <b>°259</b> . L	imonite, Brown Iron Ore. 2Fe <sub>2</sub> O <sub>3</sub> .3H <sub>2</sub> O. Massive, com-
	pactly radio-fibrous, stalactitic, brownish40
990	stalactite, concentric structure40
991	compact, botryoidal50
992+	compact, subfibrous structure, mammillary, shining
	black surface30
993*	compact, globular crust, iridescent bronze50
994	compact, globular crust, iridescent variegated. 1.00
995	ocherous, brown20
996+	ocherous, yellow20
997°	Bog Ore, porous, coherent20

86 Type Species No. No.	COMPLETE TYPE COLLECTION. DANA'S SYSTEM  Limonite—Continued
998	Bog Ore, porous, loose, plant remains50
999+	Brown clay-ironstone, compact20
1000	ditto, concretionary40
10010	ditto, "pipe ore," hollow tube50
1002	ditto, pisolitic40
10039	ditto, oölitic30
	Esmeraldaite. Hyd. Fe <sub>2</sub> O <sub>3</sub> . Massive, black.
	Hardness 2.5 and Soft
1004°260	. Xanthosiderite. Fe <sub>2</sub> O <sub>3</sub> .2H <sub>2</sub> O. Long divergent fibro-columnar, concentric, brown75
1005+261.	Bauxite. Al <sub>2</sub> O <sub>3</sub> .2H <sub>2</sub> O. Oölitic, yellowish20
10060	pisolitic, red20
1007	clay-like, Wocheinite, grayish35
	Brucite Group. R(OH) <sub>2</sub> .
	<u>-</u>
	Rhombohedral. Hardness 2.5
1008 262.	Brucite. MgO.H <sub>2</sub> O. Rhombohedral, broad tabular    base c with rhombohedron r and pyramid p, greenish-gray. 2.50
10090	ditto, rosette-like aggregate. 1.50
1010+	broad cleavage, pearly white. 1.00
1011	foliated, in serpentine. 1.00
10120	Nemalite, 4 to 5 p.c. FeO, fibrous. 1.25
1013	Manganbrucite, contains much Mn. 1.50
	Eisenbrucite. An altered brucite.
1014 263.	Pyrochroite. MnO.H <sub>2</sub> O. Rhombohedral, rounded hexa-
	gons, white becoming bronze and finally black. 2.50
10150	foliated crystalline. 1.50
	Hardness 2·5—3·5 and Soft
1016 264.	Gibbsite. Al <sub>2</sub> O <sub>3</sub> .3H <sub>2</sub> O. Monoclinic, hexagonal aspect,
	(Hydrargillite) minute, pearly greenish. 2.00
1017+	drusy, minutely radio-lamellar incrustation40
10180	stalactitic, smooth40
	Richmondite. A hydrate containing 37 p.c. P <sub>2</sub> O <sub>5</sub> .
	Zirlite. Al hydrate. Amorphous.
1019 <b>•26</b> 5.	Sassolite. B <sub>2</sub> O <sub>3</sub> .3H <sub>2</sub> O. Triclinic, minute scales, pearly white, loose (lot)75

```
Type Specie
No. No.
```

-Soft

1020 266. Hydrotalcite. Al<sub>2</sub>O<sub>3</sub>.6MgO.15H<sub>2</sub>O(?). Hexagonal, translucent pearly white. .75

Houghite (altered from spinel). .50

1022°267. Pyroaurite. Fe<sub>2</sub>O<sub>3</sub>.6MgO.15H<sub>2</sub>O(?). Hexagonal, minute six-sided tables, pearly pale yellow. 2.00

Hardness 2.5 and 5.6

1023 268. Chalcophanite. (MnZn)O.2MnO<sub>2</sub>.2H<sub>2</sub>O. Rhombohedral, druses of minute tables, bluish-black. 2.50

small botryoidal, subfibrous. .75

1025+269. Psilomelane. H<sub>4</sub>MnO<sub>5</sub>(?). Massive, bluish-black. .20

botryoidal, dull iron-black. .75

reniform, rough. .40

stalactitic, smooth. 1.00

1029 Lithiophorite. 10—15 p.c. Al<sub>2</sub>O<sub>3</sub>, 1·2—1·4 p.c. Li<sub>2</sub>O, 12·6—15·4 p.c. H<sub>2</sub>O. Botryoidal, bluish-black. 1.00

1030+ Wad. (A) Bog Manganese. Impure hydrated Mn oxide, loose earthy, black. .20

1031º ditto, dendritic on rhyolite. .40

1032 (B) Asbolite, Earthy Cobalt. Black. .50

1033° (C) Lampadite, Cupreous Manganese. 4 to 18 p.c. CuO. Earthy, black. .50

Varvicite. An altered manganite.

1034 II. Brostenite. Manganite of Mn and Fe<sup>II</sup>. Massive, black. 2.50

## Appendix to Oxides

Delafossite. CuO 47·45, Fe<sub>2</sub>O<sub>3</sub> 47·99, Al<sub>2</sub>O<sub>3</sub> 3·52. Heterogenite. Essentially CoO.<sub>2</sub>Co<sub>2</sub>O<sub>3</sub>+6H<sub>2</sub>O. Heubachite. 3(Co,Ni,Fe)<sub>2</sub>O<sub>3</sub>+4H<sub>2</sub>O(?). Massive. Namaqualite. Nearly Al(OH)<sub>3</sub>. 2Cu(OH)<sub>2</sub>. 2H<sub>2</sub>O. Rabdionite. Near asbolite. Earthy, black. Transvaalite. Co<sub>2</sub>O<sub>3</sub> 65·80, As<sub>2</sub>O<sub>5</sub> 5·79, H<sub>2</sub>O etc.

# VI. Oxygen-Salts

#### 1. Carbonates

## A. Anhydrous Carbonates

1. Calcite Group. RCO<sub>3</sub>. Rhombohedral.

Hardness 3.5-4 (Calcite 3, Smithsonite 5).

Type Species

1035 1036+

270. Calcite, Calc Spar. CaCO<sub>3</sub>. Rhombohedral.

#### A. VARIETIES BASED ON CRYSTALLIZATION AND IMPURITIES:-

#### ORDINARY

(a). Symmetrical well defined crystals, transparent to translucent, usually glassy colorless to white. Very great multiplicity of forms including highly

complex combinations. Only the commoner are here described:—simple rhombohedron r (fig.). 1.50 ditto, modified by scalenohedron v. 1.00 flat rhombohedron e, in quartz geode

flat rhombohedron e, in quartz geode
(fig.). .50

1038 ditto, parallel grouping, large. .50

1039 ditto, with short prism m, "nail-head spar" (fig.). .50

1040 cuboid rhombohedron φ (fig.). .50

1041 acute rhombohedron M. 1.00 ditto, with base c. 1.50


positive and negative rhombohedrons.
1.25

Papierspath, very thin hexagonal tables, rounded rhombohedrons prominent, flower-like aggregate, glistening icy aspect. .75

prism m and base c (fig.), ideal symmetry. 1.00



1035. Calcite





1040. Calcite

# CALCITE GROUP

Type Species No. No.	Calcite—Continued
10469	ditto, with diagonal prism a. 1.25
1047	prism $m$ , acute rhombohedron $\rho$ flat
	rhombohedron $e$ and base $c$ . 2.00 $\binom{m}{m}$ $\binom{m}{m}$
1048	tabular    base $c$ , prism $m$ , modified. 1.00
1049+	Dog-tooth Spar, scalenohedron v (fig.),
	ideal symmetry. 1.00
1050	scalenohedron $v$ and base $c$ . 1.50
1051+	scalenohedron $v$ , terminated by rhombohedron $r$ (fig.) amber-yellow, very large, loose50
1052	ditto, white with "phantom." .50
1053+	scalenohedron v, rhombohedron r and prism
	m (fig.), large75
10540	two scalenohedrons B and v with rhombo-
	hedrons s and e, ferruginous brick-red
	"phantom." .75
1055	scalenohedron v built up of small rhombo-
1056	hedrons, large75 acute scalenohedron y, spire-shaped. 1.00
1056 1057 <b>°</b>	three scalenohedrons $v$ , $y$ and $M$ with rhombohedron $r$
105/0	(fig.). 1.00
10589	complex highly modified crystal. 1.25
1059	twin, tw.pl. base c, the two rhombic individ-
	uals having the same vertical axis. 2.50
10600	twin, ditto, but scalenohedral (fig.), large.
_	1.25
1061*	twin, tw. pl. rhombohedron e, the scaleno-
	hedral individuals having vertical
	axes inclined 127° 29½' and 52° 30½' (fig.), large, loose75
1062	twin, tw.pl. r, individuals with vertical
1002	axes inclined at 90° 46′ and 89° 14′, "but-
	terfly twin" (fig.), large, loose. 2.50
	(b) cleavages, rhombohedral:—
10630	Iceland Spar, doubly refracting, clear. 2.00
1064+	ditto, pale amber. 1.00
10659	salmon-red cleavage30
1066	sky-blue cleavage50
1067	twin cleavage, tw.pl. e50 1053. Calcute
10680	asteriated cleavage50

twinning lamellæ indicated by surface striæ.

1069 (c) drusy botryoidal, milky. .30

(d) Brunnerite, bluish, chalcedony-like.

(e) Reichite, r r 74° 40', Gr. 2.67 (?).

(f) Fontainebleau Limestone, 50-63 p.c. quartz sand, acute rhombohedron f (fig.), ideal symmetry, gray, loose. .50

ditto, nodular aggregate. 1.50 1071 ditto, acute hexagonal pyramid of second 10729

order  $\gamma$ , symmetrical and definite, but ends rounded, very large, loose. 1.00

ditto, aggregate with quartz pebbles, very large. 1.00 (g) Hislopite, impure, grass-green.

minute hollow channels within (hohle canale of Rose), which are connected with

II. FIBROUS AND LAMELLAR KINDS, white Satin Spar, silky, finely fibrous. Argentine, curved lamellar, pearly. .40

Aphrite, scaly foliated, pearly. III. GRANULAR MASSIVE TO CRYPTOCRYS-TALLINE KINDS

1. Granular limestone. (Saccharoidal):-Statuary marble, Carrara, fine grained, white. .20

Architectural marbles follow:-

1078+ coarse grained, white. .20 Cipolin, white with green-1079 ish shadings. .75

1080 Giallo antico, yellowish with whitish spots. .75

10819 Siena, light yellow, with reddish veins. .30 1082

Mandelato, light red with whitish spots. .75

Bardiglio gray, clouded.

Turquois-blue, veined with white. .40

Verd-Antique, clouded greenish, due to presence of serpentine. .30




1057.

Caldite



1060. Calcite



1083 1084

1070\*

1073

10749

1075°

1076

1077

10859

# CALCITE GROUP Calcite—Continued

Type Species	Calcite—Continued
NG. 110.	2. Hard compact limestone (marbles):-
1086	black30
	yellow.
	red.
1087	fetid, black, Anthraconite, Stink- stone .40
10889	Portor (Egyptian), black, veined yellow30
	Panno-di-Morte, black with white 1062. Calcite shells.
	Marble of Languedoc, red with white fossils.
	Griotte, brown spotted red and white.
	Sarencolin, deep red, gray and yellow.
1089	Bird's-eye, gray with whitish points30
1090	Shell-marble, fossiliferous30
10919	Madreporic marble, coralline, polished75
1092+	Tennessee, encrinal, mottled reddish20
1093	Lumachelle, "fire marble," chatoyant. 1.00
10949	Ruin Marble, brownish, polished. 1.00
10959	Landscape Marble, gray, polished. 1.25
10969	Breccia Marble, cemented fragments30
1097*	Lithographic stone, smooth even-grained20
1098	Pudding-stone marble, cemented rounded pebbles30
1099+	Hydraulic Limestone, or "Cement Rock," contains Mg,
	Al and Si as impurities20 3. Soft compact limestone:—
1100+	Chalk, white20
1101	Calcareous marl, loose, earthy impure. $f = f = f$
1101	.20
	4. Concretionary massive:—
1102*	Oölite, minutely rounded granular20
	Pisolite, see ktypeite.
	5. Deposited by calcareous waters or in rozo, Calcite
	caverns:—
1103+	Stalactite, from roof of cavern, buff40
1104	Stalactite, translucent, white60
,	Stalactite, translucent, sea-green.
11059	Stalagmite, from floor of cavern40
1106+	Mexican Onyx, irregularly banded, translucent white,
1100	pale green, etc., polished60

92 Type Spec No. N	COMPLETE TYPE COLLECTION. DANA'S SYSTEM Calcite—Continued
1107	Clouded Onyx, translucent, variegated60
1108	Brecciated Onyx, variegated, polished. 1.00
1109*	Travertine, very coarse, irregularly and indistinctly
	banded, yellowish60
1110+	Calc Tufa, moss-like porous structure20
IIII	Calc Tufa, perfect leaves, porous mass60
1112	Agaric mineral, Rock-milk, soft, crumbling40
11139	Rock-meal, cotton-like, loose40
	B. VARIETIES BASED UPON COMPOSITION:—
1114	Dolomitic calcite, contains MgCO <sub>3</sub> 20
	Baricalcite, contains some BaCO <sub>3</sub> , rhombohedrons,
	grayish-white.
1115	Strontianocalcite, contains SrCO <sub>3</sub> , minute acute rhom-
Ū	bohedrons. 2.00
11160	ditto, opaque globular aggregates. 2.00
1117*	Ferrocalcite, contains FeCO <sub>3</sub> , acicular, brown50
1118	Ferrocalcite, globular, concentric structure50
	Manganocalcite, see Agnolite.
	Zincocalcite, containing ZnCO <sub>3</sub> .
11190	Plumbocalcite, contains PbCO <sub>3</sub> , rhombs. 1.25
11200	altered to dolomite. 1.00
1121	altered to siderite. 1.00
1122	altered to calamine. 1.25
11230	altered to smithsonite75
1124	altered to quartz crystals. 1.00
1125+	altered to chalcedony. 1.00
1126	altered to hyalite. 2.00
	altered to copper.
11279	Thinolite. Pseudomorphous. CaCO <sub>3</sub> . Acute tetragonal
	(?) pyramids, skeleton structure, forming tuffaceous
	aggregate, grayish. 1.00
11280	I. Ktypeite (formerly called Pisolite). CaCO <sub>3</sub> . Specific
	gravity and optical properties differ from calcite
	and aragonite. Mass of cemented pea-like con-
	cretions, yellowish-white50
21	71. Dolomite. CaCO <sub>3</sub> .MgCO <sub>3</sub> . Rhombohedral, tetartohedral,
	well defined rhombohedrons:—
	r Stanotural Variation.

rhombohedron r, ideal symmetry, transparent, vitreous.

11290

2.50

Type Species No. No.	Dolomite—Continued
11300	acute rhombohedron $m$ with obtuse rhombohedron $r$ ,
	symmetrical, dull black, loose30
1131	twin symmetrical. 2.00
11320	Pearl spar, rhombic, white. 1.00
1133+	ditto, saddle-shaped (fig.), cream.
1134	ditto, in selenite. 1.00
1135	columnar crystalline50
11360	globular drusy75
30	Miemite, pale asparagus-green.
1137	granular, fine, gray20 1133. Dolomite
1138	granular, fine white marble20
1139+	granular, coarse, white20
1140*	compact, buff20
•	Compact porcellanous, Gurhofite, conchoidal, sub-trans-
	lucent, snow-white.
2.	Varieties depending on Composition:—
	Normal Dolomite. Ca:Mg=1:1. See crystals.
11419	ferriferous, contains FeCO <sub>3</sub> , massive30
	manganiferous, contains MnCO <sub>3</sub> .
	cobaltiferous, reddish.
	zinciferous.
1142	altered to steatite. 1.25
1143 271A.	Ankerite. CaCO <sub>3</sub> .(Mg,Fe,Mn)CO <sub>3</sub> . Rhombohedral.
	rhombohedrons, pearly brown. 1.00
1144*	ditto, yellowish75
11459	crystalline granular30
1146	compact massive30
1147°272. M	Lagnesite. MgCO <sub>3</sub> . Rhombohedral, rough prism. 1.00
1148	Lamellar cleavable50
11490	Fine granular30
1150+ .	Compact, like unglazed porcelain, snow-white20
1151	earthy40
11520	Pinolite, cleavage30
1153	ferriferous, Breunnerite, 5 to 10 p.c. FeO, distinct
	rhombohedrons. 1.00
1154 272A.	MESITITE. 2MgCO <sub>3</sub> .FeCO <sub>3</sub> . Rhombohedral, perfect
	rhombohedrons, transparent. 2.50
1155•	Pistomesite. MgCO <sub>3</sub> .FeCO <sub>3</sub> , granular50

		MPLETE TYPE COLLECTION. DANA'S SYSTEM
	Type Species No. No.	
		Siderite, Chalybite, Spathic Iron. FeCO <sub>3</sub> . Rhombohedral.
	I	Ordinary Varieties:—
		(a) Crystallized, brown:—
	1156+	obtuse rhombohedron r, perfect50
	1157	acute rhombohedron d, perfect75
	1158*	ditto, with base c. 1.00
	1159	ditto, curved and built of sub-indi-
	11600	viduals75 octahedroid, $s$ (0551) with $c$ (similar to
	11000	fig.), perfect, small75
	1161	base c hexagonally banded, with rhom-
	1101	bohedrons and prism. 3.00 1160. Siderite
	11620 II.	new scalenohedron y, scalenohedron
		v, rhombohedrons $f$ and $r$ (fig.),
		minute, sharply defined, splen-
		dently iridescent. 1.00
	1163	twin, tw.pl. e. 1.25
	1164°	curved saddle-shaped crystals on / /
		fluor75
	1165+	cleavage, rhombic20
	1 166 <b>9</b>	(b) concretionary, Sphærosiderite,
		concentric globular, subfibrous,
		brown30
	1167	granular crystalline, brown20
	1168	compact massive20 oölitic.
	11690	earthy, clay iron-stone, impure20 1162. Siderite
	1109*	2. Manganiferous, Oligonite.
	1170	3. Magnesian, Sideroplesite60
•	11/0	4. Calciferous, Siderodot, green.
	1171	altered to limonite50
٠	II.	Manganspherite, contains Mn.
		Thomäite. FeCO <sub>3</sub> . Orthorhombic (?).
		Rhodochrosite, Dialogite. MnCO <sub>3</sub> . Rhombohedral,
	11/2-271.	rhombohedron r, opaque pale pink. 1.50
	1173	ditto, translucent rich pink. 4.00.
	1174°	curved rhombs with fluor. 1.50
	1175°	drusy crust, small columnar structure, fawn-color75
	1176	fine granular, coated with glistening drusy quartz, deli-
	•	cate nink 1.25

.

## 2. Aragonite Group. RCO<sub>3</sub>. Orthorhombic.

#### Range of Hardness 3.5-4


277. Aragonite. CaCO<sub>3</sub>. Orthorhombic, sharply symmetrical crystals:-

prismatic.

contact-twins, tw.pl. m, the indi-11930 viduals showing unit prism m, brachypinacoid b, brachydome k and pyramids, transparent yellow. .30 1194+

repeated twins, tw.pl. prism m, prismatic, pseudohexagonal (fig.), brown, loose.

ditto, twinning striæ on base 1195 deeply marked, aggregate, brownish-red. 2.50



96 C Type Species No. No.	OMPLETE TYPE COLLECTION. DANA'S SYSTEM  Aragonite—Continued
11960	ditto, individuals separated by re-entrant prismatic angles, forming on the base a serrated hexagonal outline, aggregate, brilliant, clear colorless. 1.50
1197	ditto, tabular, white, on lava50
11980	contact-twin, pseudohexagonal spire-shaped, sub- transparent gray loose (3)50
1199*	acicular, radiating groups, white75
1200*	columnar, divergent20
1201	fibrous silky, light blue75
I 202®	massive compact20 scaly massive, snow-white.
1203°	stalactitic, Sprudelstein, fibrous50
1204+	Flos-ferri, coralloidal, slender interlacing and twisted stems, snow-white. 1.00
1205	Flos-ferri, branching botryoidal. 1.00
1206	Tarnowitzite, contains PbCO <sub>3</sub> . 1.50
1207	Mossottite, nearly 7 p.c. SrCO <sub>3</sub> and trace of Cu, columnar radiated. 1.50
12080	altered to calcite, large symmetrical pseudohexagonal tables, loose (3)50
1209	altered to drusy calcite, cellular structure outlining twinned character of the original hexagon. 1.00
II.	
1210 <b>°278</b> .	Bromlite, Alstonite. BaCO <sub>3</sub> . CaCO <sub>3</sub> . Orthorhombic, complex twins forming sharp dihexahedral pyramids, translucent whitish. 2.00
12110279.	Witherite. BaCO <sub>3</sub> . Orthorhombic, repeated twins, tw.pl.
	prism m, symmetrical pseudohexagonal pyramids, acute. 2.00
1212	ditto, very obtuse, grayish. 2.00 tuberose.
1213+	granular crystalline, whitish20
12149280.	Strontianite. SrCO <sub>3</sub> . Orthorhombic, contact-twins, tw.
	pl. prism <i>m</i> , acute pyramids and brachydomes forming very acute well defined pseudohexagonal pyramids, translucent75
1215	contact-twins, tw.pl. prism <i>m</i> , individuals marked by serrated pseudohexagonal basal outline, translucent flesh-red. 2.00
1216+	columnar crystalline, yellowish-white20
1217	Calciostrontianite. CaCO <sub>3</sub> 13·14 p.c75

Type Species

281. Cerussite. PbCO<sub>3</sub>. Orthorhombic, crystals well defined, translucent, adamantine:—

	transfucent, adamantine.—
12180	thin tabular    b, modified (fig.), clear75
1219	prismatic, white. 1.00
I220º	pyramidal, gray. 1.00
1221	penetration-twins, tw.pl. prism m. 1.50
1222*	contact-twins, tw.pl. m75
1223	repeated twins, six-rayed stellate (fig.).
12240	repeated twins, pyramidal (fig.). 1.50
12250	reticulated twinned aggregate. 2.00
1226+	aggregate of interlacing slender prisms, twinned, satiny white. 1.25
1227	aggregate of long fluted columns, stellate twins. 2.00
1228	aggregate, sheaf-like75
1229+	fine granular, brownish.
1230	loose granular, blackish,

loose granular, blackish.

1231° compact, gray. .50

fibrous, satiny. 2.00

1223. Cerussite

3. Barytocalcite Group. Monoclinic. Hardness 4

1233\*282. Barytocalcite. BaCO<sub>3</sub>.CaCO<sub>3</sub>. Monoclinic, prismatic by extension of pyramids. 1.50

1234 massive. 1.00

—Hardness 3—3·5

283. Bismutosphärite. Bi<sub>2</sub>(CO<sub>3</sub>)<sub>3</sub>.2Bi<sub>2</sub>O<sub>3</sub>. Spherical, concentric radio-fibrous.

4. Parisite Group. Hexagonal. Hardness 4.5

284. I. Parisite. [(Ce,La,Di)F]<sub>2</sub>Ca(CO<sub>3</sub>)<sub>2</sub>. Hexagonal, acute

double hexagonal pyramids o, and base c, brownish-yellow.


pyramid r, with o, s and a, sharp.

6.00

1236 II. Cordylite. (BaF) (CeF) Ce(CO<sub>3</sub>)<sub>3</sub>. Hexagonal, minute prisms, yellowish. 5.00

Kischtimite. Ce metals fluocarbonate.

Massive, dark brownish-yellow.



	COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type Species No. No.	•
II	I. Synchisite. CeFCa(CO <sub>3</sub> ) <sub>2</sub> . Rhombohedral, rhombio
	habit, wax-yellow.
12370285	i. Bastnäsite. $(Ce, La, Di)_2C_3O_9(Ce, La, Di)F_3$ . Massive,
	brown. 4.00
	Weibyeite. Ce metals fluocarbonate. Orthorhombic,
	minute pyramids.
II	I. Ancylite. 4Ce(OH)CO <sub>3</sub> .3SrCO <sub>3</sub> .3H <sub>2</sub> O. Orthorhombic,
	small pyramids, curved faces, orange-yellow.
5.	Phosgenite Group. Chlorocarbonates. Hardness 3
1238+286	6. Phosgenite. PbCO <sub>3</sub> .PbCl <sub>2</sub> . Tetragonal, prismatic, ada-
3-	mantine, translucent, perfect. 1.25
1239	tabular   c, transparent, sharp. 2.00
12400	minute highly modified prisms with laurionite, in
	ancient slag, limpid, sharp75
1241	cleavage, transparent75
12420	crystalline mass. 1.25
	I. Northupite. MgCO <sub>3</sub> .Na <sub>2</sub> CO <sub>3</sub> .NaCl. Isometric, ideal octa-
	hedron o, gray75
1244	ditto, translucent pale-yellow. 1.00
11	I. Tychite. 2MgCO <sub>3</sub> .2Na <sub>2</sub> CO <sub>3</sub> .Na <sub>2</sub> SO <sub>4</sub> . Isometric, octahedron, colorless.
	B. Acid, Basic and Hydrous Carbonates Hardness 1.5
287	7. Teschemacherite. (NH <sub>4</sub> ) <sub>2</sub> CO <sub>3</sub> .H <sub>2</sub> CO <sub>3</sub> . Orthorhombic.
	Hardness, Malachite, Azurite 3.5-4, others 2-3
	Kalicine. Potassium bicarbonate.
12450288	3. Malachite. 2CuO.CO <sub>2</sub> .H <sub>2</sub> O. Monoclinic, acicular prisms
1245~200	adamantine, bright green. 1.50
1246+	capillary tufts75
1240	velvety crust. 2.00
1247	radiated in chrysocolla, polished. 3.00
1240	radiated in chrysocolia, polished. 3.00

tuberose, smooth, radiated fibrous. 2.00 concentric banded, polished. 2.00

massive. 1.25

1249+

1250° 1251

```
ACID, BASIC, AND HYDROUS CARBONATES
Type Species
No. No.
1252+289. Azurite, Chessylite. 3CuO.2CO<sub>2</sub>.
                   H<sub>2</sub>O. Monoclinic, tabular
                   || c, highly modified (fig.),
                   perfect, adamantine,
                   prussian-blue.
                                    1.00
              prismatic || axis b, modified, sharp. 1.50
12539
              rhombic aspect, symmetrical. 1.50
12540
12550
              ball of crystals. 2.00
              drusy incrustation. .75
1256
              tuberose, concentric radiated, azure-blue. 1.25
1257
1258+
              massive.
              massive with chrysocolla, polished. 3.00
1259
1260
              banded with malachite, polished. 4.00
1261*
              altered to malachite. 1.00
              altered to copper, loose. .75
1262
           Zinkazurite. Zn sulphate, Cu carbonate and H<sub>2</sub>O.
                   Small blue crystals.
1263 290. Aurichalcite. 2(Zn,Cu) CO<sub>3</sub>.3(Zn,Cu) (OH)<sub>2</sub>. Monoclinic
                   (?), tufts of minute very thin flat prisms, bright
                   turquois-blue. 1.50
              velvety druse of minute needles, turquois-blue. .75
1264+
12650
              globular on smithsonite, verdigris-green. .75
1266
              fine laminated, sky-blue. 1.00
       II. Rosasite. 2CuO.3CuCO<sub>3</sub>.5ZnCO<sub>3</sub>. Fibrous, greenish-blue.
1267 291. Hydrozincite. 3ZnO.CO<sub>2</sub>.2H<sub>2</sub>O(?). Massive, reniform
                   fibrous crust in detachable concentric layers. 1.50
1268°
             compact. 1.00
       II. Otavite.
                      Basic Cd carbonate. Rhombohedral, minute
                  crystals, whitish.
1269°292. Hydrocerussite. 3PbO.2CO<sub>2</sub>.H<sub>2</sub>O(?). Minute scaly hex-
                   agonal planes pearly coating on lead. 2.00
             filmy coatings on galena. 3.00
1270
                          Na<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.2CO<sub>2</sub>.2H<sub>2</sub>O.
                                                        Monoclinic (?),
1271°293. Dawsonite.
                  tufts of minute radiating needles.
             thin crusts of radiating blades.
```

others 1.5—2.5 1272 294. Thermonatrite. Na<sub>2</sub>CO<sub>3</sub> + H<sub>2</sub>O. Orthorhombic, a whitish incrustation. 1.50 I. Hydrocalcite. CaCO<sub>3</sub>.2H<sub>2</sub>O.

-Hardness Hydro-magnesite 3.5,

- 100 COMPLETE TYPE COLLECTION. DANA'S SYSTEM
  Type Species
  No. No.
  - 295. Nesquehonite. MgCO<sub>3</sub>+3H<sub>2</sub>O. Orthorhombic, prisms, whitish.
    - 296. Natron. Na<sub>2</sub>CO<sub>3</sub>+10H<sub>2</sub>O. Monoclinic. Occurs in nature only in solution.
- 1273 I. Pirssonite. CaCO<sub>3</sub>.Na<sub>2</sub>CO<sub>3</sub>.2H<sub>2</sub>O. Orthorhombic, hemimorphic, small prisms, perfect, clear, loose. 2.00
  - I. II. Dundasite. PbO.Al<sub>2</sub>O<sub>3.2</sub>CO<sub>2.4</sub>H<sub>2</sub>O. Small spherical aggregates, radio-fibrous, silky whitish.
  - 297. Gay-lussite. CaCO<sub>3</sub>.Na<sub>2</sub>CO<sub>3</sub>+5H<sub>2</sub>O. Monoclinic, elongated || a.
- 1274\* flat wedge-shaped, perfect. 1.00
- 1275°298. Lanthanite. La<sub>2</sub>(CO<sub>3</sub>)<sub>3</sub>+9H<sub>2</sub>O. Orthorhombic, minute thin four-sided plates || base c, whitish. 2.50
  - Hydroconite. CaCO<sub>3</sub>+5H<sub>2</sub>O. Rhombohedral. Recent.
  - 299. Trona. Na<sub>2</sub>CO<sub>3</sub>. HNaCO<sub>3</sub> + 2H<sub>2</sub>O. Monoclinic, tabular, well defined.
- 1276\* efflorescence, whitish, on lava. .40
- 1277\*300. Hydromagnesite. 3MgCO<sub>3</sub>.Mg(OH)<sub>2</sub>+3H<sub>2</sub>O. Monoclinic (?), minute thin clear blades, forming drusy surface of finely foliated crust, silky white. 2.00
- 1278 amorphous chalky. 1.25
  - 301. Hydrogiobertite. MgCO<sub>3</sub>.Mg(OH)<sub>2</sub>+2H<sub>2</sub>O. Spherical.
  - II. Artinite. MgCO<sub>3</sub>.Mg(OH)<sub>2</sub>.3H<sub>2</sub>O. Orthorhombic, radiofibrous, white.
  - II. Giorgissite. 4MgCO<sub>3</sub>.Mg(OH)<sub>2</sub>.4H<sub>2</sub>O. Crusts, white. 302. Lansfordite. 3MgCO<sub>3</sub>.Mg(OH)<sub>2</sub>+21H<sub>2</sub>O. Triclinic, white.
- Hydrodolomite. Hydrated Ca and Mg carbonate. A mixture? Globular, whitish. 2.00
- minute globules (Pennite), incrusting chromite. .75
- 1281+303. Zaratite. 3NiO.CO<sub>2</sub>.6H<sub>2</sub>O. Massive, minute mammillary, vitreous, translucent emerald-green, incrusting chromite. .60
- 1282 ditto, compact massive. .60
  - 304. Remingtonite. Hydrous Co carbonate. Earthy incrustation, rose-colored.
- 1283 305. Tengerite. An yttrium carbonate (?). Pulverulent, thin white coating on gadolinite. 2.50
- 1284°306. Bismutite. Bi<sub>2</sub>O<sub>3</sub>.CO<sub>2</sub>.H<sub>2</sub>O(?). Earthy amorphous, opaque straw-yellow. 1.00
  - Walthérite. Hyd. Bi carbonate. Thin longish crystals, translucent.

Type Species

- 307. Uranothallite. 2CaCO<sub>3</sub>.U(CO<sub>3</sub>)<sub>2</sub>10H<sub>2</sub>O. Orthorhombic, minute, siskin-green.
- 308. Liebigite. CaCO<sub>3</sub>.(UO<sub>2</sub>)CO<sub>3</sub>.20H<sub>2</sub>O. Concretions or coatings, transparent apple-green.
- 1285 309. Voglite. Hydrous U, Ca and Cu carbonate. Rhomboidal scales, pearly green. 4.00
  - Schröckinergite. U hyd. oxycarbonate (?). Orthorhombic(?), six-sided tables, greenish-yellow.
  - Randite. U and Ca hyd. carbonate. Incrustation of microscopic needles, canary-yellow.

#### 2. Silicates

#### A. Anhydrous Silicates

The classification here adopted for the anhydrous silicates cannot be carried through strictly, since there are many species which do not conform to any one of the groups named, and often the true interpretation of the composition is doubtful. Furthermore, within a single group there may be a wide variation in the proportion of the acidic element.

## I. Disilicates. Salts of Disilicic Acid. RSi<sub>2</sub>O<sub>5</sub>.

Polysilicates. Salts of Polysilicic Acid. R<sub>2</sub>Si<sub>3</sub>O<sub>8</sub>.

## Petalite Group. Hardness 6

- 1286°310. Petalite. Li<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.8SiO<sub>2</sub>. Monoclinic, tabular || b, transparent, Castorite, loose. .75
- 1287+ massive. .50
  - Hydrocastorite. Altered castorite. Al, Ca silicate. Mealy mass of fine needles.
- 1288°311. Milarite. H<sub>2</sub>O.K<sub>2</sub>O.4CaO.2Al<sub>2</sub>O<sub>3</sub>.24SiO<sub>2</sub>. Hexagonal, hexagonal prism, perfect, glassy colorless. 3.00
- 1289°312. Eudidymite.  $H_2O.Na_2O.2BeO.6SiO_2$ . Monoclinic, twins, tabular ||c|, well defined, loose (3). .50
- 1290° I. Epididymite. HNaBeSi<sub>3</sub>O<sub>8</sub>. Orthorhombic, tabular || c, elongated || brachydome, colorless. 1.00
  - II. Leucosphenite. 2Na<sub>2</sub>O.BaO.2TiO<sub>2</sub>. 10SiO<sub>2</sub>. Monoclinic, minute crystals, white.

## Feldspar Group. Hardness 6-6.5

Silicates of Al, with either K, Na or Ca, rarely Ba. Besides the distinct species there are intermediate compounds connected by insensible gradations, this close relationship showing in angle, habit, twinning and the various physical and optical characters.

Type Species No. No.	A. MONOCLINIC SECTION
	Orthoclase, Potash Feldspar. K <sub>2</sub> O.Al <sub>2</sub> O <sub>3</sub> .6SiO <sub>2</sub> . Monoclinic, sharply defined perfect crystals:—
1291+	1. Adularia, unit prism m, prism z, clinopinacoid b, with orthodome x and base c united in oscillatory combination, rounded summit(fig.), transparent, large, glassy75
1292	Adularia, Baveno twins, tw.pl. n, 1291. Orthoclase vicinal, coated with chlorite, large, loose. 1.50
12930	Adularia, Moonstone, parting    steep pearly hemiorthodome, transparent50
12940	Adularia, Valencianite, curved crystals, pearly white, aggregate. 1.00
1295+	2. Sanidine, glassy feldspar, 3 to 6 p.c. Na <sub>2</sub> O, twins, thin tabular    b, small transparent, in lava50
1296	Rhyacolite, tabular    $b$ , small glassy. 1.00 Orthoclase 3. Ordinary varieties and forms:—
1297+	unit prism m, clinopinacoid b, orthodome y and base c (fig.), symmetrical, gray50
1298	ditto, with prism z, loose squarish prisms (3)50
12990	ditto, with pyramid o, very perfect, yellowish. 1.00
1300	ditto, lustrous, milky, with tourmaline.
13010	m, z, b, c, y, o with pyramid n and orthopinacoid a (fig.), very large, roughly symmetrical, red-brown. 1.25
1302*	Carlsbad penetration-twin, tw. axis c (fig.), large, rough, gray50 1301. Orthoclase

## FELDSPAR GROUP


Type Species No. No.	Orthoclase—Continued
1303	ditto, very perfect, yellowish, loose40
13049	Carlsbad contact-twin, tw.pl. orthopina- coid a50
1305*	Baveno contact-twin, tw.pl. n (fig.), large squarish prism, loose50
13060	Manebach contact-twin, tw.pl. base $c$ (fig.).
1307	columnar divergent, reddish40
1308+	coarsely cleavable20
13090	granular20 1302. Orthoclase
13100	compact, jaspery, red20
1311	Loxoclase, 7.56 p.c. Na <sub>2</sub> O, $m$ , $z$ , $a$ , $b$ ,
	y, n and c, bluish opalescent, gray75
1312	Necronite, fetid75
	Lazurfeldspar, with lazurite.  Murchisonite, yellow reflections, red.  Weissigite, small twins in amygdaloid.
1313	altered to cassiterite, Carlsbad twin, loose. 1.25
13140	Perthite. An interlamination of albite 1305. Orthoclase
	and orthoclase, cleavage, Aventurine, flesh-red25
1315	Perthite, Cryptoperthite, gray chatoyant. 1.00
1316 314.	Hyalophane. K <sub>2</sub> O.BaO.2Al <sub>2</sub> O <sub>3</sub> .8SiO <sub>2</sub> . Monoclinic, unit
	prism $m$ , clinopinacoid $b$ , orthodome $x$ and base $c$ ,
	sharply defined, clear colorless. 1.50

#### B. Triclinic Section

315. Microcline. K<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.6SiO<sub>2</sub>. Triclinic, large well developed crystals:—
1317\*

1. Ordinary, unit prisms M and m, brachypinacoid b, macrodome x and base c, pale yellowish-gray, with albite. .50
1318 ditto with additional prisms z and f and macrodome y. .50
1319+ squarish perfect cleavage, cream-

yellow. .20




1306. Orthoclase

Type Species	MPLETE TYPE COLLECTION. DANA'S SYSTEM  Microcline—Continued
1320+	Amazonstone, Amazonite, unit prisms $M$ and $m$ , prisms $s$ and $f$ , brachypinacoid $b$ , macrodome $s$ , pyramid $o$ and base $c$ , bright verdigris-green50
1321	Amazonstone, ditto, capped with white albite on macrodome x. 1.50
1322	Amazonstone $M$ , $m$ , $z$ , $b$ , $o$ , $c$ and macrodome $y$ , with polyadelphite75
13230	Amazonstone, <i>Baveno</i> contact-twin, tw.pl. <i>n</i> , loose squarish prism, greenish. 1.00
13249	Amazonstone, Manebach contact-twin, tw.pl. c, bright verdigris-green, loose. 2.00
1325	Amazonstone, broad perfect cleavage, bright and trans- lucent, mottled verdigris-green. 1.00
1326	Amazonstone, massive, precious. 2.00
13279	Chesterlite, rough crystal75
1328 315A.	ANORTHOCLASE. (NaK) AlSi <sub>3</sub> O <sub>8</sub> . Triclinic, twins,
	Manebach law, tabular    c, glassy, perfectly de-
	veloped but microscopic, in obsidian lithophyses.
	·75
13290	Anorthoclase cleavage, near 90°, bluish opalescent, gray. 1.00

## Albite—Anorthite Series

Note—Between the isomorphous species Albite, NaAlSi $_8O_8$  (Ab) and Anorthite, CaAl $_2Si_2O_8$  (An), are several subspecies, regarded as isomorphous mixtures of these molecules (Ab $_n$ An $_m$ ), and defined according to the ratio in which they enter.

1330° <b>316</b> .	Albite, Soda Feldspar. Na <sub>2</sub> O.Al <sub>2</sub> O <sub>3</sub> .6SiO <sub>2</sub> .
	Triclinic, unit prisms $M$ and $m$ ,
	macrodome x in oscillatory com-
	bination with base $c$ , on rock-crystal50
13310	twin, tw.pl. b, albite law, (fig.), perfect, small, translucent50
1332	parallel grouping, curved40
1333*	cleavage, well marked polysynthetic twinning striæ, albite law20
1334	massive granular20



Type Species No. No.	Albite—Continued
No. No. 1335°	Peristerite, contact-twin, Carlsbad law, tw. axis c,
<b>*333</b> °	small, perfect, clear faintly iridescent. 1.00
	Aventurine, Sunstone.
1336	Moonstone, cleavage, flesh-colored50
1337	Moonstone, cleavage, well marked polysynthetic twin-
*331	ning striæ, albite law, pearly-gray. 1.00
1338+	Pericline, unit prism M and m,
	brachypinacoid b, macrodome
	x and base c (fig.), ideal de-
	velopment, opaque milky.
	1.00
1339°	Pericline, contact-twin, tw. axis
	b, pericline law, perfect75 1338. Albite
	Hyposclerite, contains 5 p.c. pyroxene, blackish-green.
1340+	Cleavelandite, lamellar, curved divergent, white20
	Olafite and Tschermakite are oligoclase-albite.
317. O	LIGOCLASE. Al, Na and Ca polysilicate (intermediate
	between albite and anorthite, Ab <sub>3</sub> An <sub>1</sub> ). Triclinic,
	crystallized.
1341*	cleavage, grayish-white40
13420	massive, subtransparent, whitish75
1343	Aventurine, Sunstone, squarish perfect cleavage, pale
	greenish-gray75
1344	ditto, cleavable-granular, reddish-gray40
1345+	ditto, cleavage, well marked polysynthetic twinning
	striæ, translucent brownish-red, striking golden
	fiery reflections of included microscopic crystals
	(hematite or göthite?), precious75
1346 318. A	NDESINE, Andesite, Al, Na and Ca polysilicate (intermedi-
	ate and between albite and anorthite, Ab <sub>3</sub> An <sub>2</sub>
	to Ab <sub>1</sub> An <sub>1</sub> ). Triclinic, highly modified, loose75
1347*	porphyritic phenocrysts, well defined, whitish30
1348 319. L	ABRADORITE. Al, Na and Ca polysilicate (intermediate
	between albite and anorthite, Ab <sub>1</sub> An <sub>1</sub> to Ab <sub>1</sub> An <sub>3</sub> ).
	Triclinic, twin, very thin tabular. 2.00
1349+	cleavage, well marked polysynthetic twinning striæ, albite
	law, blue and green chatoyancy, gray, precious30
1350	ditto, with also yellow, red and bronze in the play of
	colors, polished. 1.25
13519	compact massive30

106	COMPLETE	TYPE	COLLECTION.	DANA'S	SYSTEM
Type	Species				

Maskelynite (Meteoric). Al<sub>2</sub>O<sub>3</sub> 25.7 p.c., CaO 11.6 p.c., Na<sub>2</sub>O 5.1 p.c., K<sub>2</sub>O 1.3 p.c., SiO<sub>2</sub> 56.3 p.c.=100. Isometric, distorted cubic (?) grains, transparent colorless.

1352+320. Anorthite. CaO.Al<sub>2</sub>O<sub>3.2</sub>SiO<sub>2</sub>. Triclinic, small, highly modified, clear glassy, in lava. 1.00

1353° penetration-twin, Carlsbad law, tw. axis c, rough, gray, loose. .75

1354 Indianite, granular. 1.00
Amphodelite and Latrobite are reddish.

1355° Cyclopite, minute, thin tabular || b, transparent, in lava. 1.00

Tankite, cleavable, grayish.

- I. Celsian. BaAl<sub>2</sub>Si<sub>2</sub>O<sub>8</sub>. Triclinic, massive, colorless.
- II. Paracelsian.

Barsowite. CaO.Al<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>(?). Orthorhombic (or monoclinic). Partly altered anorthite (?). Pearly white.

## II. Metasilicates. Salts of Metasilicic Acid. RSiO<sub>3</sub>.

## 1. Leucite Group. Isometric. Hardness 6 and 6.5

1356\*321. Leucite. K<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.4SiO<sub>2</sub>. Isometric at 500° C., pseudo-isometric ordinarily, form near trapezo-hedron *n*, ideal symmetry (fig.), slightly altered, gray, loose. .30

1357° ditto, unaltered in lava. .75

1358+ ditto, small, subtransparent, loose (24). .30

ditto, opaque snow-white. 1.00

1360 altered to feldspar. .50

1361° altered to nephelite-feldspar mixture, Pseudoleucite.

1362 altered to kaolin. .50

1363 322. Pollucite. H<sub>2</sub>O.(Cs, Na)<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.5SiO<sub>2</sub>. Isometric, cube a and trapezohedron n, loose. 5.00

1364° massive, translucent, vitreous. 3.00



### 2. Pyroxene Group. RSiO<sub>3</sub>.

The orthorhombic, monoclinic and triclinic species of this group are closely related in optical and physical properties and in angles. All have a fundamental squarish prism with an angle of 93° and 87°, with cleavage parallel to this prism. The metasilicates of Ca, Mg and Fe III are prominent.

These species are closely related to a parallel chemical series formed by the corresponding orthorhombic, monoclinic and triclinic members of the Amphibole Group. In several cases the same chemical compound appears in both groups.

The chief external distinctions between pyroxene and amphibole proper are: prismatic angle of pyroxene 87° and 93°; amphibole 56° and 124°, with more perfect prismatic cleavage. Pyroxene prisms usually short and often complex, massive forms mostly lamellar or granular. Amphibole prisms generally long and simple, columnar and fibrous types predominating.

#### A. ORTHORHOMBIC SECTION.

Type Species No. No.	Hardness 5.5
No. No. 323.	Enstatite. MgO.SiO <sub>2</sub> . Orthorhombic.
	1. Iron-free, prismatic.
1365	massive, in meteorite. 8.00
13660	2. Ferriferous, Bronzite, cleavage, metalloidal, brownish75
1367	ditto, lamellar, greenish-brown30
1368+	ditto, sublamellar, gray30
1369	ditto, fine fibrous, gray75
13700	altered to steatite, sharply developed large prism $m$ , pinacoids $a$ and $b$ , but roughly terminated50
224	•
1371 324.	Hypersthene. (Fe,Mg)O.SiO <sub>2</sub> . Orthorhombic, small prisms. 3.00
1372*	cleavage, metalloidal schiller, brownish-black. 1.50
1373°	granular-cleavable, black. 1.00
	Amblystegite.
	Szaboite, tabular $  b$ , slightly altered.
	Alteration-products of enstatite-hypersthene:—Diaclasite,
	Bastite (Schiller Spar), Phästine.

#### B. MONOCLINIC SECTION

Range of Hardness 5.5-6.5 (Wollastonite and Pectolite 5)

Type Species
No. No. 325. Pyroxene. Mainly RSiO<sub>3</sub> (see leading varieties). Monoclinic and hemihedral forms described under the chemically classified varieties:-

#### I. VARIETIES CONTAINING LITTLE OR NO ALUMINIUM

1.	VARIETIES CONTAINING LITTLE OR NO ALUMINIUM
1374°	DIOPSIDE. CaMg(SiO <sub>3</sub> ) <sub>2</sub> . Unit prism m, orthopinacoid a, clinopinacoid b, pyramid u and base c (similar to fig.), perfect, transparent pale green, loose75
1375°	m, a, b, c, orthodome p, pyramids u, s and λ (similar to fig.), large, well developed, subtranslucent. 1.00
1376	slender prisms, transparent pale green, with cinnamon garnet. 1.50
1377	rough prisms, subtransparent, dark green, loose (12)50
1378	columnar crystalline, buff75
1379+	granular, olive-green50
	The following belong here:—
	Chrome-diopside, some Cr, bright green.
1380*	Malacolite, large rough prisms, yel-
	lowish-white, loose (6)30   m   a   m
1381	Alalite, squarish prisms, pale greenish.
	2.00
13820	Mussite, long flat implanted prisms,
	pale greenish-gray. 1.00
1383	Traversellite, long prisms, pale greenish. 1.50
	Canaanite, massive, whitish.
	Lavrovite, contains V, granular, em-
	erald-green.
1384*	HEDENBERGITE. CaFe(SiO <sub>3</sub> ) <sub>2</sub> . Prisms  m and $\chi$ , orthopinacoid a, clinopinacoid b, orthodome p and pyramids u and o (fig.), sharply developed, perfect, brilliant

blackish. 1.50

1397 S. Anomalite, Mn-Ni-Co-Pyroxene, light like pumice. 1.50


## II. ALUMINOUS VARIETIES

Augite. Chiefly CaMgSi<sub>2</sub>O<sub>6</sub> with (Mg,Fe) (Al,Fe)<sub>2</sub>Si<sub>2</sub>O<sub>6</sub> and occasionally alkalies:—

(a) Leucaugite. Contains Al, Ca, Mg, whitish.

13980

(b) Fassaite, pyramidal (fig.), pale green. 1.00



Type Species No. No.	COMPLETE TYPE COLLECTION. DANA'S SYSTEM Pyroxene—Continued
1399°	(c) Augite, minute pyramidal, bright, perfect, translucent green, in lava.
1400+	Augite, unit prism m, orthopinacoid a, clinopinacoid b and pyramid s (fig.), highly symmetrical, sharp, black, in volcanic tuff50
1401	ditto, with base $c$ 50
1402*	Augite, contact-twin, tw.pl. a (fig.). sharply developed, loose30
1403	Augite, penetration-twin75
1404	Titaniferous Augite, 0.5-4.5 p.c. TiO275
1405	Alkali-augite. 1 to 10 p.c. Na <sub>2</sub> O75
1406	Fassaite altered to talc. 1.00
1407°	Augite altered to cimolite, symmetrical, complete, loose30
1408	Augite altered to serpentine75
	Alteration-products of Pyroxene:—Hec-
	torite, Monradite, Pitkärantite,
	Hydrous diallage, Pyrallolite, 1402. Augite
	Strakonitzite.
1409	Picrophyll, lamellar, shiny grayish-green75 Uralite. See amphibole.
1410+326.	Acmite. Na <sub>2</sub> O.Fe <sub>2</sub> O <sub>3</sub> .4SiO <sub>2</sub> . Monoclinic, acutely termi-
	nated large slender prism, vertically channeled,
	black. 50
14110	Ægirite, prismatic $  $ axis $c$ , unit prism $m$ and orthopinacoid
	a prominent, sharply symmetrical, loose, large.
_	1.00
	Urbanite. (Ca,Mg)SiO <sub>3</sub> +2NaFe <sup>III</sup> (SiO <sub>3</sub> ) <sub>2</sub> . Monoclinic, pyramidal.
1412°3 <b>27</b> .	Spodumene. Li <sub>2</sub> O.Al <sub>2</sub> O <sub>3</sub> .4SiO <sub>2</sub> . Monoclinic.
	<ol> <li>Ordinary. Flat rough prisms, highly modified, loose, large. 1.25</li> </ol>
	cleavage, clear pale yellowish-green, precious. 1.00
1413°	cleavage, perfect, opaque white20
1413° 1414+	cicavage, perfect, opaque winte20
414+	2. Hiddenite, slender prisms, deeply etched, transparent emerald-green, precious, loose. 2.50
1414+ 1415	2. Hiddenite, slender prisms, deeply etched, trans-

#### PYROXENE GROUP Spodumene—Continued

Type No.	Species
No.	No.

Alteration-products of spodumene:—

1st stage,  $\beta$  spodumene, Na replacing  $\frac{1}{2}$  of original Li, compact subfibrous, whitish.

2d stage, Cymatolite or Aglaite, subfibrous, silky whitish. .75

Killinite, compact, cryptocrystalline, greenish.

1418+328. Jadeite. Na<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.4SiO<sub>2</sub>. Monoclinic (or triclinic), splintery compact, translucent leek-green, precious. 2.50

1419 ditto, greenish-gray. 1.50

1420° ditto, white spotted with green, cut. 3.00

Chloromelanite. 6.06—10.59 p.c. Fe<sub>2</sub>O<sub>3</sub>, blackish.

JADE is a popular term for jadeite, as well as the commoner nephrite, etc.

1421 329. Wollastonite. CaO.SiO<sub>2</sub>. Monoclinic, tabular || c, rough, large. 2.00

slender prisms, translucent, in lava. 1.00

1423+ radio-fibrous, whitish. .75

1424°330. Pectolite. H<sub>2</sub>O.Na<sub>2</sub>O.4CaO.6SiO<sub>2</sub>. Monoclinic, acicular, aggregated, strongly triboluminescent. 1.50

1425° capillary, aggregated, white. 1.00

long fibrous, radiated. .50

1427+ mammillary, radio-fibrous, white. .50

Osmelite, columnar radiated. Walkerite, 5·12 p.c. MgO. compact jade-like, pale green.

1428 Manganpectolite. 4.25 p.c. MnO, cleavage, gray. 1.00

1429 altered to quartz. .75

1430 331. Rosenbuschite. 6CaSiO<sub>3</sub>.2Na<sub>2</sub>ZrO<sub>2</sub>F<sub>2</sub>.(TiSiO<sub>3</sub>TiO<sub>3</sub>). Monoclinic, crystalline, light orange-gray. 2.50

#### C. Triclinic Section. Hardness 5.5—6

1433 334. Hiortdahlite. Nearly corresponds to 4Ca(SiZr)O<sub>3</sub>.Na<sub>2</sub> ZrO<sub>2</sub>F<sub>2</sub>. Triclinic, tabular || a, light yellow. 3.00

<sup>1431 332.</sup> Lavenite. (Na<sub>4</sub>, Ca<sub>2</sub>, Mn<sub>2</sub>, Zr) ([Si, Zr]O<sub>3</sub>)<sub>2</sub>. Monoclinic, prismatic, yellowish. 4.00

<sup>1432°333.</sup> Wöhlerite.  $12R(Si,Zr)O_3.RNb_2O_6$ , with R=Ca: Na<sub>2</sub>=4:1. Monoclinic, tabular || a, resin-yellow. 1.00

I. Hainite. Contains Ti, Zr, Na, Ca. Triclinic, slender needles.

	COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type Species No. No.	
335.	Rhodonite. MnO.SiO <sub>2</sub> . Triclinic. 1. Ordinary:—
1434*	Paisbergite, prisms M and m, brachypina-
	coid b, pyramid k (fig.), sharp, pink. 1.00
14350	granular massive, deep rose-pink50
1436	ditto, translucent slab, polished. 1.50
1437+	compact, pale rose-red35
	2. Ferriferous.
1438	3. Calciferous, Bustamite, 9 to 20 p.c. CaO,
	gravish-red 200
1439°	4. Zinciferous, Fowlerite,
107	5 to 7 p.c. ZnO, prisms
	M and m, macropinacoid
	a, brachypinacoid b,
	pyramids k and n and
	base c (fig.), rough tabu-
	lar    c, large, dull pink.
	2.00
1440	ditto, pyramids $q$ and $r$ addi-
-440	tional, translucent bright 1439. Rhodonite
	cherry-red, in white calcite. 3.00
14410	ditto, imperfect rounded, opaque, rose-pink75
1442	ditto, rounded squarish prism, large, pale salmon-red.
1440	1.50
1443+	ditto, cleavage, rose-pink35
	Alteration-products of rhodonite:-Marceline, Dyssnite,
	Stratopeite, Allagite and Photicite.
	Hydrorhodonite. A hydrated rhodonite?
1444°336.	Babingtonite. (Ca, Fe, Mn)SiO <sub>3</sub> with Fe <sub>2</sub> (SiO <sub>3</sub> ) <sub>3</sub> . Triclinic,
,	small well defined tables, bright, black. 5.00

## 3. Amphibole Group. Range of Hardness 5-6 (Crocidolite 4)

light red.

II. Schizolite. HNa(Ca,Mn)<sub>2</sub>(SiO<sub>3</sub>)<sub>3</sub>. Triclinic, prismatic,

Composition RSiO<sub>3</sub> with R=Ca,Mg,Fe chiefly, also Mn,Na<sub>2</sub>(K<sub>2</sub>), H<sub>2</sub>. Further often containing aluminium and ferric iron, in part as NaAl(SiO<sub>3</sub>)<sub>2</sub> or NaFe(SiO<sub>3</sub>)<sub>2</sub>; perhaps also as  $R^{II}R^{III}_{2}SiO_{6}$ .

The orthorhombic, monoclinic and triclinic members of this group are closely related optically, chemically and in form, and have a common prismatic cleavage of 54° to 56°. See Pyroxene Group. (Species No. 325).

Type Species No. No.	A. Orthorhombic Section.
No. No. 1445 337.	Anthophyllite. (Mg,Fe)SiO <sub>3</sub> . Orthorhombic, lamellar-fibrous, grayish-green60
1446	radio-fibrous, stellate, gray40
1447+	long fibres, asbestiform, brownish60 Kupfferite.
1448°	aluminous, Gedrite, bladed-granular, blackish60 Thalackerite, metalloidal. Valléite. RSiO <sub>3</sub> with R = Mg, Ca, Fe, Mn. Orthorhombic, prisms, reddish.

### B. Monoclinic Section

338. Amphibole. Composition in general analagous to the pyroxenes. See leading varieties. Monoclinic.

	pyroxenes. See reading varieties. Atomosimist
	I. CONTAINING LITTLE OR NO ALUMINIUM
14490	TREMOLITE. CaMg <sub>3</sub> (SiO <sub>3</sub> ) <sub>4</sub> . Unit prism m, clinopinacoid
	b and clinodome $r$ (fig.), grayish-white. 1.00
1450	prismatic, transparent, pale asparagus-green. 1.50
1451	prismatic long-bladed75
14520	thin columnar, pale gray75
1453+	cleavage, prismatic, greenish-gray40
1454	fibrous, gray75
14550	radio-fibrous, stellated, white75
1456+	Hexagonite, 1.37—2.39 p.c. MnO, sublam-
	ellar, luminesces red with sharp friction,
	lavender75
1457+	ACTINOLITE. Ca(Mg,Fe) <sub>3</sub> (SiO <sub>3</sub> ) <sub>4</sub> . Prismatic,
	long bright blades, dark green, in talc.
	.40
1458*	columnar bladed, light green30 . 1449. Amphibole
1459°	fibrous, blackish-green50
1460	radio-fibrous, greenish75
1461	granular massive, green40
1462+	Nephrite, Jade (see also jadeite), splintery compact,
	translucent whitish (tremolite)75
1463	ditto, dark green (actinolite), precious. 1.25
14640	Asbestus (see also chrysotile, a variety of serpentine),
	fine loose silky threads, white, Amianthus40
1465+	fibrous, grayish20

Type Species Amphibole—Continued No. No.
1466 long cohering fibres, brownish30
long cohering fibres, greenish30
1468* Mountain leather, flexible sheets, grayish50
1469 <sup>®</sup> Mountain cork, floats easily, yellowish75
1470° Mountain wood, compactly fibrous, brownish50
Byssolite, matted capillary, green, with epidote etc. 1.50
1472 <sup>9</sup> Byssolite, ditto, in calcite cleavage40
Byssolite, ditto, felt-like mass20
1474 SMARAGDITE, thin-foliated, light grass-green50
1475° URALITE, an altered pyroxene, sharply defined squarish
prisms, green. 1.50
1476° CUMMINGTONITE. (Fe, Mg)SiO <sub>3</sub> . Radiated fibro-lamellar
grayish-brown50
DANNEMORITE. (Fe, Mn, Mg) SiO <sub>3</sub> . Here belong Asbe
ferrite (asbestiform), Silfbergite, Hillängsite (like
anthophyllite). Grüneriте. FeSiO₃. Fibro-lamellar, silky.
1477° RICHTERITE. ([K,Na] <sub>2</sub> MgCaMn)SiO <sub>3</sub> , long crystals. 1.00
1478° Breislakite, wool-like, dark-brown, on lava. 1.00
1479 S. ASTOCHITE. (Mg, Mn, Ca) SiO <sub>3</sub> combined with (Na, K, H)
SiO <sub>3</sub> . Short columnar aggregates. 1.25
MARMAIROLITE, fine needles, pale yellow.
II. ALUMINOUS VARIETIES
Chiefly Ca(Mg,Fe) <sub>3</sub> Si <sub>4</sub> O <sub>12</sub> with Na <sub>2</sub> Al <sub>2</sub> Si <sub>4</sub> O <sub>12</sub> and (Mg,Fe) <sub>2</sub> (Al,Fe)
Si <sub>2</sub> O <sub>12</sub> (color darkens with the increase of Fe):—
1480* EDENITE, Al-Mg-Ca- amphibole, cleavage, pale green30
1481 fibro-lamellar, grayish50 1482° PARGASITE, crystalline, green75
1482° PARGASITE, crystalline, green75 1483° Common Hornblende, unit
prism $m$ , clinopinacoid $b$
and clinodome $r$ , sharply
symmetrical, (fig 1449.), 1485. Hornblende
large, black, with brown apatite. 1.50
1484+ $m, b, r$ and orthodome $p$ , black, loose30
1485 ditto, tabular (fig.), in calcite. 1.50
1486° terminated blades, bright, black in lava75
non-terminated blades, bright black, large. 1.50
1488+ cleavable granular, black20
1489 cleavage, greenish-black50

Type Species No. No. 1490 Amphibole—Continued

granular, black. .50

Noralite, contains only 2.25 p.c. MgO.

Gamsigradite, contains 6 p.c. MnO.

Bergamaskite, contains only 0.93 p.c. MgO.

Kaersutite, 6.75 p.c. TiO<sub>2</sub>, prismatic, black.

- II. Soretite, short prisms.
- II. Szichenyite, greenish.
- I. Hastingsite, grains in nephelite-syenite.
- I. Xiphonite, minute light honey-yellow crystals in lava.

1491° Tremolite altered to talc, fibrous, pearly white. .75

Tremolite altered to talc, bladed, grayish-green. .75

Altered amphiboles:—Kirwanite, Loganite, Paligorskite, Phäactinite, Waldheimite.

- 1493°339. Glaucophane. Essentially NaAl(SiO<sub>3</sub>)<sub>2</sub>. (Fe, Mg)SiO<sub>3</sub>.

  Monoclinic, indistinct prisms, embedded, bluish-black. .60
- 1494+ fine columnar-granular, bluish-gray. .60
  - I. Rhodusite. Fe,O, replaces Al,O,.
  - I. Crossite. Chiefly Fe, Mg, Na, Ca and Al metasilicate, monoclinic, lath-shaped crystals, blue.
- 1495\*340. Riebeckite. 2NaFe<sup>III</sup>(SiO<sub>3</sub>)<sub>2</sub>.FeSiO<sub>3</sub>. Monoclinic, embedded prisms, black. .75
- 1496°341. Crocidolite. NaFe<sup>III</sup>(SiO<sub>3</sub>)<sub>2</sub>.FeSiO<sub>3</sub>. Asbestiform, long fibrous, silky lavender-blue. .25

1497+ altered to quartz, Tiger-eye, chatoyant golden-brown. .40
1498 ditto, blue. .40

1499 ditto, golden-brown and blue, polished. 1.00.

1500\*342. Arfvedsonite. 4Na<sub>2</sub>O.3CaO.14FeO.(Al,Fe)<sub>2</sub>O<sub>3</sub>.21SiO<sub>2</sub>.

Monoclinic, long prism, loose. 1.25

I. Cataphorite. An alkali-iron amphibole.

342A. Barkevikite. Ratio of SiO<sub>2</sub>: (Al,Fe)<sub>2</sub>O<sub>3</sub>: (Fe,Mn,Ca,Mg) O: (NaK)<sub>2</sub>O=0.707: 0.148: 0.498: 0.113. Large rough prisms, deep velvet-black.

#### C. Triclinic Section

1501°343. Ænigmatite. Nearly 2Na<sub>2</sub>O. 9FeO.AlFeO<sub>3</sub>.12(Si,Ti)O<sub>2</sub>.
Triclinic, prisms, black. 2.50

Cossyrite, minute embedded crystals.

II. Rhönite.  $(Ca,Na_2K_2)_3Mg_4Fe_2^{11}Fe_3^{111}Al_4(Si,Ti)_6O_{30}$ . Triclinic.

# 4. Beryl Group.

Type Species No. No.	Hexagonal. Hardness 7.5—8
No. No. 344.	Beryl. 3BeO.Al <sub>2</sub> O <sub>3</sub> .6SiO <sub>2</sub> . Hexagonal. Transparent
	varieties are precious.
1502	1. Emerald, colored by $Cr_2O_3$ , large symmetrical unit prism $m$ and base $c$ , translucent emerald-green,
	loose. 2.50
1503+	ditto, embedded prisms. 1.00
1504	ditto, transparent prism, bright. 9.00
	2. Ordinary varieties:—
15050	(a) colorless transparent, unit prism m,
	unit pyramid $p$ , diagonal pyramid $s$ and base $s$ (similar to fig.), $s$
	small, sharply developed, with
	tourmaline, etc. 1.50
1506	(b) bluish-green transparent, Aqua-
-0	marine, slender prism m, highly
	modified termination. 5.00
1507*	ditto, massive. 1.25
1508∘	(c) apple-green, subtranslucent, unit
	prism m and base c (fig.), symmet-
	rical, large, loose75 m m m
15090	ditto, long prism parted into cross-sec-
	tions, the interstices being filled with quartz (fig.). 1.25
1510+	ditto, massive35
1511	(d) honey-yellow, golden beryl, trans-
1311	parent prism. 2.50
	(e) pale yellowish-green.
	(f) clear sapphire-blue.
1512	(g) pale sky-blue, Blue Aquamarine, trans-
•	parent, massive. 2.00
15130	(h) pale rose-red, transparent prism,
	small, sharply developed, with tour-
	maline, etc. 1.50
1514	ditto, tabular $  $ base $c$ , modified, larger.
	7.00 1509. Beryl
15150	(i) brownish-yellow, waxy, semi-opaque,
	massive35

Type Species
No. No.

1516

Beryl-Continued

- 3. Caesium Beryl, contains Cs, unit prism m, diagonal pyramid s, translucent pale pink, large. 4.00 Altered beryl, Rosterite, Pseudosmaragd.
- 5. Eudialyte Group. Range of Hardness 5-6
- 1517\*345. Eudialyte. Na<sub>13</sub>(Ca,Fe)<sub>6</sub>Cl(Si,Zr)<sub>20</sub>O<sub>52</sub>. Rhombohedral, optically positive, diagonal prism a, rhombohedrons r and z and base c (fig.), brownishred. 1.25
- 1518° Eucolite. Optically negative. Rhombohedron e prominent, rose-red.

  1.25
- 1519°346. Catapleiite. H<sub>2</sub>(Na<sub>2</sub>, Ca)(Zr(OH)<sub>2</sub>)
  (SiO<sub>3</sub>)<sub>3</sub>. Hexagonal at 1400 C°.,
  ordinarily pseudohexagonal and
  monoclinic, thin tabular hexagonal prisms with replaced edges, yellow. 1.00

1520 I. Elpidite. Essentially Na<sub>2</sub>O.ZrO<sub>2</sub>.6SiO<sub>2</sub>.3H<sub>2</sub>O. Orthorhombic, small prisms, whitish. 2.50

Natron-catapleiite, without Ca, bluish-gray.

- 6. Melanocerite Group. Range of Hardness 5-6
- 347. Cappelenite. 3BaSiO<sub>3</sub>.2Y<sub>2</sub>(SiO<sub>3</sub>)<sub>3</sub>.5YBO<sub>3</sub>. Hexagonal, thick prisms.
- 348. Melanocerite. Hypothetically 12(H<sub>2</sub>Ca)SiO<sub>3</sub>.3(Y,Ce) BO<sub>3</sub>.2H<sub>2</sub>(Th,Ce)O<sub>2</sub>F<sub>2</sub>.8(Ce,La,Di)OF. Rhombohedral, tabular.
- 1521 349. Caryocerite. 6(H<sub>2</sub>Ca)SiO<sub>3.2</sub>(Ce,Di,Y)BO<sub>3.3</sub>H<sub>2</sub>(Ce,Th)O<sub>2</sub> F<sub>2.2</sub>LaOF. Rhombohedral, tabular rhombohedrons. 5.00
- 1522 Steenstrupine. Essentially a metasilicate of Th, Ce metals, Fe and Na, with some Al, Mn and Ca. Rhombohedral, tabular rhombohedrons, sharply developed, brownish-black. 3.00
  - 350. Tritomite. 2(H<sub>2</sub>Na<sub>2</sub>Ca)SiO<sub>3</sub>.(Ce,La,Di,Y)BO<sub>3</sub>.H<sub>2</sub>(Ce,Th, Zr)O<sub>2</sub>F<sub>2</sub>. Rhombohedral, acute triangular pyramidal.
- 1523 massive, disseminated, resinous dark-brown. 5.00

#### II. Intermediate Silicates

### 1 Leucophanite Group.

Range of Hardness 4-5

Type Species 1524 351. Leucophanite. Na(BeF)Ca(SiO<sub>3</sub>)<sub>2</sub>. Orthorhombic, tabular  $\parallel$  base c. 3.00

cleavage, greenish-white. 1.50 15250

> 352. Meliphanite. NaCa<sub>2</sub>Be<sub>2</sub>FSi<sub>3</sub>O<sub>10</sub>. Tetragonal, tetartohedral, obtuse pyramid p prominent.

crystalline lamellæ, honey-yellow. 15260

- II. Taramellite. 4BaO.FeO.2Fe<sub>2</sub>O<sub>3</sub>.10SiO<sub>2</sub>. Orthorhombic (?), fibrous, reddish-brown.
- II. Weinbergerite. NaAlSiO<sub>4</sub>+3FeSiO<sub>3</sub>(?). Spherical aggregates in meteoric iron.

#### 2. Iolite Group. Hardness 7-7.5

1527 353. Iolite, Cordierite. H<sub>2</sub>O.4(Mg,Fe)O.4Al<sub>2</sub>O<sub>3</sub>.10SiO<sub>2</sub>. Orthorhombic, twins, short pseudohexagonal prisms, dark smoky-blue. 2.00

massive, translucent dark blue. 15280

coarse granular, pale blue, with pale greenish chlorophyl-1529+

Cerasite, contains regularly arranged inclusions.

Alteration-products of iolite:—(See mica group for the alkaline kinds), Bonsdorfite, Auralite, Pyrargillite, Esmarkite, Praseolite, Raumite, Peplolite, Aspasiolite, Polychroilite. also:-

Fahlunite, dark brown. .75 1530

Chlorophyllite, greenish-gray. .50 1531

## 3. Barysilite Group. Hardness 3 (except Hyalotekite 5-5.5)

- 3PbO.2SiO<sub>2</sub>. Hexagonal, curved lamellar, 15320354. Barysilite. pearly white, tarnishing. 1.50
  - 355. Ganomalite. 3PbO.2(Ca,Mn)O.3SiO<sub>2</sub>. Tetragonal, prismatic.
- granular massive, resinous grayish, with manganophyl-15330 lite. 1.50


Type Species

- 1534 I. Hardystonite. 2CaO.ZnO.2SiO<sub>2</sub>. Tetragonal, rough indistinct crystal. 3.00
- 1535° compact massive, white, with franklinite and willemite.
- 1536\* granular massive, pale yellow, with polyadelphite. .50
  356. Hyalotekite. Approximately Ca<sub>3</sub>Ba<sub>3</sub>Pb<sub>3</sub>B<sub>2</sub>(SiO<sub>3</sub>)<sub>12</sub>. Massive, coarsely crystalline.

## III. Orthosilicates. Salts of Orthosilicic Acid. R<sub>2</sub>SiO<sub>4</sub>.

1. Nephelite Group. Hexagonal. Hardness 6

1537°357. Nephelite. 3Na<sub>2</sub>O.K<sub>2</sub>O.4Al<sub>2</sub>O<sub>3</sub>.9SiO<sub>2</sub>. Hexagonal, unit prism *m*, diagonal prism *a*, unit pyramid *p* and base *c* (fig.), small but sharply defined, glassy colorless, in lava.



1538° unit prism *m* and base *c*, minute, ideal symmetry, glassy pale gray, with melilite. 1.00

1539 Elæolite, coarse crystals. 1.00

1540 Elæolite, massive, greasy brown. .40

1541+ Elæolite, massive, greasy pale grayish, in "litch-fieldite" (nephelite-syenite). .40

Alteration-products of nephelite:—See Pinite, Gieseckite, Dysyntribite, Liebenerite Lythrodes.

- 358. Eucryptite. Li<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>. Hexagonal, embedded microscopic crystals, clear colorless.
- 359. Kaliophilite. K<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>. Hexagonal, slender acicular and cobweb-like threads, silky colorless.
- 360. Cancrinite. 3H<sub>2</sub>O.4Na<sub>2</sub>O.CaO.4Al<sub>2</sub>O<sub>3</sub>.9SiO<sub>2</sub>.2CO<sub>2</sub>. Hexagonal, unit prism m and unit obtuse pyramid p.
- massive, orange-yellow, in "litchfieldite" (nephelitesyenite). .75

massive, blue. 1.50

Kalk-cancrinite. SiO<sub>2</sub> 39·82, Al<sub>2</sub>O<sub>3</sub> 33·54, CaO 17·63, Na<sub>2</sub> O 0·76, CaCO<sub>3</sub> 9·09=100·84. Granular, in lava.

1544°361. Microsommite. 4(NaK) CaAl<sub>3</sub>(SiO<sub>4</sub>)<sub>3</sub>.4(NaK) Cl.(NaK) SO<sub>4</sub>(?). Hexagonal, minute prism m, clear colorless, in leucitic lava. 4.00

- 120 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species No. No.
- 1545° I. Nasonite. (Ca,Pb)<sub>10</sub>Cl<sub>2</sub>Si<sub>6</sub>O<sub>21</sub>. Monoclinic(?), massive, greasy white. 2.00
- 1546° Davyne. SiO<sub>2</sub> 38.76, Al<sub>2</sub>O<sub>3</sub> 28.10, CaO 9.32, Na<sub>2</sub>O 15.72, K<sub>2</sub>O 1.10, CO<sub>2</sub> 5.63, H<sub>2</sub>O 1.96, Cl trace =100.59. Hexagonal, stout perfect prisms like nephelite, pearly colorless, in lava. 2.50

#### 2. Sodalite Group. Isometric. Hardness 5.5

The Sodalite and Helvite Groups may rightly be included in a broad grouping with the Garnet Group.

- 1547°362. Sodalite. Na<sub>4</sub>(AlCl) Al<sub>2</sub>Si<sub>3</sub>O<sub>12</sub>. Isometric, cube a, octahedron o and dodecahedron d, small, sharply symmetrical, subtransparent grayish-green. 1.50
- minute dodecahedrons d, glassy colorless, perfect, with neochrysolite on lava. 2.50
- 1549+ massive, cleavable-granular, azure-blue. .75 concentric nodule, chalcedony-like.
- 1550°363. Haüynite, Haüyne. Na<sub>2</sub>Ca(NaSO<sub>4</sub>.Al)Al<sub>2</sub>Si<sub>3</sub>O<sub>12</sub>. Isometric, dodecahedron d, small but sharply defined, glassy blue, in lava. 2.00
- 1551+ massive granular, green, in lava. .75
- grains, blue, embedded in lava. .75
- 1553 364. Noselite, Nosean. Na<sub>4</sub>(NaSO<sub>4</sub>.Al)Al<sub>2</sub>Si<sub>3</sub>O<sub>12</sub>. Isometric dodecahedron d. 3.00
- 1554° granular massive. 1.25
  - 365. Lazurite, Lapis-Lazuli. Essentially Na<sub>4</sub>(NaS<sub>3</sub>.Al)Al<sub>2</sub>Si<sub>3</sub>O<sub>12</sub>. Isometric, dodecahedron d, microscopic.
- fine granular in lava, blue. 1.00
- 1556° compact, ultramarine-blue, with pyrite, precious, polished. 3.00
- 1557+ compact, fine azure-blue, in white feldspar. .75
  - II. Hackmanite. Na<sub>4</sub>[Al(NaS)]Al<sub>2</sub>(SiO<sub>4</sub>)<sub>3</sub>. Isometric, reddish-violet.

## 3. Helvite Group. Range of Hardness 4.5-7

- 366. Helvite. 3(Be,Mn,Fe)<sub>2</sub>SiO<sub>4</sub>.(Mn,Fe)S. Isometric, tetrahedral, tetrahedron o<sub>1</sub> prominent with tetrahedron o.
- 1558\* tetrahedron o<sub>1</sub> (similar to fig.), small, ideal symmetry, resinous-yellow. 1.50


Type Species 1559

Helvite-Continued

ditto, minute, sulphur-yellow, on rhodochrosite. 1.00

Achtaragdite. Hydrated Al, Fe, Ca, Mg silicate. Pseudomorphous. Isometric, tetrahedral, hemitrisoctahedron n. minute.

 $(Fe,Zn,Mn)_2((Zn,Fe)_2S)$ 367. Danalite. Isometric, octa-Be<sub>3</sub>Si<sub>3</sub>O<sub>12</sub>. hedrons.



1558. Helvite

disseminated masses, reddish. 4.00 1560°

15610368. Eulytite. 2Bi<sub>2</sub>O<sub>3</sub>,3SiO<sub>2</sub>. Isometric, tetrahedral, minute hemi-trapezohedron n, sharply defined, adamantine. brownish. 2.00

twins, axes ||, minute. 3.00 1562

1563°369. Zunyite. (Al(OH,F,Cl)<sub>2</sub>)<sub>6</sub>Al<sub>2</sub>Si<sub>3</sub>O<sub>12</sub>. Isometric, tetrahedral, tetrahedron o prominent with tetrahedron  $o_1$ , sharply symmetrical, minute, whitish.

4. Garnet Group. 3RO.R<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>. Isometric, holohedral.

Range of Hardness 6.5-7.5

Ca, Mg, Fe<sub>2</sub>O<sub>3</sub>, Mn, Al, FeO or Cr orthosilicate. 370. Garnet. (See varieties). Isometric, highly symmetrical and sharply defined forms. Transparent varieties are precious:-

#### I. ALUMINIUM GARNET

A. GROSSULARITE. 3CaO.Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>. The following types are classed according to color:-

(a) white, dodecahedron d truncated by trapezohe-1564 dron n (fig. 1578).

(b) olive-green, dodecahedron d. .75 1565

olive-green, Wiluite, trapezohedron n, truncated by 1566\* dodecahedron d (fig. 1580), loose. .50

(c) amber-yellow, octahedron o truncated by dodeca-15670 hedron d. 3.00

1568+ (d) cinnamon-brown, Cinnamon Stone or Essonite, dodecahedron d, truncated by trapezohedron n(fig. 1578), brilliant.

122 CC Type Species No. No.	OMPLETE TYPE COLLECTION. DANA'S SYSTEM  Garnet—Continued
1569	(e) brownish-red, same form, with penninite. 2.00
15700	(f) pale rose-red, dodecahedron d.
	(g) emerald-green, containing Cr.
1571	B. Pyrope. 3MgO.Al <sub>2</sub> O <sub>3</sub> .3SiO <sub>2</sub> . Rounded grains in serpentine40
1572+	loose pebbles, clear deep red40
1573+	C. Almandite. 3FeO.Al <sub>2</sub> O <sub>3</sub> .3SiO <sub>2</sub> .  Dodecahedron d, coated with chlorite, very large, loose30
1574	ditto, squarish elongated60 ( / " )
1575	ditto, flattened60
15760	dodecahedron d, smooth ideal symmetry, red, large30
1577	ditto, small, in schist30 1579. Garnet
15780	dodecahedron d, truncated by tra- pezohedron n (fig.), large, ideal symmetry, bright, translucent deep red, in schist. 1.25
1579	trapezohedron n (fig.), deep red, large. 2.00
1580*	trapezohedron $n$ , truncated by dodecahedron $d$ (fig.), ideal symmetry, bright, deep red, loose30
15819	brownish-black, Common Garnet, trapezohedron n30
1582	D. Spessartite. 3MnO.Al <sub>2</sub> O <sub>3</sub> .3SiO <sub>2</sub> .  Transparent pale hyacinth-red semi-precious, loose75
15830	trapezohedron <i>n</i> , with dodeca- hedron <i>d</i> , bright, brownish- red, in rhyolite75
1584	dark hyacinth-red75
	II. IRON GARNET
	E. Andradite. 3CaO.Fe <sub>2</sub> O <sub>3</sub> .3SiO <sub>2</sub> . Sub-varieties:— 1. Calcium-iron Garnet.
1585*	(a) Topazolite, minute dodecahedron d, clear topaz- yellow. 1.50
1586	ditto, pale olive-green. 1.00
1587	Demantoid, massive, clear emerald-green. 3.00

## Monoclinic Species Related to the Foregoing

sive, black. 1.00

- 372. Partschinite. (Mn,Fe)<sub>3</sub>Al<sub>2</sub>Si<sub>3</sub>O<sub>12</sub>. Monoclinic, minute dull yellowish-red crystals in sand.
- 1603 373. Agricolite. Bi<sub>4</sub>Si<sub>3</sub>O<sub>12</sub>. Monoclinic, globular fibrous, yellowish-brown. 5.00
  - 5. Chrysolite Group. R<sub>2</sub>SiO<sub>4</sub>. Orthorhombic.

Range of Hardness 5.5—6.5

1604\*374. Monticellite. CaO.MgO.SiO<sub>2</sub>. Orthorhombic, brachyprism s, brachypinacoid b, unit pyramid e, brachydome k and macrodome d (fig.), translucent yellowish, in calcite. 2.00

124 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species Monticellite—Continued No. No.
1605 ditto, imperfect, rounded50
1606° rounded crystals in lava. 2.00
Batrachite, massive, with ceylonite. 1.50
1608 I. Glaucochroite. CaMnSiO <sub>4</sub> . Orthorhombic, embedded prisms, pale bluish-green (reddish by artificial light). 9.00
375. Forsterite. 2MgO.SiO <sub>2</sub> . Orthorhombic, white crystals in lava.
Boltonite, embedded grains, yellowish50 Monticellite
376. Chrysolite, Peridot. 2(MgFe)O.SiO <sub>2</sub> . Orthorhombic.
1610* 1. Precious, clear pale yellowish-green pebbles75
1611 2. Common, pale greenish-yellow, dull, rounded, in lava. 1.50
greenish-brown, prisms $m$ and $s$ (metalloidal), brachypinacoid $b$ , brachydome $k$ , clinodome $d$ (fig.), sharply symmetrical, loose75
1613+ Olivine, olive-green, granular30   m m a a
1614° ditto, grains in basalt30
1615 ditto, grains in kimberlite. 1.00
1616 ditto, grains in meteorite. 6.00 1612. Chrysolite
1617° 3. Hyalosiderite, 30 p.c. FeO, granular,
reddish-brown. 1.00
altered to serpentine, a,m,s,d,e,f, distinct, greenish-yel- low. 1.00
Villarsite. An altered chrysolite.
Matricite. SiO <sub>2</sub> 33·99, MgO 37·96, CaO 5·64, H <sub>2</sub> O 17·81. Concentric fibrous, pearly.
Ferrite. An alteration-product of chrysolite. SiO <sub>2</sub> 13·02, Al <sub>2</sub> O <sub>3</sub> 13·16, Fe <sub>2</sub> O <sub>3</sub> 53·47, FeO 4·51, MgO 6·63, H <sub>2</sub> O
8.39. Dark reddish-brown.
1619 376A. HORTONOLITE. (Fe, Mg) <sub>2</sub> SiO <sub>4</sub> . Orthorhombic, flattened    b, yellowish changing to dull black. 6.00
1620°376B. TITAN-OLIVINE. A chrysolite containing 6·10 p.c. TiO <sub>2</sub> .
Massive, dark reddish-brown. 3.00
I. Iddingsite. A chrysolite alteration-product(?). Fe,Ca,Mg silicate. Lamellar, brown.

Type Species No. No.

1621°377. Fayalite. 2FeO.SiO<sub>2</sub>. Orthorhombic, modified, tabular || a (fig.), microscopic but distinct, bright metalloidal iridescence, with glassy anorthoclase in obsidian lithophyses. 1.00

1622° cleavage mass, blackish-green. 2.00


1623° I. Neochrysolite. A variety of fayalite.

Slender prisms, dull black, with sodalite on lava. 2.50

1624°378. Knebelite. 2(Fe,Mn,Mg)O.SiO<sub>2</sub>. Orthorhombic, crystalline mass. 1.25

Igelströmite. About 10 p.c. more FeO, replacing MnO. 2.00

1626 379. Tephroite. 2MnO.SiO<sub>2</sub>. Orthorhombic, prismatic, grayish-brown. 2.00



1621. Fayalite

massive, grayish, with franklinite, etc. 1.25
 Hydrotephroite. (Mn,Mg)<sub>2</sub>.SiO<sub>4</sub>+<sup>2</sup>/<sub>3</sub>H<sub>2</sub>O. Reddish.
 Epigenite. (Mn,Mg).SiO<sub>4</sub>.H<sub>2</sub>O. Small bladed masses in tephroite, brownish-red.

1628 379A. ROEPPERITE. (Fe,Mn,Zn,Mg)<sub>2</sub>SiO<sub>4</sub>. Orthorhombic, large coarse crystal, yellow altering to black. 4.00

1629° massive crystalline. 1.50

## 6. Phenacite Group. R<sub>2</sub>SiO<sub>4</sub>. Rhombohedral.

Hardness 6-7

380. Trimerite. (Mn,Ca)<sub>2</sub>SiO<sub>4</sub>. Be<sub>2</sub>SiO<sub>4</sub>. Triclinic, thick pseudohexagonal tabular prisms, clear reddish.

------Hardness 5.5 and 7.5-8

1630°381. Willemite. 2ZnO.SiO<sub>2</sub>. Rhombohedral, unit prism m, pyramid p, minute, bright, translucent brownish, drusy. 1.50

1631 ditto, flesh-red. 1.50

slender hexagonal prisms, bright, clear grayish. 2.50

massive, very bright subtranslucent apple-green, fluoresces under Ra- and ultra-violet rays, with franklinite in calcite. 1.50

1634+ massive, subtranslucent brownish-apple-green, with franklinite, etc. .60

1635 ditto, reddish-brown. 1.00

1636° grains embedded in zincite, etc. .60

126 COMPLETE TYPE COLLECTION. DANA'S SYSTEM  Type Species Willemite—Continued No. No.
1637* Troostite, contains much Mn, diagonal
prism $a$ , rhombohedrons $r$ and $e$
(fig.), large, pale flesh-red, in cal-
cite. 1.50
1638 ditto, brownish, very large, rough, loose.
1.50
1639+382. Phenacite. 2BeO.SiO <sub>2</sub> . Rhombohedral,
unit and second order prisms m 1637. Willemite
and a with third order rhombo-
hedron x predominating (fig.),
bright, sharp, transparent,
loose (3). 1.00
1640 unit prism a, rhombohedron r
transparent, precious, faces m m
uneven, loose. 2.00
1641 large rough dull prism, semi-trans-
lucent 4.00
1642 <sup>®</sup> ditto, fragmentary. 1.00
Range of Hardness 4·5—5
Range of Hardness 4·5—5
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral,
Range of Hardness 4.5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a,
Range of Hardness 4.5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply sym-
Range of Hardness 4.5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald
Range of Hardness 4.5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645-384 Friedelite H (MnCl)Mn (SiO.) Rhombon
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombo-
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, minute sharp hexagonal
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombo-
Range of Hardness 4.5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, minute sharp hexagonal prism m, tabular    base c, translucent pale rose-red. 3.00
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, minute sharp hexagonal prism m, tabular    base c, translucent pale rose-red. 3.00  1646°385. Pyrosmalite. H <sub>7</sub> [(Fe,Mn)Cl](Fe,Mn) <sub>4</sub> (SiO <sub>4</sub> ) Rhombohedral thick six-
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, minute sharp hexagonal prism m, tabular    base c, translucent pale rose-red. 3.00  1646°385. Pyrosmalite. H <sub>7</sub> [(Fe,Mn)Cl](Fe,Mn) <sub>4</sub>
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, minute sharp hexagonal prism m, tabular    base c, translucent pale rose-red. 3.00  1646°385. Pyrosmalite. H <sub>7</sub> [(Fe,Mn)Cl](Fe,Mn) <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, thick six-m a m a
Range of Hardness 4.5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, minute sharp hexagonal prism m, tabular    base c, translucent pale rose-red. 3.00  1646°385. Pyrosmalite. H <sub>7</sub> [(Fe,Mn)Cl](Fe,Mn) <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, thick six-sided prism m and base c, pearly,
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, minute sharp hexagonal prism m, tabular    base c, translucent pale rose-red. 3.00  1646°385. Pyrosmalite. H <sub>7</sub> [(Fe,Mn)Cl](Fe,Mn) <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, thick six-sided prism m and base c, pearly, sharp ideal symmetry, liver-brown.
Range of Hardness 4·5—5  1643+383. Dioptase. H <sub>2</sub> O.CuO.SiO <sub>2</sub> . Rhombohedral, tetartohedral, diagonal prism a, rhombohedron s (fig.), sharply symmetrical, bright translucent emerald-green, small, loose. 1.00  1644 ditto, crystal aggregate. 5.00  1645°384. Friedelite. H <sub>7</sub> (MnCl) Mn <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, minute sharp hexagonal prism m, tabular    base c, translucent pale rose-red. 3.00  1646°385. Pyrosmalite. H <sub>7</sub> [(Fe,Mn)Cl](Fe,Mn) <sub>4</sub> (SiO <sub>4</sub> ) <sub>4</sub> . Rhombohedral, thick six-sided prism m and base c, pearly, sharp ideal symmetry, liver-brown.  2.50

7. Scapolite Group. Tetragonal. Hardness 6
Type Species
No. No.

1648\*386. Meionite. 4CaO.3Al<sub>2</sub>O<sub>3</sub>.6SiO<sub>2</sub>. Tetragonal, diametral prism a, truncated by unit prism m, unit pyramid r (fig.), transparent glassy colorless, on lava. 1.50

1649\*387. Wernerite, Scapolite. Al, Ca, and Na chloro-silicate. SiO<sub>2</sub>
46—54 p.c., Al<sub>2</sub>O<sub>3</sub> 31—24, CaO 17—10, Na<sub>2</sub>O 3—11,
Cl 1—3. Tetragonal, pyramidal hemihedrism,
unit and diametral prisms *m* and *a*, pyramid *r*(fig.), large and finely developed, brownish. 1.00

1650 $\circ$  ditto, with base c, gray. 1.00

ish. 1.00 highly modified, large, rough, whit-

1652+ massive, coarse cleavable-granular, pink. .20

massive, Glaucolite, bluish. 2.00

1654° massive, yellow. 1.00

1655 388. Mizzonite. Al, Na and Ca chlorosilicate. SiO<sub>2</sub> 54—60 p.c., Al<sub>2</sub>O<sub>3</sub>
24—20, CaO 10—6, Na<sub>2</sub>O 3—11, Cl 0—3. Tetragonal, minute prisms m and a, base c, clear whitish, on lava. 2.00

Dipyre, slender square prisms, embedded. .75
Prehnitoid, resembling prehnite.

1657 389. Marialite. Na<sub>4</sub>Al<sub>3</sub>Si<sub>9</sub>O<sub>24</sub>Cl. Tetragonal, minute squarish prisms, clear whitish, in lava. 5.00

Altered Scapolites:—Atheriastite, Stroganovite, Algerite, Terenite, Gabbronite, Pseudo-Scapolite, Paralogite.

16580 Wilsonite. Square cleavage prisms, pearly lavender. .50

1659°390. Sarcolite. 3RO.Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>, with R=Ca: Na<sub>2</sub>=9: 1. Tetragonal, pyramidal hemihedrism, aspect cubooctahedral, highly modified, glassy, in lava. 2.50

## 8. Melilite Group. Tetragonal. Hardness 5 and 5.5

1660\*391. Melilite. Na<sub>2</sub>(Ca,Mg)<sub>11</sub>(Al,Fe)<sub>4</sub>Si<sub>9</sub>O<sub>36</sub>(?). Tetragonal, short square prisms, minute but distinct, brown, with nephelite on lava. .75

Humboldtilite, octagonal prisms (unit and diametral m, a) glassy, sharply defined, in lava. 4.00

Type Species No. No.	COMPLETE TYPE COLLECTION. DANA'S SYSTEM  Melilite—Continued
16620	ditto, with calcareous coating. 2.00
1663 I.	Fuggerite. SiO <sub>2</sub> 34·04, Al <sub>2</sub> O <sub>3</sub> 17·97, Fe <sub>2</sub> O <sub>3</sub> 3·49.CaO 37·65, MgO 4·89, Na <sub>2</sub> O 2·04. Tetragonal(?), tabular, apple-green. 2.00
	Gehlenite. 3CaO.Al <sub>2</sub> O <sub>3</sub> .2SiO <sub>2</sub> . Tetragonal, short square prisms75
1665	Cacoclasite. Pseudomorphous. SiO <sub>2</sub> 32·67, Al <sub>2</sub> O <sub>3</sub> 19·63, CaO 36·38, P <sub>2</sub> O <sub>3</sub> 3·36, CO <sub>2</sub> 4·25. Tetragonal(?), large square prisms, cubo-octahedral aspect75
9.	Vesuvianite Group. Tetragonal. Hardness 6.5
1666 393.	Vesuvianite, Idocrase. $H(OH)_3Ca_{12}(Al,Fe)_6(SiO_4)_{10}$ (?). Tetragonal, unit and diametral prisms $m$ and $a$ and base $c$ , sharply symmetrical, vitreous greenish.
16670	ditto, with unit pyramid $p$ (fig.), ideal $m \mid a \mid m \mid a$ symmetry, loose, large. 2.00
1668	ditto, with ditetragonal prism f, brown, loose. 1.00
1669*	$m$ , $a$ , $c$ , pyramids $p$ and $\vartheta$ , distinct, 1667. Vesuvianite bright, dark olive-green. 1.00
1670	prisms $m$ deeply furrowed, terminated by six brilliant pyramids and base $c$ , translucent asparagus-green. 1.50
16710	pyramid p, sharp, yellowish. 1.50
1672	pyramid p, truncated by prisms m and a (fig.), ideal form but quite rough faces, very large, yellowish. 8.00
1673°	highly modified, adamantine, translucent brownish, in lava.  1.00 Possible Programme P
1674+	columnar radiated, brownish40

granular massive. .40

Mangan-vesuvianite, 12.49 p.c. MnO, reddish. 2.00 Cyprine, with trace of Cu, columnar, bluish-green. 1.00

1675 <sub>.</sub> 1676

1677°

# 10. Zircon Group. RSiO<sub>4</sub>. Tetragonal.

Type Spec	Hardness 7.5 and 5
	94. Zircon. ZrO <sub>2</sub> .SiO <sub>2</sub> . Tetragonal, unit pyramid p, ideal sym-
•	metry, reddish-brown, loose60
1679	ditto with base $c$ . 4.00
16800	unit pyramid $p$ truncated by unit prism $m$ (fig.), sharply symmetrical, bright,
	dark brown40
1681+	unit prism m, terminated by unit pyramid  p (fig.), ideal symmetry, bright, gray- ish-brown, loose (6)40
16820	ditto, with extra unit pyramid $u$ (fig.), ideal symmetry, adamantine, sub-translucent reddish-brown, loose40
16830	m, a, p and ditetragonal pyramid, or zirconoid, x (similar to fig. 1687), minute, perfect, transparent (3)40
1684	diametral prism a and unit pyramid p equally developed, resembling sharp rhombic dodecahedron, adamantine, 1681. Zircon sub-translucent reddish-brown, in feldspar. 1.00
1685	highly modified, adamantine, small. 1.50
16860	contact-twins, tw.pl. $e$ , geniculated (fig.),
	adamantine, brownish-red. 1.50
1687	Precious, Hyacinth, zirconoid (ditetragonal
	pyramid) $x$ , unit pyramid $p$ and diam- $m$ $m$
	etral prism a (fig.), adamantine,
	transparent red, small, loose75
1688+	ditto, much water-worn, lot40
1689	Precious, Jargon, clear yellowish, water-
	worn. 1.00
-6	Beccarite, olive-green.  II. Naëgite, contains UO <sub>3</sub> , ThO <sub>2</sub> , Cb <sub>2</sub> O <sub>5</sub> ,
1690	Ta <sub>2</sub> O <sub>5</sub> , Y <sub>2</sub> O <sub>3</sub> . 5.00 $m$
	Altered Zircon:—Tachyaphaltite,
	Erstedite, Auerbachite.
16910	Malacon, unit and diametral prisms
- <del>- y -</del>	m and a, unit pyramid p (fig.),
	bright, sharp, grayish, loose40 1686. Zircon

COMPLETE TYPE COLLECTION. DANA'S SYSTEM 130 Type Species No. No. Cyrtolite. SiO, 27.66, ZrO, 41.78, Er<sub>2</sub>O<sub>3</sub>, Y<sub>2</sub> 16920 O<sub>3</sub> 8.49, Ce<sub>2</sub>O<sub>3</sub> 3.98, CaO 5.06, MgO 1.10, H,O 12.07=100.14. Tetragonal, diametral pyramid e, strongly curved, brown. 1.00 Alvite. Essentially SiO<sub>2</sub> 20.33, ThO<sub>2</sub>(?) 15.33, 1693 1687. Zircon ZrO<sub>2</sub> 3.92, Y<sub>2</sub>O<sub>3</sub> 22.01, Al<sub>2</sub>O<sub>3</sub>, BeO 14.11,  $Fe_2O_3$  9.66,  $H_2O$  9.32. Tetragonal. 1.50 ThO, SiO, Highly radio-active. 16940395. Thorite. Tetragonal, unit prism m, terminated by unit pyramid p (fig.), brownish, loose. 2.50 unit pyramid p, truncated by unit prism m, 1695 blackish, complete, loose. 4.00 compact massive, resinous brownish-black. 1696+ 2.50 1691. Zircon Orangite, unit prism m, unit pyramid p, 1697 resinous orange-yellow. Orangite, compact massive, resinous-yellow. 1698\* Uranothorite, resinous, dark red-brown. 5ThSiO<sub>4</sub>.2Ca<sub>2</sub>SiO<sub>4</sub> + 10H<sub>2</sub>O. Calciothorite. Massive, translucent garnet-red. Eucrasite. Essentially Th, Y, Ce silicate. Freyalite. Essentially Th, Cesilicate. Resinous.

#### (RO) R<sub>3</sub>SiO<sub>4</sub>. Danburite—Topaz Group. Orthorhombic. 11. Range of Hardness 7-8

Auerlite. ThO<sub>2</sub>. (SiO<sub>2</sub>.  $\frac{1}{3}$ P<sub>2</sub>O<sub>5</sub>) + 2H<sub>2</sub>O(?). Tetragonal, square prism with pyramid,

1699°396. Danburite. CaO.B<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>. Orthorhombic, large squarish prism (fig.), opaque pale vellow. 3.00 large highly modified prism, brilliantly 1700

terminated, transparent, loose. 8.00 1701\* ditto, small, loose (12). 1.00 ditto, with adularia, etc. 3.00 1702

resinous.



1694.

Thorite

Danburite

Type Species	
397.	Topaz. [Al(O,F <sub>2</sub> )]AlSiO <sub>4</sub> . Orthorhom-
•	bic, crystals brilliant and sharply
	developed. Transparent varieties
	are precious:—
1703	unit and brachyprisms $m$ and $l$ , unit $\begin{vmatrix} l & m \\ l & m \end{vmatrix} = l$
	pyramids $u$ and $i$ , brachydome $  i   i''  $
	y and base c (fig.), transparent
	nale blue large loose 7.00
1704°	m and $l$ , uneven unit pyramid $u$
, ,	(fig.), adamantine, deeply striated,
	clear rich wine-yellow, large and
	slender, loose. 1.00 $u/u$
17050	unit and brachyprisms m and l, brachy-
	pinacoid b, acute unit pyramid o and
	brachydome y (fig.), transparent $\binom{l}{m}m$
	reddish, loose40
1706+	unit prism m and two brachyprisms land n,
•	pyramids $u, i$ and $x$ , brachydome $f$ and
	base c (fig.), transparent pale straw- 1704. Topaz
	yellow, loose (2)35
1707	ditto, on quartz75
1708+	m and $l$ , unit and obtuse pyramids $o$ and $u$ ,
•	brachydome y, macrodome d and base
	c, transparent colorless, loose (2)35
1709°	ditto, with extra brachydome f and pyra-
	mid <i>i</i> (similar to fig.), loose40 $l_m m l_b$
1710	ditto, pale wine-yellow, in rhyolite75
1711	unit and brachyprism $m$ and $l$ , brachy-
	domes $y$ , $f$ and $X$ , pyramid $u$ , macro-
	dome $d$ (fig.), clear colorless, loose. 1.50
17120	perfect basal cleavage, limpid. 1.00
1713	water-worn pebbles, limpid75
1714*	granular massive, grayish75
1715	compact massive, milky75
17160	Pycnite, columnar, pale yellowish75
1717+398.	Andalusite. $Al_2O_3.SiO_2$ . Orthorhombic, $n \mid n \mid n$
	unit prism m, base c truncated by
	brachydome s (fig.), coarse, large
	and symmetrical, grayish. 1.00
1718	massive, flesh-red75 1706. Topes
•	,,

132 CO Type Species No. No.	OMPLETE TYPE COLLECTION. DANA'S SYSTEM Andalusite—Continued
1719	imperfectly columnar, reddish-brown75
1720*	Chiastolite, cruciform arrangement of car-
2,00	bonaceous impurities in interior (fig.),
	polished cross-sections of prisms, loose
	- II V 1 III
•	(5). 1.00 m m lab
I.	Manganandalusite. Contains Mn.
	Hardness 6—7 and 5—7
1721+399.	Sillimanite, Fibrolite. Al <sub>2</sub> O <sub>3</sub> . SiO <sub>2</sub> . Orthorhom-
	bic, slender embedded prisms, gray40
	densely compact (prehistoric European 1709. Topaz
	"jade").
17229	Fibrolite, columnar-fibrous30
	Bamlite, subplumose, silky.
	Xenolite, rolled pebbles (Sp. gr. 3.58).
	Wörthite, hydrous (impure altered?),
	white.
•	Westanite. $SiO_2$ 42·53, $Al_2O_3$ 51·14. $l \mid m \mid m \mid l$
	(An altered andalusite?).
	Prismatic.
1722 400 (	Swanite Disthene ALO, SiO, Trica
1/23 100.	linic, blue prisms $M$ and $m$ ,
	macropinacoid a, brachypinacoid
	b, pyramid q and base c, trans-
	parent, loose. 2.00
1724°	blue, long flat prisms, transparent, in m
	paragonite. 1.00
1725	blue, cruciform-twins crossing at 60°.
	2.50
1726+	blue, curved bladed-columnar30
1727	green, transparent terminated crystal,
	loose. 1.00
_	
-	



1728\* green, bladed-columnar. .75
1729° white, Rhætizite, curved fibro-columnar. .75
II. Hibschite. CaO.Al<sub>2</sub>O<sub>3</sub>.2H<sub>2</sub>O.2SiO<sub>2</sub>. Isometric, minute crystals, colorless.

#### **12**. Datolite Group.

Monoclinic. Range of Hardness 5-7 Type Species 401. Datolite. H,O.2CaO.B,O,.2 SiO<sub>2</sub>. Monoclinic. 1. Glassy Crystals, rhombo-1730+ hedral aspect, highly modified (fig.), sharp and perfect. .60 ditto, large, colorless. 1731 ditto, greenish. 1.50 17320 thin sphenoidal by predom-1733° Datolite inance of orthodome x 1730. and clinodome t, truncated by several prisms and pyramids (fig.), triclinic aspect, highly perfect. 3.00 2. coarse subgranular massive. 1.00 1734° 3. compact massive, porcelain-like. 2.00 1735 4. botryoidal, Botryolite, radiated columnar. altered to Quartz, Haytorite. II. Bakerite. 8CaO.5B<sub>2</sub>O<sub>3</sub>.6SiO<sub>2</sub>.6H<sub>2</sub>O. Compact, greenish-

white.

1736**°402**. Homilite. 2CaO.FeO.B<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>. Monoclinic, tabular || base c (fig.), dark brown. 2.00 % octahedroid (M and m prominent.)

> Erdmannite. Chiefly silicate of Ce metals and Ca, with formula similar to datolite and gadolinite.



1737 403. Euclase. H<sub>2</sub>O.2BeO.Al<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>. Monoclinic, striated prism, well terminated, clear glassy. 9.00

17380 ditto, fragment. 3.00

2BeO.FeO.2Y,O3.2SiO2. Monoclinic, very 1739 404. Gadolinite. large rough prism with acute pyramid. 5.00

1740+ massive, vitreous black. 2.50

1741 405. Yttrialite. Chiefly silicate of Th and Y metals. Amorphous, vitreous greenish-black. 8.00

I. Thalénite. H<sub>2</sub>Y<sub>4</sub>Si<sub>4</sub>O<sub>15</sub>. Monoclinic, massive, flesh-red. 3.00 1742+

S. Rowlandite. 2Y<sub>2</sub>O<sub>3.3</sub>SiO<sub>2</sub>. Massive, dark drab-green. 5.00 1743

13.	Epidote Group. Range of Hardness 6-7
Type Species No. No.	dicite (CoO eA) O (SiO II O Oct och cochic cochic)
1744 406. 2	coisite. 4CaO.3Al <sub>2</sub> O <sub>3</sub> .6SiO <sub>2</sub> .H <sub>2</sub> O. Orthorhombic, vertical-
T # 4 # 0	ly furrowed prism, brownish, bright. 2.00
17450	ditto, greenish. 2.00
1746+	columnar-crystalline, ash-gray40
1747	fibrous aggregate, grayish-white60
1748*	rose-red, Thulite, massive, deep rose variegated with
	white quartz40
1749	ditto, disseminated in feldspar40
17500	ditto, minute indistinct crystals, pale rose40
1751+ <b>407.</b> E	Epidote. H <sub>2</sub> O.4CaO.3(Al,Fe) <sub>2</sub> O <sub>3</sub> .6SiO <sub>2</sub> . Monoclinic, slen-
	der prismatic    axis b, prominent planes are pyra-
	mid $n$ , orthopinacoid $a$ , orthodomes $r$ and $i$ and
	base c, truncated by lesser planes, sharply and bril-
	liantly defined, transparent greenish-black. Pleo-
	chroism strong: vibrations    c green, b brown and
	strongly absorbed, a yellow. Large, loose40
17520	ditto, twins, tw.pl. a (fig.)40
1753	ditto, group of crystals with byssolite.
17540	3.00 very large elongated crystal in quartz,
1754°	opaque pistachio-green. 1.00
17550	large well defined elemented empty 4
1755°	bright, ash-gray. 1.00
17560	short stout prismatic, small but sharply
1/30-	defined, pistachio-green75
1757	divergent columner crystale Disenite
1/3/	dark. 1.50
1758	acicular with feldspar40
1759*	crystalline columnar, pistachio-green40
1759*	granular massive, coarse60
1761+	granular massive, fine, green40
1701	Scorza (sand).
	Bucklandite, unit prism $m$ , pyramid $n$ and clinodome $o$ ,
	equally developed, affording a symmetrical bi-pyramid-like form, instead of the usual elongated habit.
600	·
17620	Withamite, small radial aggregates, embedded, deep
	red to pale yellow, strongly pleochroic. 1.50
1763	Beustite, grayish. 1.50

Type	Species No.
No.	No.

1769

S. Fouquéite, monoclinic but same composition as zoisite, elongated crystals.

Picroepidote. A magnesium-epidote(?). Prismatic || b, whitish, translucent.

1764 408. Piedmontite. H<sub>2</sub>O.4CaO.3R<sub>2</sub>O<sub>3</sub>.6SiO<sub>2</sub>, with R=Al: Mn: Fe=3:2:1 (?). Monoclinic, prismatic || axis b, vitreous dark reddish-brown. 2.00

1765\* bladed prismatic, embedded, dark purplish-brown. .75

1766 fibro-columnar in schist. 1.00

1767° massive, dark reddish-brown. .75

409. Allanite, Orthite. (Ca,Fe)<sub>2</sub>(AlOH) (Al,Ce,Fe)<sub>2</sub>(SiO<sub>4</sub>)<sub>3</sub>.

Monoclinic, varieties:—

1768°

1. Allanite, flat tabular || orthopinacoid a (fig.), large, rough.

slender acicular || axis b, very large. 1.50

1770 granular massive, brownish-black.
.50

k. r768. Allanite

1771+ compact massive, black. .50

2. Uralorthite, nearly anhydrous, large prisms.

- 3. Bagrationite. Contains Ce. Nearly symmetrical (not lengthened).
- 4. (Orthite originally included the very long *straight* prisms. Some authors now use it as the comprehensive name for the species).
- 5. Xanthorthite, hydrous, altered.
- 6. Pyrorthite, contains 30 p.c. carbonaceous impurity, altered, long prisms.

1772<sup>®</sup> I. Hancockite. Pb, Ca, Sr, Al, Fe<sup>III</sup> silicate. Monoclinic, microscopic flat prisms, bright, transparent brownish-red, with franklinite. 1.50

massive with polyadelphite. 1.00

Wasite. Much altered allanite, brownish-black.

Muromontite and Bodenite are chiefly silicates of Y earths and Ce metals with Fe, Al, etc.

14. Axinite Group. Triclinic. Hardness 6.5—7
Type Species
No. No.

1774° 410. II. Axinite. R"<sub>7</sub>R""<sub>4</sub>B<sub>2</sub>(SiO<sub>4</sub>)<sub>8</sub>, with R"=chiefly Ca, and R"=chiefly Al. Triclinic, unit prisms M and m, macropinacoid a, macrodome s and pyramids r and x (fig.), acute wedge-shaped, brown. 1.00

the area of the ar

1775\* unit prisms M and m, macropinacoid a, brachyprism w, distinct, clear brachypinacoid b, brachy-

dome  $\gamma$ , pyramids r and n and base c (similar to fig.), obtuse-edged table, with datolite. 1.50

1776 crystalline mass, brown. .50

yellow, highly modified, minute, clear glassy. 2.00

17780 yellow, crystalline mass. .50

Other Orthosilicates. Range of Hardness 5.5—6.5

1779 411. Prehnite. H<sub>2</sub>Ca<sub>2</sub>Al<sub>2</sub>Si<sub>3</sub>O<sub>12</sub>. Orthorhombic, tabular || base c, distinct. 2.50

1780 thin tabular aggregate of crystals united by c, pale green. 1.50

1781\* ditto, thick tabular. 1.00

17820 ditto, wheel-shaped. 1.00

1783 ditto, barrel-shaped, pearly. 1.50


1784+ drusy globular, translucent green.

.50

1785° drusy globular, radiated fibrolamellar, whitish. 1.50

II. Hillebrandite. 2CaO.SiO<sub>2</sub>.H<sub>2</sub>O. Orthorhombic, fibrous, white.

Uigite. SiO<sub>2</sub> 45.98, Al<sub>2</sub>O<sub>3</sub> 21.93, Na<sub>2</sub>O 4.69, CaO 16.15, H<sub>2</sub>O 11.25. Radiated sheafs of plates, pearly.



412. Harstigite.  $H_7(Ca, Mn)_{12}Al_3Si_{10}O_{40}(?)$ . Orthorhombic, prismatic || axis c, colorless.

<sup>1786 413.</sup> Cuspidine. Ca<sub>2</sub>SiO<sub>4</sub> with CaF<sub>2</sub>(?). Monoclinic, contacttwins, tw.pl. a, small spear-shaped, whitish. 6.00

#### IV. Subsilicates. Division 1. R<sub>5</sub>Si<sub>2</sub>O<sub>0</sub>


Humite Group. Hardness 6.5. See Ap. I.

These formulæ vary progressively by an increase of one molecule of (Mg<sub>2</sub>SiO<sub>4</sub>); likewise the vertical axes vary in the ratio 3:5:7:9, corresponding to the total number of magnesium atoms present. Hence the following transposition of the species 415 and 414.

Type Species

I. Prolectite.  $Mg[Mg(F,OH)]SiO_4(?)$ . Not yet analyzed. Monoclinic. Measured by Sjogren and named in allusion to Penfield and Howe's earlier prophecy of its discovery.

1787\*415. I. Chondrodite.  $Mg_3[Mg$ (F,OH)<sub>b</sub> $[SiO_a]_b$ . Monoclinic, highly modified, flattened || b, lenticular (fig.), translucent garnetred, bright, with clinochlore, etc. 1.50



1788 rounded, transparent yellow. 1.25 minute, brightly defined, pale 1789º yellowish, in lava. rounded, reddish-brown, in lime-1790 stone. 1.25 coarse crystalline grains, embedded, 1791+

vellow. .40

ditto, reddish-brown. .60 1792 1793


17940


massive, yellowish. altered to talc, highly modified (fig.), distinct. 2.00

1795\*414. I. Humite.  $Mg_s[Mg(F,OH)]_2[SiO_4]_3$ . Orthorhombic, very highly modified (fig.), small but brightly defined, translucent yellowish, in lava. 2.00

large distinct crystal in limestone. 8.00 1796 penetration-twins in lava. 17979

1798\*416. I. Clinohumite. Mg,[Mg(F,OH)],[SiO<sub>4</sub>]. Monoclinic, minute, very highly modified, yellowish, in lava. 4.00





COMPLETE TYPE COLLECTION. DANA'S SYSTEM 138 Type Species

1799 II. Leucophoenicite. 7MnO.3SiO<sub>2</sub>.H<sub>2</sub>O. Monoclinic(?), light purplish-red, with franklinite, etc. 1.00

1800 grayish, with hardystonite. 1.00

-Range of Hardness 6-6.5

1801 417. Ilvaite, Lievrite. H<sub>2</sub>O.CaO.4FeO.Fe<sub>2</sub>O<sub>3</sub>.4SiO<sub>2</sub>. Orthorhombic, bright distinct prisms, iron-black. 2.00

1802° large rough prism. 2.00

columnar massive. 1803\*

> $2R^{II}O.3R_2^{III}O_3.4SiO_2.3H_2O$ , with  $R^{II}=Ca$ II. Hellandite. chiefly, and R =Al, Fe, Mn and Ce metals. Monoclinic, prismatic, brown.

18040418. Ardennite. 5H<sub>2</sub>O.8MnO.4Al<sub>2</sub>O<sub>3</sub>.V<sub>2</sub>O<sub>5</sub>.8SiO<sub>2</sub>(?). rhombic, indistinct prisms, yellowish. .75

1805°419.S. I. Langbanite. 37Mn<sub>3</sub>SiO<sub>7</sub>10Fe<sub>3</sub>Sb<sub>2</sub>O<sub>8</sub>(?). Rhombohedral, minute distinct hexagonal prisms with base, black, on paisbergite. 2.00

#### Hardness 6.5 (Kentrolite 5) Kentrolite Group.

420. Kentrolite. 2PbO.Mn<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>(?). Orthorhombic, minute prism with pyramid.


compact massive, black. 2.00 1806°

> II. Molybdophyllite. RSiO<sub>4</sub> + H<sub>2</sub>O, with R =Pb, Mg. Hexagonal, foliated, colorless.

421.I. Melanotekite.  $(Fe_4O_3)Pb_3(SiO_4)_3$ . Orthorhombic, minute prism with pyramid.

crystalline druses, disseminated. 1.00 1807\* 1808°

massive, black. 1.00



Calamine 1810.

- 1809°422. Bertrandite. H<sub>2</sub>O.4BeO.2SiO<sub>2</sub>. Orthorhombic, hemimorphic, small tabular, glassy. 2.00
  - II. Stokesite. CaO.SnO<sub>2</sub>.3SiO<sub>2</sub>.2H<sub>2</sub>O(?). Orthorhombic, pyramidal, colorless.

#### Division 2. Hardness 5.5 (Tourmaline 7)

1810\*423. Calamine. H<sub>2</sub>O.2ZnO.SiO<sub>2</sub>. Orthorhombic, hemimorphic. group of small sharply defined crystals, short and thick, some showing the upper end (the analogous pole) highly modified with brachy- and macrodomes prominent, truncated by base c, others showing the lower (antilogous) pole with only brachypyramid v (fig.). 1.00

tabular || b (fig.), distinct, small. 1.50 18110

1812+ sheaf-like groups of tabular crystals, small, clear colorless. 1.00

wheel-shaped groups, milky, large. 18130 twins, tw.pl. c, axes || and antilogous poles of individuals together.

1814+ drusy crystalline. .50

18150 botryoidal, white. 1.50

botryoidal, blue. 2.00 1816

massive. .50 1817 carbonated, pisolitic.

1818 argillaceous, mixed with clay, soapy feel.

I. Clinohedrite. H<sub>2</sub>ZnCaSiO<sub>4</sub>. Monoclinic-clinohedral, minop181 ute, highly modified (fig.), adamantine, transparent pale amethystine. 9.00

1820 Moresnetite. SiO<sub>2</sub> 30·31, Al<sub>2</sub>O<sub>3</sub> 13·68, NiO 1·14, ZnO 43.41, H<sub>2</sub>O 11.37. Massive, green. 1.00

fibrous, straw-yellow. .75 18210

I. Lawsonite. H<sub>4</sub>CaAl<sub>2</sub>Si<sub>2</sub>O<sub>10</sub>. Orthorhombic, octahedroid, 1822 unit prism m, brachydome d, small. 1.50

tabular, grayish-blue, in margarite schist. 1823\* .75

twins, tw.pl. m. 1.50 1824°

1825°424. Carpholite. 2H<sub>2</sub>O.MnO.Al<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>. Monoclinic, radio-stellate tufts, silky strawyellow. 1.00

 $_{3H_{2}O.2}(Ca,Fe)O._{3}Ce_{2}O_{3}.6SiO_{2}(?).$ 425. Cerite. Orthorhombic, short prismatic.

massive, purplish-gray. 1.50 1826+

II. Beckelite. Ca<sub>3</sub>(Ce,La,Di)<sub>4</sub>Si<sub>3</sub>O<sub>15</sub>. Isometric, 1819. Clinohedrite small crystals, yellow.




1811. Calamine

	MPLETE TYPE COLLECTION. DANA'S SYSTEM
Type Species No. No. <b>426</b> . I	. Tourmaline. H <sub>9</sub> Al <sub>3</sub> (BOH) <sub>2</sub> Si <sub>4</sub> O <sub>19</sub> . Rhombohedral, hemimorphic. Usually a well-formed trigonal prism, rounded by striation, terminated by several rhombohedrons. Transparent varieties are precious.  I. VARIETIES BASED ON COLOR:—
1827°	(a) Rubellite, translucent dark purplish-red, short thick prism with rhombs, adamantine, loose. 2.00
1828	rich transparent pink, prism with rhombs, bright, loose. 1.50
1829*	pale clear pink, prism with base c, bright, loose50
1830	ditto, one end base c, other end several rhombohedrons, loose. 2.00
1831+	subtranslucent pink, large slender prisms radiating in pale lilac lepidolite50
18320	(b) Indicolite, indigo-blue, terminated prism. 2.00
1833	(c) Brazilian Sapphire, transparent prussian-blue. 3.00
1834	(d) Brazilian Emerald or Brazilian Peridot, clear green, bright, prism and acute rhombohedron y. 2.00
1835	opaque pale green, short thick prism with rhombs, in limestone. 1.50
1836+	subtransparent green, prism with rhomb $r$ , loose50
1837	(e) Ceylon Peridot, clear yellow. 3.00
18380	(f) Achroite, clear colorless terminated ma a ma prism50
1839*	(g) Black, unit and diagonal prisms m and a, rhombohedron r (fig.), very large, rough, loose. I.00
18400	obtuse rhombohedron r prominent at one end and acute rhombohedron o at the other end (fig.) 1.00
1841+	large slender bright prisms in white quartz40
18420	ditto, bent and broken, cross- sections later joined by quartz75
1843°	radiated acicular, in schist40
1844	capillary60
18450	columnar60

# SUBSILICATES, DIVISIONS 2 AND 3 Tourmaline-Continued

Type Species No. No. fibrous. .60 1846 massive compact. 1847\* .20 1848° (h) Brown, translucent unit and diagonal prisms m and a, rhombohedrons r and o, sharply defined, loose. Magnesium Tourmaline. .40 1849




1850+ ditto, indistinct, embedded. .40


18520

(i) Bi-colored, terminated prism, subtrans-1851 parent. 2.00

exterior green, interior red. 1.50 III.VARIETIES BASED ON COMPOSITION graduate



subtransparent polished cross-section,



141

1856. Staurolite

- (a) Alkali Tourmaline, Na and Li or both, also K.
- (b) Iron Tourmaline, usually black.
- (c) Magnesium Tourmaline.

Tourmaline altered to mica. 1853 .75

#### Range of Hardness 6.5-7.5 Division 3.

**427.** Dumortierite. 4Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>(?). Orthorhombic, fibrous, blue.


massive, dark violet-blue. 1.00 18540 2H<sub>2</sub>O.6(Fe, Mg)O.12Al<sub>2</sub> 1855 428. Staurolite. O<sub>3</sub>.11SiO<sub>2</sub>(?). Orthorhombic, unit prism m, brachypinacoid

b and base c (similar to fig.), ideal symmetry, large, loose, brownish. .40


1856° ditto, macrodome r prominent (fig.), large, loose. .30

1857º growth parallel to axis c, of cyanite within staurolite, both bright and sharply defined prisms, brownish-black.

cruciform-twins, tw.pl. x, individ-1858 uals crossing nearly at right angles (fig.), ideal symmetry, large. 1.50.



1858 Staurolite



1859. Staurolite

142 COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type Species Staurolite—Continued
No. No.

ditto, tw.pl. z (fig.), crossing at nearly 60°. .75

Nordmarkite, contains Mn.

- II. Grandidierite. 7SiO<sub>2</sub>.11(Al,Fe)<sub>2</sub>O<sub>3</sub>.7(Mg,Fe,Ca)O.2(Na, K,H)O. Orthorhombic, bluish-green.
- 429. Kornerupine. MgO.Al<sub>2</sub>O<sub>3</sub>.SiO<sub>2</sub>. Orthorhombic, with sapphirine, etc.
- 1860° Prismatine, slender embedded prisms, gray. 2.00 Kryptotil. A prismatine alteration-product.
- 1861°430. Sapphirine. 5MgO.6Al<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>. Monoclinic, granular, pale blue. 2.00
  - II. Sevendibite. Al, Ca, Mg basic silicate. Irregular grains, blue.
  - II. Silicomagnesiofluorite. Ca, Mg fluosilicate. Radio-fibrous.
- 1862° I. Roeblingite. 5H<sub>2</sub>CaSiO<sub>4</sub>. 2CaPbSO<sub>4</sub>. Compact mass of microscopic prisms, white. 4.00

### Appendix to Anhydrous Silicates

Barylite. 4BaO. Al<sub>2</sub>O<sub>3</sub>. 7SiO<sub>2</sub>. Tabular prisms. Monzonite. SiO<sub>2</sub> 52·60, Al<sub>2</sub>O<sub>3</sub> 17·10, FeO 9·00, CaO 9·65,

Na<sub>2</sub>O 6.60. Compact, light grayish-green.

Neociano. Anhydrous Cu silicate(?). Monoclinic, microscopic tables, blue, sublimate on lava. 2.00
 Sphenoclase. Chiefly Al and Ca silicate. Massive, yellowish.

### B. Hydrous Silicates

True hydrous compounds, containing water of crystallization (e.g. the Zeolites), also hydrous amorphous clays, as well as certain acid or basic silicates (Micas, Talc, etc.), which yield water on ignition and which bear a close relationship to the true hydrous species. Finally are included certain species in which the chemical constitution and the part played by water, remain in doubt.

#### I. Zeolite Division

# 1. Indroductory Subdivision

Hardness 6 and 4-4.5

1864 431. Inesite. 2(Mn,Ca)SiO<sub>3</sub>+H<sub>2</sub>O. Triclinic, small prisms. 2.50

1865° divergent fibrous, rose-red. 1.25

Type Species

1878

432. Ganophyllite. 6H<sub>2</sub>O.7MnO.Al<sub>2</sub>O<sub>3</sub>.8SiO<sub>2</sub>. Monoclinic, short prisms terminated by acute clinodome e and base c.

foliated micaceous, brown. 5.00

—Hardness 4.5—5, 3—4 and 4.5—5

1867 433. Okenite. 2H<sub>2</sub>O.CaO.2SiO<sub>2</sub>. Orthorhombic(?), mass of minute white prisms. 2.00

1868 434. II. Gyrolite (=Zeophyllite?). 3H<sub>2</sub>O. 2CaO.3SiO<sub>2</sub>. Rhombohedral, tetartohedral, white. 3.00

1869 II. Agnolite (formerly Manganocalcite). 3MnO.4SiO<sub>2</sub>.2H<sub>2</sub>O. Triclinic, radiating fibrous, pale red. .75

1870°435. Apophyllite. K<sub>2</sub>O.8CaO.16SiO<sub>2</sub>.16H<sub>2</sub>O. Tetragonal, cuboid, striated diametral prism a, pearly base c (similar to fig.), sharp ideal symmetry, white. 1.25

octahedroid (fig.), brilliant limpid, small, with copper. 1.00

1872 ditto, opaque milky, with pink drusy natrolite. .75

1873+ ditto, elongated, brilliant glassy, subtransparent. .75

unit pyramid p, diametral prism a (fig.), sharply symmetrical, brilliant, rosepink. 3.00

1875\* ditto, clear colorless, small. .75


thin tabular || base c, with prism a and pyramid p (fig.), clear colorless. 3.00 187


1877° lamellar massive, pink. 1.00
Albine, altering to calcite.

Xylochlore, contains Fe, olivegreen.


Tesselite, cuboid, tessellated structure.

Leucocyclite. Basal sections show, in polarized light, a black cross with alternate white and violet-black rings. 2.00









144 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species No. No.

II. Astrolite. (Na,K)<sub>2</sub>.Fe(Al,Fe)<sub>2</sub>.(SiO<sub>2</sub>)<sub>5</sub>.H<sub>2</sub>O(?). Globular, radio-stellate structure, siskin-green.

OTHER HYDROUS CALCIUM SILICATES, IMPERFECTLY DE-FINED.

Xonotlite.  $4\text{CaSiO}_3 + \text{H}_2\text{O}(?)$ . Massive.

Tobermorite. Chiefly hyd. Ca silicate. Granular.

Chalcomorphite. Chiefly hyd. Ca silicate. Hexagonal, minute acicular prisms.

Plombierite. CaSiO<sub>3</sub>+2H<sub>2</sub>O. Massive.

#### 2. Zeolites

A family of well defined hydrous silicates, closely related in composition and all occurring as secondary minerals in cavities and veins of basic igneous rocks. They are silicates of aluminium with chiefly sodium and calcium, rarely barium and strontium. The Zeolites are analogous to the Feldspar Group, except that the former include independent groups of diverse form and distinct composition. Intumescence under the blowpipe is marked.

### Mordenite Group. Hardness 3—4

1879°436. Ptilolite. (Ca, K<sub>2</sub>, Na<sub>2</sub>) Al<sub>2</sub>Si<sub>10</sub>O<sub>24</sub> + 5H<sub>2</sub>O. White spongy mass of minute crystalline colorless needles. 1.50

437. Mordenite.  $3RAl_2Si_{10}O_{24} + 20H_2O$ , with  $R=K_2$ :  $Na_2$ : Ca=1: 1: 1. Monoclinic, tabular || clinopinacoid b, minute, pearly.

I. Erionite. CaO.K<sub>2</sub>O.Na<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.6SiO<sub>2</sub>6H<sub>2</sub>O. Orthorhombic, aggregates of slender fibers, pearly white.

Steeleite. Partly altered mordenite. Chalky balls.


Pseudonatrolite. Hydrous Ca, Al silicate. Minute needles.

Heulandite Group. Monoclinic. Range of Hardness 3.5—4.5 1880 438. Heulandite.  $5H_2O.CaO.Al_2O_3.6SiO_2$ . Monoclinic, unit prism m, orthodomes s and t and clinopinacoid b, tabular ||b| (pearly), small, brilliantly symmetrical,

unit prism m, clinopinacoid b (pearly), orthodomes s and t and base c (fig.), yellowish-white. .75

1882 ditto, curved, brilliant snow-white, large. 1.50

1883° saddle-shaped group of nearly parallel individuals, large. .75



Type Species No. No.


Heulandite-Continued

1884\*

tabular || clinopinacoid b (pearly), copper-red. 1.00

18850439. Brewsterite. (Sr, Ba, Ca) O. Al<sub>2</sub>O<sub>3</sub>.6SiO<sub>2</sub>. 5H<sub>2</sub>O. Monoclinic, minute stout prisms, brightly defined, translucent pale yellowish-gray.

1886°440. Epistilbite. CaO.Al<sub>2</sub>O<sub>3</sub>.6SiO<sub>2</sub>.5H<sub>2</sub>O. Monoclinic, twins, tw.pl. orthopinacoid a, prismatic (fig.).



#### Phillipsite Group. Monoclinic. Range of Hardness 4-4.5

1887 441. Phillipsite.  $(K_2,Ca)Al_2Si_4O_{12} + 4\frac{1}{2}H_2O.$ Monoclinic, simple penetrationtwins, tw.pl. base c. 1.50

1888

cruciform penetration-twins (preceding twinned, tw.pl. e, fig.), small, perfect, opaque white. 1.00

1889+

ditto, clear glassy, minute. 1.00

1890°

1892

complex penetration-twins (twinning of three of the preceding double twins, tw.pl. m, fig.), with phacolite.

1888. Phillipsite


18919 drusy, globular white. 1.00

Spangite. A variety of phillipsite.

S. Offrétite.  $(K_2Ca)_2Al_6Si_1O_{30}$ . 17H<sub>2</sub>O. Hexagonal, microscopic hex-

agonal prisms, white. 1.00

1893\*442. Harmotome. (K<sub>2</sub>,Ba)O.Al<sub>2</sub>O<sub>3</sub>.5SiO<sub>2</sub>. 5H<sub>2</sub>O. Monoclinic, cruciformpenetration-twins, tw.pl. c, united as fourlings with tw.pl. e (fig. 1888), small, brilliant ideal symmetry, white.



1894

ditto, aspect of square prism (without re-entrant angle), terminated by diagonal pyramid.

I. Wellsite. (BaCaK<sub>2</sub>)O.Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>.3H<sub>2</sub>O. Monoclinic, complex-twins, transparent whitish.

1895°443. Stilbite, Desmine. (Na<sub>2</sub>,Ca)O.Al<sub>2</sub>O<sub>3</sub>.6SiO<sub>2</sub>.6H<sub>2</sub>O. Monoclinic, penetration-twins, tw.pl. base c (fig.), thin tabular || clinopinacoid b (pearly), sharply defined, white. 1.00

146 Type Species	COMPLETE TYPE COLLECTION. DANA'S SYSTEM Stillbite—Continued		
18960	sheaf-like groups of preceding in parallel growth (fig.), creamyellow, large60		
1897+	ditto, rounded, brown40		
1898	lamellar-columnar40		
1899*	stellate, radio-fibrous60 m		
1900	globular50		
1901	foliated, yellowish40 1895. Stilbite		
19020	foliated, brick-red75		
1903	Foresite. Chiefly hyd. Al, Ca silicate. Monoclinic, like		
	stilbite, minute. 2.00		
	Hardness 4·5		
10049444	Gismondite. CaAl <sub>2</sub> Si <sub>4</sub> O <sub>12</sub> +4H <sub>2</sub> O. Monoclinic, complex		
.,04	twins, pseudo-tetragonal octahedroids, faces rough		
	composite, small. 2.00		
1905 II	I. Bavenite. 3CaO.Al <sub>2</sub> O <sub>3</sub> .6SiO <sub>2</sub> .H <sub>2</sub> O. Monoclinic, minute		
prisms in spherical groups, whitish. 4.00			
1906*445	i. Laumontite. 4H <sub>2</sub> O.CaO.Al <sub>2</sub> O <sub>3</sub> .4SiO <sub>2</sub> . Mono-		
	clinic, square prism m, obliquely termin-		
	ated by orthodome e, embedded, copperred40		
TAA=0	ditto, sharply developed, white. 1.00		
19070	crystalline amygdules in diabase, salmon-		
1908			
	red40		
1909	Leonhardite, altered. 1.00		
446. Laubanite. 2CaO.Al <sub>2</sub> O <sub>3</sub> .5SiO <sub>2</sub> +6H <sub>2</sub> O. Fibrous			
	radiating, white, exterior yellowish. 1896. Stilbite		


Chabazite Group. Rhombohedral. Range of Hardness 4.5-5

1910 447. Chabazite. (Ca,Na<sub>2</sub>)Al<sub>2</sub>Si<sub>4</sub>O<sub>12</sub>+6H<sub>2</sub>O.
Rhombohedral, cuboid rhombohedron r, ideal symmetry, glassy white. .50

1911\* ditto, flesh-red, Acadialite. .75

1912+ penetration-twins, tw. axis c (fig.), ideal symmetry, lustrous

white. .50 1913 ditto, brown. .75



1912. Chabasite

Type Species No. No.	Chabazite—Continued	-4,
	Haydenite, twinned    R, yellow-	
1914		
	ish, small. 1.00	Milland
1915	Phacolite, Herschelite or See-	8 8
	bachite, penetration-twins,	3 3
	ideal pseudo-hexagonal tables,	1/1/1/1
	c prominent, small. 1.50	1916. Chabazite
19160	ditto, lenticular (fig.). 1.50	1910. Chaoaste
19170	ditto, highly composite cruciform-	
	twins, spherical aspect. 2.00	
1918	ditto, drusy-globular. 1.00	
19190448.	Gmelinite. $(Na_2,Ca)Al_2Si_4O_{12}+6$	(m)
	H <sub>2</sub> O. Rhombohedral, cuboid,	m
	rhombohedrons $r$ and $\rho$ with	
	prism m (fig.), small, brightly	
	defined. 1.50	1919. Gmelinite
1920*	penetration-twin,tw.axisc, pseudo-	
- )	hexagonal (fig.), ideal sym-	
	metry, small, flesh-red. 1.50	
1921	ditto, rounded lenticular, white.	
1921	1.50	m
	•	7 10
	Groddeckite. Hyd. Fe, Al, Mg, Na	
	silicate. Rhombohedral, clear	
	colorless.	1920. Gmelinite
449.	Levynite. $CaAl_2Si_3O_{10}+5H_2O$ .	
	Rhombohedral, twins, whit-	$\int \int n $
	ish.	
1922+450.	Analcite. Na <sub>2</sub> O.Al <sub>2</sub> O <sub>3</sub> .4SiO <sub>2</sub> .2H <sub>2</sub> O.	$\binom{n}{n}$
-	Isometric, trapezohedron n,	
	ideal symmetry (fig.), milky.	
	.75	
1923	ditto, reddish-white, large. 1.50	1922. Analcite
1924	composite group of preceding	1922. Analote
- 7-4	(similar to fig.). 2.00	180
T025#	cube a, truncated by trapezo-	AHA X
1925*		HATTER
	hedron n, (fig.), brilliantly	
	symmetrical, limpid, small,	RETURN THE
	on lava. 1.00	
	Euthallite, compact, greenish.	The state of the s
	Eudnophite, cleavages, unusually	All S
	strong double refraction.	1924. Analcite

ZEOLITES. CHABAZITE GROUP


147

- 148 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species No. No.
- 1926°451. Faujasite. Na<sub>2</sub>O.CaO.2Al<sub>2</sub>O<sub>3</sub>.10SiO.20H<sub>2</sub>O(?). Isometric, small octahedrons, sharply defined, grayish. 1.00
- 1927°452. I. Edingtonite. BaO.Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>.3H<sub>2</sub>O(?). Orthorhombic, hemihedral, prismatic cleavage piece, white. 4.00

### Natrolite Group. Hardness 5

453. Natrolite. Na<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>+2 H<sub>2</sub>O. Orthorhombic. 1. Ordinary varieties:— 1928\* (a) square stout unit prism m

(a) square stout unit prism m bright, obtuse square unit pyramid o dull, ideal symmetry, gray. 1.50



1929. Natrolite

1929 ditto, slender, clear colorless, brilliant (fig.), diverging group.

brilliant (fig.), diverging group. 3.00 ditto, very slender, forming surface of large ball with fibro-columnar radiating structure, yellowish-white 1.50

1931 acicular, clear colorless. 1.25

19300

19350


druse of minute prisms, flesh-red, with apophyllite. .75

1933+ (b) radio-fibrous mass, white. .75 1934 (c) solid amygdules, radiated. .75

(d) compact massive, chalk-white. 1.00

2. Fargite, 4.31 p.c. CaO, red.

Iron-natrolite, 10 p.c. iron oxides as impurity, dark green.



1936 454. Scolecite. CaO.Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>.3H<sub>2</sub>O. Monoclinic, large prisms, brilliantly terminated, interlacing aggregate, white. 2.00

1937\* columnar, divergent. 1.25 radio-fibrous.

1938 455. Mesolite. Hyd. Ca, Na, Al silicate. Intermediate between natrolite and scolecite. Monoclinic and triclinic, acicular. 2.50

	LITES. NATROLITE AND THOMSONITE GROUPS 149		
Type Species No. No.	Mesolite—Continued		
1939*	downy tufts of diverging hairs. 1.50		
19409	· ·		
	fibrous stalactites, radiated structure.		
1941	amorphous, chalk-white. 1.50		
1942 I.	Gonnardite. (Ca,Na <sub>2</sub> ) <sub>2</sub> Al <sub>2</sub> Si <sub>5</sub> O <sub>15</sub> +5½ H <sub>2</sub> O. Orthorhombic (?) radio-fibrous amygdules, silky-white. 1.00		
	Thomsonite Group. Hardness 5-5.5		
456.	Thomsonite. (Na <sub>2</sub> ,Ca)O.Al <sub>2</sub> O <sub>3</sub> .2SiO <sub>2</sub> .½H <sub>2</sub> O. Orthorhombic.		
	1. Ordinary varieties:—		
19430	(a) Rectangular prisms, base c prominent, glassy pearly. 2.50		
1944*	(b) Slender prisms, small, indistinct, forming surface of a radiated encrustation, with analcite. 1.00		
	(c) Radio-fibrous.		
1945+	(d) Spherical concretions (amygdules), compactly radio- fibrous with concentric zones of white and shades of red, precious50		
1946	ditto, rolled pebbles (lot)50		
1947	Lintonite, spherical amygdules, compact, translucent sage-green75		
19480	spherules, compact, translucent pearly75		
1949	filmy coating on calcite rhombs, translucent pearly75		
1950	Ozarkite, radiated, white75		
1951	2. Mesole, Faroelite, radio-lamellar spherules. 1.00		
	3. Chalilite, compact reddish-brown.		
	Picrothomsonite. Hyd. Al, Mg, Ca silicate. Radio-lamellar masses, pearly.		
1952 457.	Hydronephelite. 2Na <sub>2</sub> O.3Al <sub>2</sub> O <sub>3</sub> .6SiO <sub>2</sub> .7H <sub>2</sub> O. Hexagonal		
	(?), radiated massive, altered from sodalite. 1.25		
	Ranite, altered from elæolite.		
II.	Lotrite. $4SiO_2.2(Al,Fe)_2O_3.3(Ca,Mg)O.2H_2O$ . Massive, green.		
TT	Legallite Man Al O sSin al/H (1/2) Fibrous snow.		

II. Melite. 2(Al,Fe)<sub>2</sub>O<sub>3</sub>.SiO<sub>2</sub>.8H<sub>2</sub>O. Prismatic, massive, bluish-brown.

white.

Type Species No. No.	Appendix to Zeolites
No. No. 1953	Chlorastrolite. Impure hyd. Al, Ca, etc. silicate. Amygdules, stellated-mosaic structure, pearly sage-green.
19540	ditto, rolled pebbles, precious (lot)50
1955	Zonochlorite. Impure hyd. Al, Ca, Fe, etc. silicate. Banded amygdules, sage-green. 3.00
	Sasbachite. Hyd. Al, Ca, K silicate. Massive.
	Sloanite. Chiefly hyd. Al, Ca silicate. Radiated masses, pearly.

### II. Mica Division

Monoclinic species with highly perfect basal cleavage, easily yielding thin laminæ. Their often closely related forms have a rhombic or hexagonal aspect.

1. Mica Group. Monoclinic. Range of Hardness 2.5—3

Laminæ more or less elastic.

458. Muscovite. Potash Mica. Generally 2H<sub>2</sub>O.K<sub>2</sub>O.3Al<sub>2</sub>O<sub>2</sub>.

150. 1-24500 vito, 1 otabii viica. Otherany 21120.1120.5111203.
6SiO <sub>2</sub> . Monoclinic.
1. Ordinary Muscovite, tapering
rhombic aspect, prism M,
base c (rough), large75 $M$
ditto, with clinopinacoid b, 1957. Muscovite
hexagonal outline, tabular
(similar to fig.), green, in lava75
ditto, base (bright cleavage), gray, very large20
rhombic outline, prism $M$ , bright cleavage $   c$ , very
large, gray30
2. Damourite, small silky-gray scales, coating corun-
dum40
Damourite, curved scales, pearly gray40
Margarodite, scaly granular, pearly yellowish-white,
with tourmaline, dravite40
Margarodite, very coarse scaly-granular, pearly-
gray, with topaz60
Gilbertite, small spherical groups of hexagons, pale
olive-green, with fluor, etc75

	MICA GROUP	
Type Species No. No.	Muscovite—Continued	
1965	ditto, yellowish-white, pearly75	
1966	Ivigtite, disseminated in cryolite50	
19679	Sericite, fine scaly-fibrous schist, silky30	
	Pycnophyllite, spherical masses, greasy feel, green.	
19680	3. Oncosine, compact, green. 1.00	
1969*	Fuchsite, 1 to 4 p.c. Cr <sub>2</sub> O <sub>3</sub> . Very fine scaly-granular,	,
	greenish50	
1970	Avalite. 14.59 p.c. Cr <sub>2</sub> O <sub>3</sub> . Earthy mass of microscopic	:
	scales, with cinnabar, etc. 1.50	
	Oellacherite. 4.65 to 5.82 p.c. BaO.	
I.	Baddeckite. 25.82 p.c. Fe <sub>2</sub> O <sub>3</sub> . Small scales, pearly copper-	-
	red.	
	E is a general term for numerous alteration-products.	
	ntially aluminium and potassium hydrous silicate, often	
	rresponding to muscovite, and is probably a compact and	l
usually ve	ry impure variety of this species.	
19719	Pinite. Altered iolite. Octagonal prisms with base, dis-	
19/10	tinct, loose30	•
	Gigantolite. Altered iolite. Very large 12-sided prisms.	
1972	Gieseckite. Altered nephelite. Large hexagonal prisms	,
	grayish75	
	Lythrodes. Regarded as altered nephelite.	
1973	Liebenerite. Altered nephelite. Small hexagonal prisms	,
	embedded, greenish75	
1974	Dysyntribite. Altered nephelite. Massive, waxy, mot-	-
	tled greenish and reddish40	
	Rosite. Altered anorthite. Granular, red.	
	Polyargite. Altered anorthite. Lamellar, reddish.	
1975	Pinitoid. Altered feldspar. Massive, green50	
19760	Agalmatolite, Pagodite. A general term for a soft, com-	
	pact, easily carved, mottled pinite. (Includes also	
	compact pyrophyllite and steatite). Carved piece	٠
	1.00	
1977	Oösite. Altered iolite. Reddish prisms40	
1978	Cataspilite. Altered iolite, with a little more CaO than	
	the foregoing. Rounded gray prisms in schist50	)
1979@459.	Paragonite, Sodium Mica. 2H <sub>2</sub> O.Na <sub>2</sub> O.3Al <sub>2</sub> O <sub>3</sub> .6SiO <sub>2</sub>	•
717	Massive, microscopic scales, laminated, pearly	7
	grayish-white, with cyanite50	
	6	

152 COMPLETE TYPE COLLECTION. DANA'S SYSTEM  Type Species Paragonite—Continued No. No.
Cossaite, compact, greenish.
1980 Euphyllite. Na-K-mica between muscovite and para-
gonite. Pearly white, with corundum. 1.25
1981 º 460. Lepidolite, Lithia Mica. KLi[Al(OH,F)2]Al(SiO3)3. Aggre-
gates of short prisms, slightly rounded termination,
pearly pale reddish-lilac. 1.00
1982 ditto, tabular, whitish. 1.00
1983° cleavable plates, gray40
1984+ coarse scaly-granular, deep lilac20
1985 fine scaly-granular, pale lilac20
II. Irvingite.
I. Cookeite. Monoclinic. Hyd. lithia mica. Slender six- sided prisms.
1986• fine scaly-granular, whitish50
1987°461. Zinnwaldite, Lithium-iron Mica. (K,Li) <sub>3</sub> FeAl <sub>3</sub> Si <sub>5</sub> O <sub>16</sub> (OH,
F) <sub>2</sub> . Monoclinic, rosette-like groups of six-sided
tables, gray. 1.50
1988* very coarse cleavable-granular, pearly dark-gray40
Rabenglimmer, 19.78 p.c. Fe <sub>2</sub> O <sub>3</sub> . Dark gray.
1989 Cryophyllite, only 16 p.c. Al <sub>2</sub> O <sub>3</sub> . Strongly pleochroic:
<b>c</b> violet, <b>b</b> greenish-gray75
Polylithionite, only 12 p.c. Al <sub>2</sub> O <sub>3</sub> .
Protolithionite. A dark lithium-iron mica.
462. Biotite, Magnesium-iron Mica. (H,K) <sub>2</sub> (Mg,Fe) <sub>2</sub> (AlFe) <sub>2</sub> (SiO <sub>4</sub> ) <sub>3</sub> . Monoclinic. Pleochroism strong.
NOTE-Tschermak classes biotite thus: I. Meroxene. Ax.pl.    b, including
nearly all varieties. II. Anomite. Ax.pl. $\perp b$ , rare.
1990* six-sided tables, translucent green, small, in lava60
1991 ditto, scale-like, clear brown75
1992+ broad cleavage, basal, black20
1993 <sup>o</sup> Barytbiotite, 6.84 p.c. BaO. 1.25
Chromglimmer, 5.90 p.c. Cr <sub>2</sub> O <sub>3</sub> .
Siderophyllite. $3H_2O.6(K,Na,Li)_2O.21FeO.10Al_2O_3$ .
30SiO <sub>2</sub> . Black75
Haughtonite. Mg largely replaced by Fe. Blackish.
1995° Manganophyllite. 5.41 to 21.40 p.c. MnO. Tabular,
bronze-red. 1.25
1996 mass of fine scales75
1997 <sup>o</sup> Rubellan. Altered biotite. Hexagonal forms in basalt,
copper-red50

Type Species No. No.

Hydrated biotites:-Eukamptite, Voigtite, Rastolyte, Hvdrobiotite.

Pseudobiotite. Altered biotite. Chiefly Al, Fe, Mg silicate. Bastonite. Altered iron mica. Pearly.

1998°462A. Phlogopite, Magnesia Mica. R, Mg, Al

(SiO<sub>4</sub>), with R=H,K,MgF. Monoclinic, very large coarse six-sided tabular prism with basal cleavage. Very thin sheets show strong asterism (six-rayed-star), when held close to the eye, in viewing a candleflame. Phenomenon due to minute acicular inclusions. Pearly bronzebrown. .75

1999

large coarse tapering six-sided prism (fig.), copper-red. .75

2000+

cleavage showing parting on edge, asteriated (see 1998) pearly bronzebrown. .20

1999. Phlogopite

cleavage, green. .40 200I

2002°462B. LEPIDOMELANE. (H,K)<sub>2</sub>Fe<sub>3</sub>(FeAl)<sub>4</sub>(SiO<sub>4</sub>)<sub>5</sub>. Monoclinic, small six-sided tables, adamantine black.

Pterolite. An altered lepidomelane. Scaly massive, pearly.

I. Alurgite. HR<sub>2</sub>(AlOH)Al(SiO<sub>3</sub>)<sub>4</sub>. Monoclinic, scaly massive, purple.

-Soft

2003°463. Roscoelite.  $H_8K(Mg,Fe)(Al,V)_4(SiO_3)_{12}(?)$ . Small scales in fan-shaped groups, pearly dark-brown. 3.00

II. Moravite. H<sub>4</sub>Fe<sub>2</sub>(Al,Fe)<sub>4</sub>Si<sub>7</sub>O<sub>24</sub>. Foliated, black.

Clintonite Group. Monoclinic. Hardness 4.5 2. (Ottrelite 6-7). Basic. Laminæ brittle

2004 464. Margarite. H<sub>2</sub>CaAl<sub>4</sub>Si<sub>2</sub>O<sub>12</sub>. Monoclinic, very thin tabular c, gray. 2.00

laminated, pearly grayish-pink. .75 2005+

schistose, pearly grayish green, with lawsonite. .50 20060

2007\*465. Seybertite. 3H<sub>2</sub>O.10(Mg,Ca)O.5Al<sub>2</sub>O<sub>3</sub>.4SiO<sub>2</sub>. Monoclinic. 1. Clintonite, foliated crystalline, pearly submetallic

reddish-brown. .75

154 COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type Species Seybertite—Continued
No. No.


2008

2. Brandisite, hexagonal prisms. 1.25

465A. XANTHOPHYLLITE. H<sub>8</sub>(Mg,Ca)<sub>14</sub>Al<sub>16</sub>Si<sub>5</sub>O<sub>52</sub>(?). Monoclinic, crystalline crust.

2009 Waluewite, tabular || c, green. 1.25 466. Chloritoid. H<sub>2</sub>(Fe, Mg) Al<sub>2</sub>SiO<sub>7</sub>. Monoclinic or triclinic.

 Original chloritoid, large curving laminæ, mottled green.



2010

2. Sismondine, with glaucophane.

3. Salmite. 8.40 p.c. Mn. Saccharoidal masses, gray.

2011+

20129

4. Masonite, mass of plates, blackish-green. .35

467. OTTRELITE. H<sub>2</sub>(Fe, Mn) Al<sub>2</sub>Si<sub>2</sub>O<sub>9</sub>(?). Monoclinic or triclinic, hexagonal crystalline scales.

Venasquite, H<sub>2</sub>FeAl<sub>2</sub>Si<sub>3</sub>O<sub>11</sub>. Crystalline radio-lamellar. Phyllite, small black crystalline scales, in schist. .30

I. Cosmochlore. 'Cr silicate. Monoclinic(?), emerald-green.

#### 3. Chlorite Group


Monoclinic. Hardness 2.5 (Prochlorite 1-2)

Ferrous iron gives to most of these species a green color. They are closely related to the micas in their monoclinic form, basal cleavage and optical characters, but their laminæ are comparatively inelastic. The Chlorites are essentially silicates of aluminium with ferrous iron and magnesium, and chemically combined water, manganese rarely replacing the ferrous iron. Tschermak calls those members of the group which occur in distinct crystals or plates, Orthochlorites; and the fine scaly or indistinctly fibrous forms, Leptochlorites.

468. Clinochlore, Ripidolite. 4H<sub>2</sub>O.5MgO.Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>. Monoclinic.

· •

I. Ordinary varieties:—
(a) hexagonal crystal, tabular
|| base c, (fig.), dark green,
with chondrodite. 1.00



2014

20139

large rhombic crystal. 2.50

twins, penninite law, tw.pl. base c (fig.).

2016+


cleavage plate, green. .50

CHLORITE	GROUP
Clinochlore—C	Continued

Type Species	CHLORITE GROUP 155 Clinochlore—Continued		
No. No.			
2017	(b) foliated, green75		
	(c) massive, green.		
20180	2. Leuchtenbergite, large rough tabular hexagon, pale		
	grayish-green. 2.00		
	3. Kotschubeite, 4 to 11.39 p.c. Cr <sub>2</sub> O <sub>3</sub> , rhombic habit, red.		
4. Manganiferous, 2.3 p.c. MnO, steep rhombs built			
	up of lamellæ in twinning position.		
468A.	PENNINITE. H <sub>8</sub> (Mg,Fe) <sub>5</sub> Al <sub>2</sub> Si <sub>3</sub> O <sub>18</sub> . Monoclinic, pseudo-rhombohedral.		
2019*	1. Penninite, hexagonal prisms, pearly base, dark green.		
2019	1.00		
2020	tapering trigonal to hexagonal prisms, loose (6)75		
202 I	small crested groups of hexagonal tables. 1.00		
2022 <sup>©</sup>	twins, penninite law, tw. pl. c (fig.), on chromite, small.		
	.50		
2023	2. Kämmererite, small hexagonal		
	forms bounded by steep six-		
	sided pyramids, red, on chro-		
•	mite. 2.00		

Kämmererite, fibro-lamellar, pale 20240 violet. .75

> 3. Loganite, altered amphibole, brown.



2022. Penninite

Pseudophite, compact talc-like, green.

2025 469. Prochlorite, Chlorite. H<sub>40</sub>(Fe,Mg)<sub>23</sub>Al<sub>14</sub>Si<sub>13</sub>O<sub>90</sub>. Monoclinic, six-sided prism, vermicular, green. 2.00

spheroidal groups of six-sided tables. 20269

foliated. .75 2027

2028+ fine scaly-granular, green. .30

Pycnochlorite, contains Mg, massive, grayish-green. II. Chiefly hyd. Mg, Al silicate. Grochauite. Monoclinic (?), small rough tabular hexagons.


2029 470. Corundophilite. H<sub>20</sub>Mg<sub>11</sub>Al<sub>8</sub>Si<sub>6</sub>O<sub>45</sub>. Monoclinic, six-sided tables, dark green, with emery and diaspore. 2.00

foliated, with emery, etc. .75 20300

> Approximately H<sub>4</sub>(MgFe)<sub>2</sub>Al<sub>2</sub>SiO<sub>9</sub>. Amesite. talc-like, pearly apple-green, with diaspore.

-Range of Hardness 1-3.5

- 471. Daphnite. H<sub>56</sub>Fe<sub>27</sub>Al<sub>20</sub>Si<sub>18</sub>O<sub>121</sub>. Monoclinic, spherical aggregates, concentric radio-foliated structure, pearly dark green.
- 2031 Metachlorite. Hyd. Al, Fe silicate. Foliated-columnar, dull leek-green. 1.25
- 2032 Klementite. SiO<sub>2</sub> 27·13, Al<sub>2</sub>O<sub>3</sub> 24·70, Fe<sub>2</sub>O<sub>3</sub> 5·84, FeO 9·72, MnO 1·98, MgO 20·52, H<sub>2</sub>O 11·35=100·24. Thin scales, olive-green. 1.00
  - II. Brunsvigite. 6SiO<sub>2</sub>.2Al<sub>2</sub>O<sub>3</sub>.9MgO.8H<sub>2</sub>O. Crypto-crystalline, radio-foliated masses, olive-green.
- 2033°472. Cronstedtite.  $4\text{FeO.2Fe}_2\text{O}_3.3\text{SiO}_2.4\text{H}_2\text{O}(?)$ . Rhombohedral, hemimorphic, tapering trigonal unit pyramid x and base c (fig.), small, ideal symmetry, brilliant black. 2.00



- 2035°473. Thuringite. 8FeO.4(Al,Fe)<sub>2</sub>O<sub>3</sub>.6SiO<sub>2</sub>.9H<sub>2</sub>O. Massive, Green. .50
- 2036 Chamosite, Berthierine. Hyd. Fe, Al silicate. Massive. .50
  - II. Stilpnochloran. Alteration-product of Thuringite.
  - 474. Stilpnomelane. 2(Fe,Mg)O.(Fe,Al)<sub>2</sub>O<sub>3</sub>.5SiO<sub>2</sub>.3H<sub>2</sub>O(?). Crystalline plates.
- 2037° Chalcodite, velvety coating of microscopic scales, brass-like luster, brown, with ankerite. .75
  - 475. Strigovite. 2FeO.(Fe,Al)<sub>2</sub>O<sub>3.2</sub>SiO<sub>2.2</sub>H<sub>2</sub>O (at 100°), or with 3H<sub>2</sub>O (air-dried). Microscopic hexagonal prisms, dark green altering to brown.
- 2038 476. Diabantite. 12(Fe,Mg)O.2Al<sub>2</sub>O<sub>3</sub>.9SiO<sub>2</sub>.9H<sub>2</sub>O. Monoclinic (?), massive, greenish-black. .50
- 2039 477. Aphrosiderite. H<sub>10</sub>Fe<sub>6</sub>(Fe,Al)<sub>4</sub>Si<sub>4</sub>O<sub>25</sub>(?). Mass of microscopic hexagonal scales, clear olive-green. .50
- 2040°478. Delessite. H<sub>10</sub>(Mg,Fe)<sub>4</sub>(Al,Fe)<sub>4</sub>Si<sub>4</sub>O<sub>23</sub>(?). Massive, scaly fibrous, green. .75
  - 479. Rumpfite. 7MgO.8Al<sub>2</sub>O<sub>3</sub>.1oSiO<sub>2</sub>.14H<sub>2</sub>O. Massive granular, vermicular groups of microscopic hexagonal scales, greenish-white.

II. Spodiophyllite. (Na<sub>2</sub>K<sub>2</sub>)<sub>2</sub>(Mg,Fe)<sub>3</sub>(Fe,Al)<sub>2</sub>(SiO<sub>3</sub>)<sub>8</sub>. Hexagonal micaceous prisms, gray.

### Other Chloritic Minerals, Imperfectly Defined

Epichlorite. Hyd. Al, Fe, Mg silicate. Fibro-columnar, dull leek-green.

Euralite. Hyd. Fe, Al, Mg silicate, near diabantite. Massive, greenish-black.

2041 Chlorophæite. Near delessite and hisingerite. Granular amygdules, blackish-green. .40

Epiphanite. SiO<sub>2</sub> 37·11, Al<sub>2</sub>O<sub>3</sub> 21·13, FeO 20·00, MgO 14·03, H<sub>2</sub>O 7·83=100·10.

Melanolite. Chiefly hyd. Fe silicate. Crusts.

Ekmannite. Chiefly Fe, Mn silicate. Foliated.

Berlauite. Chiefly hyd. Al, Fe, Mg silicate. Scaly mass, green.

Steatargillite. Hyd. Al, Fe, Mg silicate. Earthy amygdules, whitish.

Pattersonite. Hyd. Al, Fe, Mg, K silicate. Scaly.

### Appendix to Micas-Vermiculites. Soft

Indefinite alteration-products of the micas, etc. Remarkable vermiform exfoliation on ignition. Laminæ generally pearly.

2042\*480. Jefferisite. Approximately  $R_3(AlO_2)MgSiO_4.3H_2O + H_2$   $Mg_2Al_2(SiO_4)_3.3H_2O$ . Broad crystalline cleavage plates, yellowish-brown. .40

II. Tænislite. (K,Li)<sub>2</sub>O.MgO.3SiO<sub>2</sub>.2H<sub>2</sub>O(?). Monoclinic, micaceous blades, colorless with blue tinge.

Vermiculite. SiO<sub>2</sub> 35.74, Al<sub>2</sub>O<sub>3</sub> 16.42, FeO 10.02, MgO 27.44, H<sub>2</sub>O 10.30=99.44. Scaly-massive. .75 Kerrite. A trihydrated phlogopite. Fine scales.

Lucasite, Philadelphite, Maconite and Dudleyite are Hyd. Mg, Fe, Al, K silicates.

Lennilite. Hyd. Al, Fe, Mg silicate. Foliated, green. .75

2045 Hallite. Hyd. Mg, Fe, Al silicate. Large rough hexagonal micaceous prisms. .75

2046 Protovermiculite. Hyd. Fe, Mg, Al silicate. Broad micaceous plates, silvery yellowish. .40

158	COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type Species No. No.	) 
	Vaalite. Hyd. Mg, Fe, Al silicate. Hexagonal prisms.
2047	Pyrosclerite. Hyd. Mg, Al silicate. Disseminated scales
2049	apple-green. 1.25
2048	Roseite. SiO <sub>2</sub> 35·38, Al <sub>2</sub> O <sub>3</sub> 30·30, MgO 14·66, H <sub>2</sub> O 19·88= 100·32. Spherical groups of small distinct hexagons
	pearly drab50
	Willcoxite. Chiefly hyd. Al, Mg and alkali silicate. Pearly
	whitish talcose scales.
	III. Serpentine and Talc Division
	Range of Hardness 2.5—3.5 (Talc 1)
	range of Hardrass 2-5 - 5-5 (Faic 1)
481	. Serpentine. 3MgO.2SiO <sub>2</sub> .2H <sub>2</sub> O. Monoclinic.
	A. In CRYSTALS, Pseudomorphs. See altered chrysolite
	pyroxene, chondrodite, etc.
2049	In crystals, i.e. perfect cubic parting (pseudomorphous?), in part crystalline, pearly. 1.50
	B. Massive.
	1. Ordinary massive:—
2050	(a) Noble, translucent pale oil-green, veined, polished75
2051*	Noble, translucent rich oil-green30
2052	(b) Common, compact, dark green30
2053+	common, granular, light green20
2054	2. Resinous, Retinalite, waxy translucent yellowish40
	3. Porcellanous, compact smooth.
20550	4. Bowenite, very fine granular, translucent pale apple-
	green40 C. Lamellar.
	5. Antigorite, thin lamellar, brownish-green.
2056*	6. Williamsite, sublamellar, impure, translucent leek-
20304	green30
2057	ditto, more compact, precious, polished. 1.00
37	D. Thin Foliated.
2058°	7. Marmolite, pearly whitish50
•	8. Thermophyllite, pearly brownish.
	E. Fibrous.
2059+	9. Chrysotile or Serpentine Amianthus, the principal
	Asbestus of commerce. See also amphibole. Olive-
	green solid mass of extremely fine and long white
	silken threads, easily separable50

2. Massive, Steatite or Soapstone:—

2075+ 20769 (a) Coarse granular-schistose, grayish. .20

(b) Fine granular, French Chalk, white.

160	COMPLETE	<b>TYPE</b>	COLLECTION.	DANA'S SYSTEM
Type No.	Species No.		Talc—Continued	

2077

- (c) Indurated, impure slaty, dark green, dull. .50
- 3. Pseudomorphous:—
- (a) Fibrous, altered from enstatite.

20789

- (b) Rensselaerite, wax-like. .30
- (c) Pyrallolite, partly altered pyroxene.
- 2079+485. Sepiolite, Meerschaum. 2H<sub>2</sub>O.2MgO.3SiO<sub>2</sub>. Very compact earthy, smooth feel, white. .40
  - 486. Connarite. 2H<sub>2</sub>O.2NiO<sub>2</sub>.3SiO<sub>2</sub>(?). Hexagonal(?), small crystals, greenish.
  - 487. Spadaite. 5MgO.6SiO<sub>2</sub>.4H<sub>2</sub>O(?). Massive, greasy luster, translucent flesh-red.
    - I. Batavite. 4H<sub>2</sub>O.4MgO.Al<sub>2</sub>O<sub>3</sub>.4SiO<sub>2</sub>. Micaceous, pearly hexagonal scales.

-Soft

- 2080\*488. Saponite. Hyd. Mg, Al silicate, impure(?). Massive. .40 2081°489. Celadonite. Fe, Mg, K silicate. Earthy, celandine-green. .50
- 2082 490. Glauconite. Chiefly hyd. Fe, K silicate, variable mixture.

  Amorphous, earthy chloritic, green, in rock. .40
  2083\* sand, "marl", grayish-green. .20
- 491. Pholidolite. Approximately 5H<sub>2</sub>O.K<sub>2</sub>O.12(Fe, Mg)O.Al<sub>2</sub>
  O<sub>3</sub>.13SiO<sub>2</sub>. Minute crystalline scales, grayish-yellow.
- IV. Kaolin Division. Hardness 1-2 (Allophane, Schrötterite 3)
  - 492. Kaolinite. 2H2O.Al2O3.2SiO2. Monoclinic, 1. Crystals.
    - 2. Ordinary:-
- 2084+
- (a) argilliform, soft clayey, white. .20
- 2085
- (b) fariniform, loose mealy. .20
- **2086**0
- (c) indurated, Lithomarge, firm compact. .50
- 3. Ferruginous, red lithomarge.
- 2087 Rectorite. 2HAlSiO<sub>4</sub>+ H<sub>2</sub>O. Monoclinic(?), leathery plates, soapy feel, pearly whitish. 1.00
  - Leverrierite. 2Al<sub>2</sub>O<sub>3</sub>.5SiO<sub>2</sub>.5H<sub>2</sub>O(?). Orthorhombic(?), hexagonal prisms, pearly brownish.
  - 493. Halloysite.  $2H_2O.Al_2O_3.2SiO_2 + H_2O.$  Massive, clayey.
- 2088\*
- I. Ordinary, opaque waxy white. .40
- 2. Smectite, transparent when moist.
- 3. Lenzinite, compact, opaline white.

Type Species No. No. 2089

manoyate—Continued

4. Bole. Impure(?). Some Fe and 24 p.c. H<sub>2</sub>O. .40

II. Termierite. Hyd. Al silicate. Clay-like.

494. Newtonite. Al<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>.5H<sub>2</sub>O. Rhombohedral, soft compact mass of microscopic cuboid rhombs, white.

2090°495. Cimolite. 2Al<sub>2</sub>O<sub>3</sub>.9SiO<sub>2</sub>.6H<sub>2</sub>O. Amorphous clayey, adheres to the tongue, whitish. .50

2091°496. Montmorillonite.  $H_2Al_2Si_4O_{12} + nH_2O(?)$ . Massive clayey, rose-red. .50

Razoumovskyn. Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>+6H<sub>2</sub>O(?). Clayey, green.

2092+497. Pyrophyllite. H<sub>2</sub>O.Al<sub>2</sub>O<sub>3</sub>.4SiO<sub>2</sub>. Monoclinic(?), radiated fibro-lamellar, greasy feel, pearly whitish. .75

2093 ditto, brownish. .75

2094° compact massive, steatitic, grayish. .50

Neurolite. Hyd. Al silicate. Fibrous, yellow.

Biharite. Hyd. Al, Mg, Ca, K silicate. Massive.

498. Allophane. Al<sub>2</sub>SiO<sub>5</sub>+5H<sub>2</sub>O. Amorphous, mammillary incrustation, translucent yellowish.

2095+ ditto, sky-blue, cupriferous. .75

2096 ditto, compact mass. .75

Plumballophane, contains some Pb, stalactitic.

2097 Carolathine. Hyd. Al silicate. Mammillary, yellow. 1.00 Samoite. 2Al<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>.10H<sub>2</sub>O(?). Stalactitic, whitish.

499. Collyrite. 2Al<sub>2</sub>O<sub>3</sub>.SiO<sub>2</sub>.9H<sub>2</sub>O. Amorphous, greasy feel, adheres to the tongue, white.

2098 500. Schrötterite. 8Al<sub>2</sub>O<sub>3</sub>.<sub>3</sub>SiO<sub>2</sub>.<sub>3</sub>oH<sub>2</sub>O. Amorphous. 1.25 I. Alexandrolite. Contains H<sub>2</sub>O<sub>3</sub>,Cr<sub>2</sub>O<sub>3</sub>,SiO<sub>2</sub>. Amorphous, green.

# Appendix to Clays

See the "System of Mineralogy" for brief description of numerous other hydrous aluminous silicates, mostly impure clays and all of doubtful character.

# V. Concluding Division. Range of Hardness 3-5.5

2099 501. I. Cenosite. Ca(Y,Er)<sub>2</sub>(SiO<sub>3</sub>)<sub>4</sub>.CaCO<sub>3</sub>.2H<sub>2</sub>O. Orthorhombic, small short prisms, greasy brownish. 4.00

2100\*502. I. Thaumasite. [(CaOH)CO<sub>2</sub>][(CaOH)SO<sub>3</sub>][(CaOH)HSiO<sub>4</sub>] + 13H<sub>2</sub>O. Hexagonal, loose mass of minute prisms, white. .50 162 COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type Species
No. No.

II. Spurrite. 2Ca<sub>2</sub>SiO<sub>4</sub>.CaCO<sub>3</sub>. Monoclinic(?), granular, gray. 2101°503. Uranophane. CaO.2UO<sub>3</sub>.2SiO<sub>2</sub>+6H<sub>2</sub>O. Orthorhombic, massive, lemon-yellow. 2.00

-Range of Hardness 2-4

2102+504. Chrysocolla. CuSiO<sub>3</sub>+2H<sub>2</sub>O. Cryptocrystalline, deep turquois-blue. .50

ditto, finely banded, agate-like. 1.00 ditto, banded with malachite. 2.00

ditto, brecciated, polished. 4.00 botryoidal crust, bluish-green.

botryoidal crust, bluish-green. 1.00
ditto, coated with drusy quartz, affording glistening
translucent turquois-blue surface. 4.00

II. Plancheite. 15CuO.12SiO2.5H2O. Fibrous, blue.

2108\*505. Chloropal. Fe<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>.5H<sub>2</sub>O(?). Amorphous, opal-like, greenish-yellow. .50

Anthosiderite. 2Fe<sub>2</sub>O<sub>3</sub>.9SiO<sub>2</sub>.2H<sub>2</sub>O. Fibrous flowery tufts, yellowish.

I. Hoeferite. 2Fe<sub>2</sub>O<sub>3</sub>.4SiO<sub>2</sub>.7H<sub>2</sub>O. Amorphous, green.

II. Müllerite. Fe<sub>2</sub>O<sub>3</sub>.3SiO<sub>2</sub>.2H<sub>2</sub>O. Massive, yellowish-green.

506. Hisingerite. A hydrated ferric silicate of doubtful homogeneity. Amorphous, compact, brownish.

Scotiolite, contains much Mg, black.

Jollyte. Hydrated ferric silicate. Compact. 2.00
Jollyte. Hyd. Al, Fe, Mg silicate. Compact.
Melanosiderite. 4Fe<sub>2</sub>O<sub>3</sub>.SiO<sub>2</sub>.6H<sub>2</sub>O. Amorphous, compact, vitreous black.

II. Morencite. Silicate of Fe<sup>III</sup> with H<sub>2</sub>O(?). Fibrous, brownish-yellow.

- 2110°507. Bementite. Approximately 2MnSiO<sub>3</sub>.H<sub>2</sub>O. Foliated-stellate mass, pale grayish-yellow. 2.00
- 2111 508. Caryopilite. Approximately 4MnO.3SiO<sub>2</sub>.3H<sub>2</sub>O. Massive, minutely reniform crust, brown. 1.00
- 2112 509. Neotocite. Hyd. Mn, Fe silicate. Amorphous, black. 2.50 Penwithite. MnSiO<sub>3</sub>+2H<sub>2</sub>O. Massive, clear glassy brownish.
  - II. Bityite. Hyd. Ca, Al silicate, also contains Be, Li, Mg, Na and K. Pseudo-hexagonal, minute plates.
  - II. Aloisiite. Hyd. silicate containing FeO, CaO, MgO, Na<sub>2</sub>O. Amorphous cement in tuff, brown to violet.

### Appendix to Hydrous Silicates

Under this heading in the "System of Mineralogy," will be found brief description of a large number of amorphous, massive and often heterogeneous compounds, mostly of doubtful chemical constitution. They are mainly silicates of magnesium, very frequently with aluminium, iron, calcium, etc.

# Titano-Silicates, Titanates. Hardness 5.5-6.5

Hardness 6 and 5.5 fkinite. Chiefly Th and Ce metals titanon A heterogeneous alteration-product. Mas reous velvet-black. 3.00	
 hyllite. $R_4R_4Ti(SiO_4)_4$ , with $R=H$ , Na, K, Fe, Mn chiefly, including also the $Fe_2O_3$ . rhombic, very long thin blades, elongated    by development of brachypinacoid, pearly br	Ortho- cleavage

- 2126 ditto, slender squarish prisms, stellated. .75
  - II. Lorenzenite. Na<sub>2</sub>O.2TiO<sub>2</sub>.2SiO<sub>2</sub>. Orthorhombic, minute needles, nearly colorless.
  - I. Lamprophyllite. Contains SiO<sub>2</sub>, Ti, Fe, Mn, Na. Minute flattened prisms, yellow-brown.
- 2127 II. Benitoite. BaO.TiO<sub>2</sub>.3SiO<sub>2</sub>. Rhombohedral, transparent blue. 7.00
  - II. Narsarsukite. Fe<sup>III</sup> and Na acidic titano-silicate. Tetragonal, tabular, honey-yellow.
- Range of Hardness 4—5
  2128 515. Johnstrupite. A complex Ce, Ca and Na titano-fluo-silicate. Monoclinic, brownish-green. 1.50
- 2129°516. Mosandrite. Ce, Ca and Na titano-fluo-silicate. Monoclinic, very rough large flat prism, not terminated, brown. 1.00
  - 517. Rinkite. (F<sub>8</sub>Ti<sub>4</sub>) Na<sub>9</sub>Ca<sub>11</sub>Ce<sub>3</sub>(SiO<sub>4</sub>)<sub>12</sub>(?). Monoclinic, flattened || a, yellowish-brown.
    - Hardness 5.5
- 2130\*518. Perovskite. CaTiO<sub>3</sub>. Isometric or pseudo-isometric, cube, brownish, loose. .50
- 2131 highly modified, adamantine blackish, small. 1.00
- 2132° I. Knopite. RO.TiO<sub>2</sub>, with R=Ce,Zr,Y,Si,Fe,Ca,Mn,Mg, K,Na. Isometric, small cubo-octahedrons, blackish lead-gray. 1.50
  - I. Zirkelite. (Ca,Fe)O.2(Zr,Ti,Th)O<sub>2</sub>. Isometric, octahedrons, black.
- 2133 I. II. Geikielite. MgO.TiO<sub>2</sub>. Rhombohedral, rolled pebbles, black. 4.00
- 2134\*519. Dysanalyte. Approximately 6(Ca,Fe)TiO<sub>3</sub>.(Ca,Fe)Nb<sub>2</sub>O<sub>6</sub>. Isometric, perfect cubes, splendent iron-black, loose (6) .50
- 2135 ditto, cubo-octahedrons (fig.), (6). .25
- 2136 ditto, with monticellite, small. 1.00

II. Yttrocrasite. Y earths and Th hyd. titanate. Orthorhombic, pitch-black.

Hydrotitanite. Altered dysanalyte, perfect 2137 cubo-octahedrons, dull yellowish-gray, loose (6). .25



II. Delorenzite. 2FeO.UO<sub>2.2</sub>Y<sub>2</sub>O<sub>3.24</sub>TiO<sub>2</sub>(?). Orthorhombic, prismatic, black.

#### Columbates, Tantalates 3.

(Columbates is the latest international usage; Niobates is employed in the "System of Mineralogy.") Chiefly salts of metacolumbic and metatantalic acid, RCb<sub>2</sub>O<sub>6</sub> and RTa<sub>2</sub>O<sub>6</sub>.

1. Pyrochlore Group. Isometric. Range of Hardness 5-5.5

II. Chalcolamfrite. R''O.(Cb<sub>2</sub>O<sub>5</sub>).R"F<sub>2</sub>. R"O.SiO<sub>2</sub>(?). Nb<sub>2</sub>O<sub>5</sub> 2138 59.65 p.c., SiO<sub>2</sub> 10.86, ZrO<sub>2</sub> 5.71, CaO 9.08, Na, O 3.99, F 5.06. Isometric, small octahedrons, dark grayish-brown inclining to red. 1.25

2139\*520. Pyrochlore. Chiefly Ca, Na and Ce metals columbate with Ti, Th and F. Isometric, octahedron o, perfect, brown. 1.25

ditto, with dodecahedron d, trape-

zohedron m (fig.). 2.00


II. Marignacite.

2140


2141 520A. KOPPITE. Essentially Ce and Ca pyrocolumbate. Isometric, minute dodecahedrons. clear brown. 1.00

> 521. Hatchettolite. U and Ca tantalocolumbate. Isometric. resinous yellowish-brown.


2142°522. Microlite. Essentially Ca, Ta, O<sub>7</sub>. Isometric, octahedron o,



Pyrrhite. (Microlite?). Isometric, microscopic octahedrons, orange-yellow.



2140. Pyrochlore



#### 2. Fergusonite Group

Tetragonal. Hardness 5.5-6

Type Species

2144

2145

2143+523. Fergusonite. (Y,Er,Ce) (Cb,Ta)O<sub>4</sub>. Highly radio-active. Tetragonal, hemihedral pyramid s prominent, unit pyramid s, base c (similar to fig.), distinct, dull grayish-brown externally, brilliantly vitreous brownish-black fracture, loose. 1.50




ditto, large, imperfect, in feldspar. 1.50 ditto, fragments with autunite (lot). 1.50 2143. Fergusonite

2146 524. Sipylite. Chiefly Er CbO<sub>4</sub>. Tetragonal, massive, brownish-black. 3.00

Adelpholite. Fe, Mn columbate. Tetragonal.

### 3. Columbite Group. Orthorhombic. Hardness 6

2147 525. Columbite. (FeMn) Cb<sub>2</sub>O<sub>6</sub> with (Fe, Mn) Ta<sub>2</sub>O<sub>6</sub>. Orthorhombic, macropinacoid a, brachypinacoid b, macrodome k, pyramids o and u, base c, flattened || a (fig.), large, distinct, ironblack, loose. 2.00





2148° unit prism m, prism g, macropinacoid a, macrodomes h, k and l, several pyramids (similar to

fig.), brilliantly defined short prism, loose. 2.00

2149+ imperfect tables, in pegmatite. 1.00
2150° massive. 2.00

Note:—Normal Columbite, the nearly pure columbate, graduates into normal Tantalite, the nearly pure tantalate.





Type Species No. No.	Tantalite—Continued
2152	disseminated in pegmatite. 1.50
2153	water-worn grains, lot. 1.50
2154	conglomerate of pebbles, ferruginously cemented. 1.50
21550	Manganotantalite (high in Mn), macropinacoid a, brachypinacoid b and base c, all prominent, dull iron-black, large, loose. 3.00

2156+ Manganotantalite, massive. 1.50

2157 526A. SKOGBÖLITE. FeTa<sub>2</sub>O<sub>6</sub>. Orthorhombic, prisms, black. 2.00 II. Neotantalite. Near tantalite in composition. Isometric, octahedral, clear yellow.

Ixiolite. Fe, Mn columbo-tantalate with some Sn. Orthorhombic, rectangular prisms, dark-gray.

2158 II. Stibiotantalite. (SbO)<sub>2</sub>(Ta,Cb)<sub>2</sub>O<sub>6</sub>. Orthorhombic, hemimorphic, adamantine, yellowish. 9.00

2159° crystalline rolled pebble, with tantalite, resinous. 2.50

2160 527. Tapiolite. Fe(Ta,Cb)<sub>2</sub>O<sub>6</sub> where Ta: Cb=4: 1. Tetragonal, square octahedroids, black. 8.00

21610 massive. 4.00


I. Mossite. Fe(Cb,Ta)<sub>2</sub>O<sub>6</sub>. Tetragonal, small twins, tw.pl. e, black.

II. Strüverite. FeO.(TaCb)<sub>2</sub>O<sub>5</sub>.4TiO<sub>2</sub>(?). Tetragonal, ironblack.

# 4. Samarskite Group. Orthorhombic. Range of Hardness 5-6

2162\*528. Yttrotantalite. Essentially RR<sub>2</sub>(Ta,Cb)<sub>4</sub>O<sub>15</sub>+4H<sub>2</sub>O., with R=Fe,Ca; R=Y,Er,Ce,etc. Orthorhombic, prisms.

2163°529. Samarskite.  $R_3R_2(Cb,Ta)_6O_{21}$ , with  $R=Fe,Ca,UO_2,$  etc.; R=Ce and Y metals chiefly. Highly radioactive. Orthorhombic, macropinacoid a, brachypinacoid b and macrodome e, all prominent (similar to fig.) dull but distinct faces, large, loose. 2.50



2164+ massive, splendent velvet-black. 2.50 S. Hydrosamarskite, 10 p.c. H<sub>2</sub>O. 168 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species No. No.

Nohlite. Chiefly U, Y, Fe columbate. Massive, brown. Vietinghofite. An iron-samarskite. Amorphous.

- II. Loranskite. Chiefly Ta<sub>2</sub>O<sub>3</sub>, Y<sub>2</sub>O<sub>3</sub>, Ce<sub>2</sub>O<sub>3</sub>, CaO, FeO, ZrO, H<sub>2</sub>O. Massive, black.
- 530. Annerödite. Essentially U and Y pyro-columbate. Orthorhombic, prisms.

2165° massive, black. 4.00

2166°531. Hielmite. Y, U, Fe, Mn and Ca stanno-tantalate and columbate. Orthorhombic, indistinct crystal, black. 2.00

### Aeschynite Group

Orthorhombic. Range of Hardness 6-6.5

- 2167 532. Æschynite. Chiefly Ce metals columbate and titanate (thorate). Orthorhombic, flat prism, distinct. 2.50 massive, brownish-black. 1.50
- 2169°533. Polymignite. Ce metals, Th, Fe, Ca columbate and titanate (zirconate). Orthorhombic, slender prisms, black.
  - 534. Euxenite. Y, Er, Ce, U columbate and titanate. Highly radio-active. Orthorhombic, prismatic.

2170+ massive, bright vitreous black. 1.50

- 2171 535. Polycrase. Y, Er, Ce, U columbate and titanate. Orthorhombic, prisms tabular || b, black. 3.00
- 2172° II. Epistolite. Containing Cb<sub>2</sub>O<sub>5</sub>, SiO<sub>2</sub>, TiO<sub>2</sub>, Na<sub>2</sub>O, H<sub>2</sub>O, F(?).

  Monoclinic, tabular, pearly-gray. 2.00
  - II. Blomstrandine, Priorite. Y, Er, Ce, U columbate and tantalate. Orthorhombic, tabular, brownish-black.
  - II. Endeiolite. R<sup>II</sup>O.(Cb<sub>2</sub>O<sub>5</sub>)H<sub>2</sub>O.R<sup>II</sup>O.SiO<sub>2</sub>. Cb<sub>2</sub>O<sub>5</sub> 59-93,
     SiO<sub>2</sub> 11·48, ZrO<sub>2</sub> 3·78, Al<sub>2</sub>O<sub>3</sub> 4·43, CaO 7·89, Na<sub>2</sub>O 3·58, H<sub>2</sub>O 4·14. Isometric, minute crystals, dark chocolate-brown.

# Appendix to Columbates, Tantalates

Blomstrandite. Chiefly U tantalo-columbate and titanate. Massive, vitreous black.

2173 Rogersite. Y etc., columbate. Encrusting, white. 1.50

#### Phosphates, Arsenates, Vanadates, Antimonates 4.

# A. Anhydrous Phosphates, Vanadates, Arsenates, **Antimonates**

1. Introductory Subdivision. Hardness 5


Type Species 2174°536. Xenotime. Essentially Y<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>. Tetragonal obtuse unit pyramid z, truncated by narrow unit prism m (fig.). 2.00 prism *m* predominating. 2175 2176+ massive, dull brown. 1.50 2174. Xenotime Hussakite, with small amount SO<sub>3</sub> II. 2177°537. Monazite. Essentially (Ce,La,Di)PO. Monoclinic, flattened || orthopinacoid a, orthodome x also prominent, with prism m, pyramids v and r distinct, opaque dull brown, loose. 1.00 highly modified (fig.), small, brilliantly defined, trans-21789 parent yellowish-brown. 2.00 contact-twin, tw.pl. a, opaque, dull, loose. 1.50 2179 water-worn pebbles, brown (lot). .60 2180 2181+ sand, containing 4 or 5 p.c. ThO<sub>2</sub>. .40 II. Britholite. Ce metals and Ca silicate and phosphate. Orthorhombic, prisms, brown. II. Erikite. Containing SiO<sub>2</sub>, P<sub>2</sub>O<sub>5</sub>, ThO<sub>2</sub>, (Ce, La, Di)<sub>2</sub>O<sub>3</sub>, Al<sub>2</sub>O<sub>3</sub>,

Na<sub>2</sub>O,H<sub>2</sub>O(?). Orthorhombic, prismatic, brown.

-Hardness 5, 6

538. Berzeliite.  $R_3As_2O_8$ , with R=Ca, Mg, Mn. Isometric, trapezohedron n truncated by cube a and dodecahedron d. 21829 massive, resinous yellow. 2.00 5 p.c. Na<sub>2</sub>O. I. Soda-berzeliite.

Pseudoberzeliite. R<sub>3</sub>.As<sub>2</sub>O<sub>8</sub>, with R=Ca, Mg, Mn. Orthorhombic(?). Massive, vellow.



- 170 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species No. No.
  - 539. Monimolite. R<sub>3</sub>Sb<sub>2</sub>O<sub>8</sub>, with R=Pb: Fe=3: 1. Isometric, octahedrons. Varieties:—
    - 1. With Ca.
    - 2. Without Ca.

#### Hardness 3, 2

- 2183 II. Graftonite. R<sub>3</sub>P<sub>2</sub>O<sub>8</sub>, with R=Fe,Mn,Ca. Monoclinic, salmon-pink. 8.00
- 2184°540 Caryinite. R<sub>3</sub>As<sub>2</sub>O<sub>8</sub>, with R=Pb,Mn,Ca,Mg. Monoclinic(?), massive, greasy brown. 2.00
  - 541. Carminite. Pb<sub>3</sub>As<sub>2</sub>O<sub>8</sub>.10FeAsO<sub>4</sub>(?). Orthorhombic, acicular, carmine.

### -----Hardness 4

- 2185 542. Pucherite. Bi<sub>2</sub>O<sub>3</sub>.V<sub>2</sub>O<sub>5</sub>. Orthorhombic, tabular || c, minute, distinct. 2.50
- 21860 minute short needles, adamantine, brown. 2.50

### 2. Triphylite Group. Orthorhombic. Hardness 4.5-5

2187+543. Triphylite. Li(Fe,Mn)PO<sub>4</sub>. Orthorhombic, massive, bluishgray. .60

NOTE:—Triphylite with increasing Fe and decreasing Mn, graduates into Lithiophilite.

- 2188\*544. Lithiophilite. Li(Mn,Fe)PO<sub>4</sub>. Orthorhombic, cleavage, resinous pale yellowish-brown. .60
  - Heterosite. Hyd. Mn, Fe phosphate. Altered triphylite. Cleavages, resinous greenish and bluish-gray, submetallic violet on exposure.
  - Pseudotriplite. Chiefly hyd. Fe phosphate. Altered triphylite. Incrustation.
  - Alluaudite. Hyd. Mn, Fe phosphate. Altered triplite(?). Cleavages, brown.
  - Melanchlor. Hyd. Fe phosphate. Altered triphylite(?). Blackish-green.
  - 545. Natrophilite. Na<sub>3</sub>PO<sub>4</sub>.Mn<sub>3</sub>P<sub>2</sub>O<sub>8</sub>. Orthorhombic, massive cleavable, clear wine-yellow.

Hardness	6,	5
----------	----	---

- 2189 546. Beryllonite. Na<sub>3</sub>PO<sub>4</sub>.Be<sub>3</sub>P<sub>2</sub>O<sub>8</sub>. Orthorhombic, highly complex, colorless. 4.00
- 2190° crystal fragment, transparent. 1.00

Type Species No. No. 21910547. I. Herderite. Monoclinic, small short (CaF) BePO<sub>4</sub>. prism, yellowish-white. -Hardness 4·5  $[Al(OH)_2]_3[SrOH]P_2O_7$ . Rhombohedral, 2192 548. I. Hamlinite. minute, transparent. 8.00 II. Florencite.  $3Al_2O_3.Ce_2O_3.2P_2O_5.6H_2O.$ Rhombohedral, clear pale yellow. 3. Apatite Group. Hexagonal with pyramidal hemihedrism. Hardness 5, 3.5 and 3 Phosphates, Arsenates, Vanadates of calcium and lead, with chlorine and fluorine. 549. Apatite, ordinary or Fluor-apatite, 3Ca<sub>3</sub>P<sub>2</sub> O<sub>8</sub> + CaF<sub>2</sub> and Chlor-apatite, 3Ca<sub>3</sub>P<sub>2</sub> O<sub>8</sub> + CaCl<sub>2</sub>, also intermediate compounds. Hexagonal with pyramidal hemihedrism. 1. Ordinary varieties, crystals bright and of ideal symmetry and perfection:-21939 unit prism m, unit pyramid x (fig.), large, greenish-blue. .50 Apatite m, x with base c (fig.), very large, brown, loose. 2194+ .50 2195 ditto, large green, in calcite. .50 ditto, with second order prism a, truncated 21960 by unit pyramid r and second order pyramid s, transparent pale violetblue, with cassiterite. 1.50 highly modified (fig.), brilliant, clear color-21970 less, with epidote. 2.00 ditto, milky, with adularia. 1.00 2198\* Apatite 2194. thin tabular || base c, unit pyra-21999 mid r (similar to fig.), white, small. 1.00 2200 ditto, truncated by unit prism m, translucent pale red. 1.50 acicular prism, clear colorless, **220I** 

in lava.

2202+

2203

1.50 granular massive, sea-green. .20

2197. Apatite

granular massive, brown. .20

172 Type Species No. No.	COMPLETE TYPE COLLECTION. DANA'S SYSTEM  Apatite—Continued
22040	compact massive, yellowish-
	white20
2205°	Asparagus-stone, unit prism
	m, unit pyramid x (similar 2199. Apatte
	to fig.), clear pale yellowish-green, brilliant. 1.50
	Lasurapatite, sky-blue crystals with lapis.
2206	Francolite, globular groups of small distinct hexagonal
	tables, translucent greenish-white. 1.50
	2. Manganapatite, Mn replaces Ca.
	Cupro-apatite. Contains 20.93 p.c. CuO(?).
2207	3. Fibrous concretionary, Phosphorite60
2208	4. Earthy apatite, Osteolite, impure altered40
	Pseudoapatite, altered pyromorphite.
2209*	Staffelite, botryoidal concentric incrustation, compact
	radio-fibrous, translucent yellowish-green50
	Hydroapatite. A hydrous apatite, mammillary concre-
	tions, chalcedony-like, milky.
22109	Phosphatic Nodules, fossiliferous, impure, gray20
2211+	Phosphate Rock, fossiliferous, whitish20
2212	Phosphate Rock, granular, brown20
22130	Guano, organic origin, earthy, brown20
550	Pyromorphite. 3Pb <sub>3</sub> P <sub>2</sub> O <sub>8</sub> .PbCl <sub>2</sub> . Hexagonal, pyramidal
	hemihedrism. 1. Ordinary varieties:—
22140	(a) unit prism $m$ , base $c$ , bright, sharply symmetrical, brown. 1.00
2215	ditto, translucent pale yellowish-green, small. 1.25
2216+	ditto, dark green75
2217 <sup>©</sup>	ditto, barrel-shaped75
2218	ditto, wax-yellow. 2.00
2219*	ditto, tapering parallel grouping, brown
	(fig.)30
2220	(b) acicular, brown. 1.50
222Iº	moss-like group, brown. 1.00
2222	(c) concretionary group. 1.00
	(d) fibrous.
2223+	(e) granular massive75
	(f) earthy, incrusting. 2219. Pyromorphite
	2. Polysphærite, contains CaO. Globular groups.
2224	3. Chromiferous, short acicular, bright orange. 2.50
	4. Arseniferous, pale green.

Type Species No. No.	Pyromorphite—Continued
2225°	altered to Galena. 1.50
	Svabite. H <sub>2</sub> O.10CaO.3As <sub>2</sub> O <sub>3</sub> . Six-sided prisms. 2.50
	Mimetite. 3Pb <sub>3</sub> As <sub>2</sub> O <sub>8</sub> .PbCl <sub>2</sub> . Hexagonal, pyramidal hemi-
	hedrism. 1. Ordinary:—
2227+	(a) minute groups of prisms, pale yellowish-brown. 1.50
2228	minute globular groups, yellow. 1.50
	(b) capillary, somewhat asbestiform.
	(c) concretionary.
	2. Calciferous.
2229°	3. Campylite. 3.34 p.c. P <sub>2</sub> O <sub>5</sub> . Nearly spherical barrel-
	shaped hexagons, resinous brownish-red, small,
	distinct. 2.00
2230+	Endlichite. Nearly equal amounts of Pb <sub>3</sub> As <sub>2</sub> O <sub>8</sub> and
	$Pb_3V_2O_8$ with PbCl (between Mimetite and Vanadinite). Hexagonal, unit prism $m$ , base $c$ , ada-
	mantine, transparent straw-yellow, ideal symmetry,
	small. 1.00
2231	ditto, brownish. 1.00
2232	bi-colored slender unit prism $m$ , clear straw-yellow,
2232	termination red and etched, loose (12)50
2233°	unit prism $m$ , unit pyramids $x$ and $y$ , base $c$ , red, loose
<b>22</b> 33°	(3)50
2234	spherical groups, pale yellow. 1.00
2235°	massive, orange. 1.50
	Vanadinite, 3Pb, V.O., PbCl., Hexagonal
· ·	pyramidal hemihedrism, unit prism
	m, base $c$ , truncated by unit pyra-
	mid $x$ and dihexagonal pyramid $u$
	(fig.), minute, ideal symmetry, ada-
	mantine, translucent red. 1.00
2237	unit prism $m$ , unit pyramids $x$ and
	y, base c, perfect, clear yellow-
	ish-red, minute. 1.50
2238*	hollow prisms in tapering groups
	(fig.), distinct, bright red,
2220	loose (6). 1.00 ditto, brown on descloizite. 1.50
2239 2240º	barrel-shaped prism $m$ , base $c$ ,
22409	ideal symmetry, adamantine,
	brown, small. 1.00 2238. Vanadinite
	210 min, Simula. 1.00 szja. Vassamite

Type Species No. No.	Vanadinite—Continued
No. No. 2241	acicular, clear brownish-yellow, small. 1.00
2241 2242°	globular incrustation, resinous brownish-yellow. 1.50
•	Hedyphane. A calcium-mimetite. Hexagonal, highly
1	
	complex pyramidal.
22430	massive, resinous, whitish. 1.50
11.	. Georgiadésite. Pb <sub>3</sub> (AsO <sub>4</sub> ) <sub>2</sub> .3PbCl <sub>2</sub> . Orthorhombic, white.
4. Wa	gnerite Group. Monoclinic. (RF) RPO4
	Range of Hardness 4—5
2244 553.	Wagnerite. Mg <sub>3</sub> P <sub>2</sub> O <sub>8</sub> .MgF <sub>2</sub> . Monoclinic, complex. 4.00
2245	Kjerulfine, large rough crystal. 3.00
2246*	Kjerulfine, massive, pale yellowish. 2.00
·	Cryphiolite. P <sub>2</sub> O <sub>5</sub> 47·59, MgO 33·72, CaO 14·74. Mono-
	clinic, tabular   a, small, clear honey-yellow, in lava.
554.	Spodiosite. Ca <sub>3</sub> P <sub>2</sub> O <sub>8</sub> .CaF <sub>2</sub> (?). Orthorhombic(?), flattened
	b, prisms, grayish.
2247*555.	Triplite. Fe, Mn, Ca, Mg phosphate, with F. Monoclinic,
	massive, resinous-brown50
	Zwieselite, Fe and Mn only, clove-brown.
	Talktriplite, much Mg and Ca, grains, yellowish.
2248°	Griphite. Mn, Al, Ca, Na, Fe phosphate. Massive, resin-
•	ous blackish-brown40
	Sarcopside. Impure altered trip-
	lite(?).
2249°556.	Triploidite. 4(Mn,Fe)O.P <sub>2</sub> O <sub>5</sub> .H <sub>2</sub>
1,7	O. Monoclinic, crystalline,
	clear yellowish. 2.00
S. I.	Adelite. (MgOH) CaAsO <sub>4</sub> . Mono-
	clinic, grayish.
I.	Tilasite (Fluor-Adelite). (Mg.F)
	CaAsO <sub>4</sub> . Massive, granular.

2250°557. Sarkinite. 4MnO.As<sub>2</sub>O<sub>5</sub>.H<sub>2</sub>O. Monoclinic, elongated || axis b, flattened || a, minute, rose-red. 2.50

# 5. Amblygonite Group

Monoclinic, Triclinic. Hardness 5 and 6

2251°558. Durangite. AlAsO<sub>4</sub>.NaF. Monoclinic, oblique pyramids m and  $\pi$  predominating, (fig.) small, distinct, orange-red, loose (6). 1.00

Type Species

559. Amblygonite. AlPO<sub>4</sub>.LiF. Triclinic, large coarse crystal. cleavage, white. .50 2252+

> S. Morinite. Contains H<sub>2</sub>O,F,P<sub>2</sub>O<sub>5</sub>,Al<sub>2</sub>O<sub>3</sub>,Na<sub>2</sub>O. An amblygonite alteration-product. Monoclinic, crystals.

# B. Acid and Basic Phosphates, Arsenates, Etc.

Hardness 3.5

2253 560. Monetite. 2CaO.P<sub>2</sub>O<sub>5</sub>.H<sub>2</sub>O. Triclinic, clear yellowishwhite. 1.00 Natrophite. HNa<sub>2</sub>PO<sub>4</sub>.

### Olivenite Group. Orthorhombic. Range of hardness 3-4

2254+561. Olivenite. 4CuO.As,Os.H.O. Orthorhombic, octahedroid, unit prism m and brachydome e prominent, ideal symmetry, adamantine, blackish-green, small. 1.50

unit prism m, macro- and brachypinacoids 2255° a and b, macro- and brachydomes vand e (fig.). 1.50

acicular, clear olive-green, small. 1.50 2256 2257° fibrous diverging, concentric, green-

> ish. 2.00

earthy felt-like mass, whitish. 2.00 2258 2259\*562. Libethenite. 4CuO.P.Os.H.O. Orthorhombic, octahedroid, unit prism m and brachydome e predominating (fig.), minute, ideal symmetry, brilliant, dark green. 2.00


2260 563. Adamite. 4ZnO.As,O<sub>5</sub>.H,O. Orthorhombic, prismatic || axis b by extension of macrodome d, terminated by several prisms and brachypinacoid b (fig.), small, ideal sym-

> metry, brilliant translucent green. 2.00

2259. Libethenite

2255.

Olivenite



ditto, colorless, minute. 1.00

2261

176 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species Adamite-Continued 2262\* drusy incrustation, bright green. II. Tarbuttite. 4ZnO.P<sub>2</sub>O<sub>5</sub>.H<sub>2</sub>O. Triclinic, striated crystals, transparent. 564. Descloizite. 4RO.V<sub>2</sub>O<sub>5</sub>. H<sub>2</sub>O., with R=Pb, Zn chiefly. Orthorhombic, prismatic. 976 pyramid o predominating (fig.), 2263\* minute, ideal symmetry, brilliant, dark brown. 1.50 drusy globular, crystalline, red. 2264 Descloizite 1.00 mammillary crust, radio-fibrous, brownish-red. 1.00 2265 22669 Cuprodescloizite, drusy botryoidal, dull greenish-black. 1.00 Eusynchite. Massive descloizite(?). Dechenite. PbO.V<sub>2</sub>O<sub>5</sub>(?). Massive. 2267°565. Calciovolborthite. 4(Cu,Ca)O.V<sub>2</sub>O<sub>5</sub>,H<sub>2</sub>O(?). Rosette-like aggregates of small thin scales, pearly green. 4.00 fine crystalline granular, gray. 2268 566. Brackebuschite. R<sub>3</sub>V<sub>2</sub>O<sub>8</sub>+H<sub>2</sub>O., with R=Pb chiefly, also Fe, Mn.(?). Monoclinic(?), small flat prisms, black. 2.50 2269 567. Psittacinite.  $4RO.V_2O_5.2H_2O_7$ , with R=Pb: Cu=1:1(?). Cryptocrystalline coating, green. 6.00 Mottramite (Psittacinite?). Pb and Cu vanadate. Crystalline incrustation, resinous velvety-black. -Range of Hardness 3-4.5 2270 568. Erinite. 5CuO.As,Os.2H2O. Crystalline groups, concentric mammillary, fibrous structure, fine emeraldgreen. 2.00 2271 569. Dihydrite. 5CuO.P,O,.2H,O. Monoclinic or triclinic, hemispherical aggregates of small crystals, adamantine, dark emerald-green. 2272°570. Pseudomalachite. In part 6CuO.P<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. Massive, reniform radio-fibrous, dark emerald-green. 1.50 Ehlite. 5CuO.P<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. 1.50 2273 2274 571. Clinoclasite. 6CuO.As<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. Monoclinic, minute prisms, vitreous dark green. hemispherical radio-fibrous. 2.50 2275\*

572. Chondrarsenite. Perhaps 6MnO.As<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. Embedded grains, translucent yellow.

Xantharsenite. Essentially 5MnO.As,O5.5H2O(?).

-Range of Hardness 3.5—5

(Arseniosiderite 1-2)

Partly  $2\text{Fe}_2\text{O}_3$ .  $P_2\text{O}_5$ .  $3\text{H}_2\text{O}$ . 2276 573. Dufrenite. Orthorhombic, drusy radio-fibrous. .75

diverging fibro-columnar, blackish-green. .50 2277+


2278 574. Lazulite.  $(Fe, Mg)O.Al_2O_3.P_2O_5.H_2O.$ Monoclinic, unit pyramids  $\phi$  and e, ideal symmetry, azure-blue. .75

ditto, with orthodome t, flattened by extension of one 2279 pair of pyramidal planes. .75

2280+ contact-twins, tw.axis c (fig.). 22810 massive, pale greenish-blue. 1.00

I. Gersbyite. P<sub>2</sub>O<sub>5</sub> 32·26, Al<sub>2</sub>O<sub>3</sub> 46·68, CaO,

- FeO, MnO 6.66, MgO 5.33, H<sub>2</sub>O 9.07=100. Grains, blue.
- 3 Ca O. Al<sub>2</sub> O<sub>3</sub>. P<sub>2</sub> O<sub>5</sub>. 3 H<sub>2</sub> O. 575. Tavistockite. Microscopic acicular crystals, pearly white.
- 576. Cirrolite. 6 CaO. 2 Al<sub>2</sub>O<sub>3</sub>. 3 P<sub>2</sub>O<sub>5</sub>. 3 H<sub>2</sub>O(?). Compact, pale yellow.
- 2282°577. Arseniosiderite. 6 CaO. 4 Fe<sub>2</sub>O<sub>3</sub>. 3 As<sub>2</sub>O<sub>5</sub>.9 H<sub>2</sub>O. Tetragonal or hexagonal(?), fibro-lamellar concretion, silky golden-brown. 1.50



2280. Lazulite

- I. Retzian. Mn, Ca and rare earths basic arsenate. Orthorhombic, prismatic, dark-brown.
- 2283°578. Allactite. 7MnO.As<sub>2</sub>O<sub>5.4</sub>H<sub>2</sub>O. Monoclinic, tabular || a, minute but distinct, adamantine, translucent pale red. 2.00
- 2284°579. Synadelphite.  $2(Al,Mn)AsO_4.5Mn(OH)_2$ Monoclinic. minute sharp pyramids, bright brownish-black. 4.00
  - I. Basiliite. 11 (Mn<sub>2</sub>O<sub>3</sub>.Fe<sub>2</sub>O<sub>3</sub>)Sb<sub>2</sub>O<sub>5</sub>.21H<sub>2</sub>O. Foliated, steelblue.
  - 580. Flinkite. 4MnO.Mn<sub>2</sub>O<sub>3</sub>.As<sub>2</sub>O<sub>5</sub>.4H<sub>2</sub>O. Orthorhombic, thin tabular || c, minute, transparent greenish-brown.
  - (AlMn) AsO<sub>4</sub>.4Mn(OH)<sub>2</sub>. Rhombohedral, 581. Hematolite. rhomboids, red, blackening on the surface.

Type Species No. No. 2285°582. Arseniopleite. 9RO.R<sub>2</sub>O<sub>3</sub>.3As<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O, with R=Mn,Ca also Pb,Mg; R=Mn also Fe. Rhombohedral (?), massive cleavable, brownish-red. 2.00

583. Manganostibiite. 10MnO.Sb<sub>2</sub>O<sub>5</sub>(?). Orthorhombic(?), compact, black.

Ferrostibian and Stibiatil. Mn, Fe antimonates. Monoclinic(?), black.

2286°584. Atelestite. 3Bi<sub>2</sub>O<sub>3</sub>.As<sub>2</sub>O<sub>5</sub>.2H<sub>2</sub>O. Monoclinic, tabular || a, minute, adamantine, clear sulphur-yellow. 2.00

### C. Hydrous Phosphates, Arsenates, Etc.—Normal Division Range of Hardness 2—2.5

2287°585. Struvite. NH<sub>4</sub>MgPO<sub>4</sub>+6H<sub>2</sub>O. Orthorhombic, hemimorphic, macrodomes s s<sub>1</sub>, brachypinacoid b, base c (similar to fig.), distinct, loose. .50

2288

unit prism m, macrodome s, base c, small, loose (3). .50

Guano Minerals: See "System of Mineralogy" for brief reference to numerous doubtful compounds.

II. Dittmarite. MgNH<sub>4</sub>PO<sub>4</sub>.2Mg<sub>2</sub>H<sub>2</sub> (PO<sub>4</sub>)<sub>2</sub>+8H<sub>2</sub>O. Orthorhombic(?), transparent.

II. Schertelite. Mg(NH<sub>4</sub>)<sub>2</sub>H<sub>2</sub>(PO<sub>4</sub>)<sub>2</sub>+4H<sub>2</sub>O. Small crystals, transparent.

586. Collophanite. 3CaO.P<sub>2</sub>O<sub>5</sub>.H<sub>2</sub>O. Amorphous, opaline.

587. Hopeite.  $Zn_3P_2O_8 + H_2O(?)$ . Orthorhombic, minute prisms.

—Hardness 4—4·5

2287. Struvite

II. Parahopeite. 3ZnO.P<sub>2</sub>O<sub>5</sub>.4H<sub>2</sub>O. Triclinic, striated crystals, transparent.

588. Dickinsonite.  $3R_3P_2O_8 + H_2O$  with R=Mn,Fe,Na<sub>2</sub>, chiefly also Ca,K<sub>2</sub>,Li<sub>2</sub>. Monoclinic, pseudo-rhombohedral tables, green.

589. Fillowite.  $3R_3P_2O_8 + H_2O$ , with R=Mn: Fe(+Ca): Na<sub>2</sub>= 6:2:1(?). Monoclinic, pseudo-rhombohedral cuboid.

#### Roselite Group.

Type Species No. No. Triclinic. Hardness 3.5, 5 and 3.5

- 2289°590. Roselite. (Ca,Co,Mg)<sub>3</sub>As<sub>2</sub>O<sub>8</sub>.2H<sub>2</sub>O. Triclinic, minute complex crystal, glassy translucent dark rose-red. 2.50
- 2290°591. Brandtite. 2CaO.MnO.As<sub>2</sub>O<sub>5</sub>.2H<sub>2</sub>O. Triclinic, highly modified, prismatic by development of several brachydomes, base c prominent, minute, divergent groups, vitreous white. 1.50
  - 592. Fairfieldite. Ca<sub>2</sub>MnP<sub>2</sub>O<sub>8</sub>+2H<sub>2</sub>O. Triclinic, prisms, white.

    Range of Hardness 3-3.5
- 2291 593. Messelite. (Ca,Fe)<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>+2½H<sub>2</sub>O. Triclinic, minute indistinct tables. .75
  - II. Anapäite. (Ca, Fe)<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>.4H<sub>2</sub>O. Triclinic, tabular, greenish-white.

#### -----Hardness 3.5

- 594. Reddingite. Mn<sub>3</sub>P<sub>2</sub>O<sub>8</sub>+3H<sub>2</sub>O. Orthorhombic, octahedroids, clear whitish.
- 595. Picropharmacolite. R<sub>3</sub>As<sub>2</sub>O<sub>8</sub>+6H<sub>2</sub>O, with R=Ca,Mg. Spherical, radio-foliated, white.
  - -----Hardness 2.5
- 596. Trichalcite. Cu<sub>3</sub>As<sub>2</sub>O<sub>8</sub>+5H<sub>2</sub>O. Radio-columnar groups, silky verdigris-green.
- 2292 Lavendulan. Hyd. Cu arsenate with Co and Ni. Amorphous, lavender-blue. 2.00
  - Chlorotile. Cu<sub>3</sub>As<sub>2</sub>O<sub>8</sub>+6H<sub>2</sub>O. Orthorhombic, minute capillary.

#### Vivianite Group. Monoclinic. Range of Hardness 1-2.5

- 2293 597. Vivianite. Fe<sub>3</sub>P<sub>2</sub>O<sub>8</sub>+8H<sub>2</sub>O. Monoclinic, large sharply defined prism, flattened || a, translucent dark blue, brilliant. 3.00
- 2294+ ditto, dull. .75
- 2295 ditto, rounded lenticular. .75
- 2296° stellated group in pyrite. .75
- 2297° acicular, replacing fossils. .75
- 2298°598. Symplesite. Fe<sub>3</sub>As<sub>2</sub>O<sub>8</sub>+8H<sub>2</sub>O(?). Monoclinic, small prisms, translucent greenish. 2.00

- 599. Bobierrite. Mg<sub>3</sub>P<sub>2</sub>O<sub>8</sub> + 8H<sub>2</sub>O. Monoclinic, microscopic prisms, white, in guano.
  - I. Hautefeuillite. (Mg,Ca)<sub>3</sub>P<sub>2</sub>O<sub>8</sub>+8H<sub>2</sub>O. Monoclinic, lamellar masses, radiated, colorless.
- 600. Hærnesite. Mg<sub>3</sub>As<sub>2</sub>O<sub>8</sub> + 8H<sub>2</sub>O. Monoclinic, prismatic, flexible folia, white.
- 2299 601. Erythrite. Co<sub>3</sub>As<sub>2</sub>O<sub>8</sub>+8H<sub>2</sub>O. Monoclinic, acicular, translucent purplish-red. 2.00
- 2300 ditto, globular, drusy surface. 1.50
- 2301° foliated-columnar, stellated. 1.50
- 2302+ earthy, Cobalt Bloom, dull purplish-red. 1.00
- 2303°602. Annabergite. Ni<sub>3</sub>As<sub>2</sub>O<sub>8</sub>+8H<sub>2</sub>O. Monoclinic, earthy, applegreen. 1.00
- 2304°603. Cabrerite. (Ni,Mg)<sub>3</sub>As<sub>2</sub>O<sub>8</sub>+8H<sub>2</sub>O. Monoclinic, minute distinct prisms, flexible folia, clear brilliant applegreen. 4.00
  - 604. Köttigite. Zn<sub>3</sub>As<sub>2</sub>O<sub>8</sub>+8H<sub>2</sub>O. Co and Ni replace some Zn. Monoclinic, light red.

#### -Hardness 3.5

- 605. Rhabdophanite. RPO₄ + H₂O, with R=La, Di, Y. Massive, brown.
- 606. Churchite. CePO<sub>4</sub>+4H<sub>2</sub>O. Monoclinic(?), minute crystals, pale reddish-gray.

#### Scorodite Group. Orthorhombic. Hardness 3.5

- 2305 607. Scorodite. Fe<sub>2</sub>O<sub>3</sub>.As<sub>2</sub>O<sub>5</sub>.4H<sub>2</sub>O. Orthorhombic, octahedroid, unit pyramid p prominent, sharply symmetrical, vitreous translucent bluish-green, small. 3.00
- 2306\* ditto, minute, pale leek-green. 1.00
- 2307°608. Strengite. Fe<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.4H<sub>2</sub>O. Orthorhombic, drusy globular, radio-fibrous, red. 2.00
  - Range of Hardness 3.5—5.5
  - II. Purpurite. 2(Fe,Mn)PO<sub>4</sub>+H<sub>2</sub>O. Orthorhombic(?), massive, reddish-purple.
  - 609. Phosphosiderite. Fe<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.3½H<sub>2</sub>O. Orthorhombic, prisms, b prominent, clear reddish.
  - 610. Barrandite. (AlFe)<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.4H<sub>2</sub>O. Spheroidal concretions, grayish.

- 2308\*611. Variscite. Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.4H<sub>2</sub>O. Orthorhombic, drusy globular incrustation, translucent deep apple-green. .75
- 2309° massive, opaque pale green, precious. 1.50
  - Planerite. Chiefly Al hyd. phosphate. Subcrystalline layers in rock, green.
  - 612. Callainite. Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.5H<sub>2</sub>O. Massive, wax-like, translucent mottled green.
  - 613. Zepharovichite. AlPO<sub>4</sub>.3H<sub>2</sub>O. Crystalline, whitish.
- 2310°614. Koninckite. Fe<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.6H<sub>2</sub>O. Spherical, radiated, transparent yellow. 1.50
  - I. Minervite. Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.7H<sub>2</sub>O. Massive, plastic.
  - II. Gorceixite. BaO.2Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.5H<sub>2</sub>O. Pebbles, white.

#### Hydrous Phosphates, Etc.—Acid Division. Hardness 2

- 615. Stercorite. HNa(NH<sub>4</sub>)PO<sub>4</sub>+4H<sub>2</sub>O. Monoclinic, crystalline masses, clear whitish.
- 2311°616. Haidingerite. 2CaO.As<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. Orthorhombic, minute' crystals, small botryoidal groups, clear whitish. 3.00

### Pharmacolite Group. Monoclinic. Hardness 2-2.5

- 2312\*617. Pharmacolite. 2CaO.As<sub>2</sub>O<sub>5</sub>.5H<sub>2</sub>O. Monoclinic, minute needles, stellated, white. 1.25
  - 618. Brushite. 2CaO.P<sub>2</sub>O<sub>5</sub>.5H<sub>2</sub>O. Monoclinic, small prisms, pearly clear whitish.
    - II. Stoffertite, 2 CaO.P<sub>2</sub>O<sub>5</sub>.6½ H<sub>2</sub>O.
      - Range of Hardness 2.5-3 (Hureaulite 5)
  - 619. Metabrushite. 2CaO.P<sub>2</sub>O<sub>5</sub>.4H<sub>2</sub>O. Monoclinic, imperfect crystals, yellowish-white.
  - 620. Martinite. 5CaO.P<sub>2</sub>O<sub>5</sub>.½H<sub>2</sub>O. Rhombohedral, microscopic rhombs, clear whitish.
- 2313 621. Newberyite. 2MgO.P<sub>2</sub>O<sub>5.7</sub>H<sub>2</sub>O. Orthorhombic, composite tabular crystal built of distinct individuals (tabular || a), arranged parallel, vitreous translucent gray, loose. 1.00
- 2314° cavernous group of preceding composite tables. .50
- 2315 622. Wapplerite. 2CaO.As<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O. Monoclinic (or triclinic), crystalline incrustation, white. 1.50

- 182 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species No. No.
  - Rösslerite. HMgAsO<sub>4</sub>+7H<sub>2</sub>O. Crystalline plates, whitish.
  - 623. Hannayite. (NH<sub>4</sub>)<sub>2</sub>O.3MgO.2P<sub>2</sub>O<sub>5</sub>.10H<sub>2</sub>O. Triclinic, small slender prisms, yellowish.
  - 624. Hureaulite. 5MnO.2P<sub>2</sub>O<sub>5</sub>.5H<sub>2</sub>O. Monoclinic, short prisms, clear glassy reddish.
- 2316 625. Forbesite. H<sub>2</sub>(Ni,Co)<sub>2</sub>As<sub>2</sub>O<sub>8</sub>+8H<sub>2</sub>O. Fibro-crystalline, whitish. 4.00
  - II. Palmerite. HK<sub>2</sub>Al<sub>2</sub>(PO<sub>4</sub>)<sub>3</sub>.7H<sub>2</sub>O.

## Hydrous Phosphates, Etc.—Basic Division

Hardness 1.5-3


- 626. Isoclasite. 4CaO.P<sub>2</sub>O<sub>5</sub>.5H<sub>2</sub>O. Monoclinic, minute dull crystals, whitish.
- 627. Hemafibrite. 6MnO.As<sub>2</sub>O<sub>5</sub>.5H<sub>2</sub>O. Orthorhombic, prisms, red, blackening.

Range of Hardness 3—5 (Tyrolite and Chalcophyllite soft, Turquois 6)

- 2317\*628. Conichalcite. 4(Cu,Ca)O.As<sub>2</sub>O<sub>5</sub>.1½H<sub>2</sub>O. Massive globular, vitreous emerald-green. 1.00
- 2318°629. Bayldonite. 4(Pb,Cu)O.As<sub>2</sub>O<sub>5.2</sub>H<sub>2</sub>O. Minute mammillary concretions, drusy, resinous green. 3.00
  - 630. Tagilite. 4CuO.P<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. Monoclinic, green.
- 2319 631. Leucochalcite. 4CuO.As<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O(?). Acicular, silky greenish-white. 1.00
- 2320\*632. Euchroite. 4CuO.As<sub>2</sub>O<sub>5</sub>.7H<sub>2</sub>O. Orthorhombic, small distinct octahedroids, vitreous emerald-green. 2.00
  - 633. Volborthite. (Cu,Ca,Ba)<sub>3</sub>(OH)<sub>3</sub>VO<sub>4</sub>+6H<sub>2</sub>O(?). Minute six-sided tables.
- 2321 incrustation, green. 3.00
  - 634. Cornwallite. 5CuO.As<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. Massive, green.
- 2322+635. Tyrolite. Perhaps 5CuO.As<sub>2</sub>O<sub>3</sub>.9H<sub>2</sub>O. Orthorhombic, fanshaped foliations, green. 1.00
- 2323 636. Chalcophyllite. 7CuO.As<sub>2</sub>O<sub>5</sub>.14H<sub>2</sub>O(?). Rhombohedral, small six-sided tables, rhombohedral, helpedrang bases (6x) possible.


bohedron r, base c (fig.), pearly verdigris-green. 3.00

2324\* foliated massive, emerald-green.
2.00



- 637. Veszelyite. (CuZn)<sub>7</sub>(OH)<sub>8</sub>(As,P)<sub>2</sub>O<sub>8</sub>+5H<sub>2</sub>O. Monoclinic (or triclinic?), incrustation, greenish-blue.
- 2325°638. Ludlamite. 7FeO.2P<sub>2</sub>O<sub>5</sub>.9H<sub>2</sub>O. Monoclinic, tabular || c, minute, distinct, vitreous pale green. 3.00
- 2326 639. Wavellite. 3Al<sub>2</sub>O<sub>3</sub>.2P<sub>2</sub>O<sub>3</sub>.12H<sub>2</sub>O. Orthorhombic, crystal terminations forming surface of radio-fibrous hemispheres, bright green. 2.00
- 2327° globular, radio-fibrous, yellowish-white. .75
- 23289 stalactitic, radio-fibrous, grayish-white. 1.50
- 2329+ stellated fibrous, bright green. .40
- 2330 stellated fibrous, grayish. .75
- 2331 reniform, chalcedony-like, brownish. .75
  - 640. Fischerite. 2Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O. Orthorhombic, minute crystals, green.
  - 641. Peganite. 2Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.6H<sub>2</sub>O. Orthorhombic, indistinct prisms, greenish.
- 2332+642. II. Turquois. [Al(OH)<sub>2</sub>.Fe(OH)<sub>2</sub>.Cu(OH).H]<sub>3</sub>PO<sub>4</sub>. Massive in matrix, sky-blue, precious. .75
- 2333° massive, greenish. .50
- 2334 massive, grayish. .50
- 2335° I. Wardite. 2Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.4H<sub>2</sub>O. Massive, concretionary, light green. 1.25
  - 643. Sphærite. 5Al<sub>2</sub>O<sub>3</sub>.2P<sub>2</sub>O<sub>5</sub>.16H<sub>2</sub>O(?). Globular concretions.
- 2336°644. Liskeardite. 3(Al,Fe)<sub>2</sub>O<sub>3</sub>.As<sub>2</sub>O<sub>5</sub>.16H<sub>2</sub>O. Microscopic needles on fibrous incrustation, white. 2.00
- 2337°645. Evansite. 3Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.18H<sub>2</sub>O. Massive, white. 1.00
- 2338 Cœruleolactite. 3Al<sub>2</sub>O<sub>3</sub>.2P<sub>2</sub>O<sub>5</sub>.10H<sub>2</sub>O(?). Cryptocrystalline, pale sky-blue. .40
  - Taranakite. Al, K, Fe hyd. phosphate. Massive, yellowish-white.
  - Berlinite. 2Al<sub>2</sub>O<sub>3</sub>.2P<sub>2</sub>O<sub>5</sub>.H<sub>2</sub>O. Compact.
  - Trolleite. 4Al<sub>2</sub>O<sub>3</sub>.3P<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. Compact, pale green.
  - I. Augelite. 2Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. Monoclinic, tabular, red. Attacolite. Al, Mn, Ca, Fe hyd. phosphate. Massive, red.
- 2339\*646. Pharmacosiderite. 4Fe<sub>2</sub>O<sub>3</sub>.3As<sub>2</sub>O<sub>5</sub>.15H<sub>2</sub>O(?). Isometric, tetrahedral, minute distinct bright cubes, translucent brown. 1.50
- 2340° ditto, small, green. 2.50
- 2341 cube a, tetrahedron o, distinct. 4.00

- 184 COMPLETE TYPE COLLECTION. DANA'S SYSTEM
  Type Species
  No. No.
- 2342°647. Cacoxenite. 2Fe<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>3</sub>.12H<sub>2</sub>O. Radiated tufts, brownish-yellow. 1.00
- velvety incrustation, drusy botryoidal. 1.00
  - II. Kertschenite. Hyd. basic ferric phosphate, fibrous, dark green.
- 2344°648. Beraunite. 3Fe<sub>2</sub>O<sub>3</sub>.2P<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O. Monoclinic, drusy incrustation. 1.00
- Eleonorite, small tabular prisms, brownish-red. 1.50 Globosite. Chiefly hyd. Fe fluo-phosphate. Globular. Picite. Chiefly hyd. Fe phosphate. Amorphous, brown. Delvauxite. 2Fe<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.24H<sub>2</sub>O.
- 2346\*649. Childrenite. (Fe,Mn)Al(OH)<sub>2</sub>.PO<sub>4</sub>+2H<sub>2</sub>O, (Fe predominates). Orthorhombic, unit prism *m*, pyramid *r*, minute, brilliantly defined, translucent brown. 1.50
- 2347 650. Eosphorite. (Mn,Fe)Al(OH)<sub>2</sub>PO<sub>4</sub>+2H<sub>2</sub>O, (Mn predominates). Orthorhombic, indistinct minute prisms in crystalline mass, translucent yellowish. 5.00
  - ----Range of Hardness 2.5-4.5
- 2348 651. Mazapilite. 3CaO.2Fe<sub>2</sub>O<sub>3</sub>.2As<sub>2</sub>O<sub>5</sub>.6H<sub>2</sub>O. Orthorhombic, small prisms, sharply defined, black. 4.00
  - 652. Calcioferrite. 6CaO.3Fe<sub>2</sub>O<sub>3</sub>.4P<sub>2</sub>O<sub>5</sub>.19H<sub>2</sub>O. Monoclinic(?), foliated mass.
  - 653. Borickite. Ca<sub>3</sub>Fe<sub>2</sub>(PO<sub>4</sub>)<sub>4</sub>.12Fe(OH)<sub>3</sub>+6H<sub>2</sub>O(?). Reniform massive, reddish-brown.
- 2349 Richellite.  $4\text{FeP}_2\text{O}_8$ .  $\text{Fe}_2\text{OF}_2(\text{OH})_2 + 36\text{H}_2\text{O}$ . Massive, yellow. 1.00
- 2350\*654. Liroconite. 18CuO.4Al<sub>2</sub>O<sub>3</sub>.5As<sub>2</sub>O<sub>5</sub>.
  55H<sub>2</sub>O(?). Monoclinic, thin rhombic octahedroids, unit prism *m*, clinodome *e* (fig.), small, sharply symmetrical, translucent bright blue. 2.00



- I. Kehoeite. ZnO.4Al<sub>2</sub>O<sub>3</sub>.5P<sub>2</sub>O<sub>5</sub>.9H<sub>2</sub>O. Amorphous, massive.
- 655. Chenevixite. 2CuO.Fe<sub>2</sub>O<sub>3</sub>.As<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O(?). Massive, greenish.
- Henwoodite. Chiefly hyd. Al, Cu phosphate. Botryoidal, crystalline structure, turquois-blue. 2.00

- 2352°656. Chalcosiderite. CuO.3Fe<sub>2</sub>O<sub>3</sub>.2P<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O. Triclinic, minute distinct crystals in sheaf-like groups, vitreous, translucent dark green. 1.25
  - Andrewsite. 5Fe<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.5H<sub>2</sub>O. Radio-globular disks, bluish-green.
  - 657. Goyazite. 3CaO.5Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.9H<sub>2</sub>O. Tetragonal or hexagonal, rounded grains, clear whitish.
- 2353°658. Plumbogummite. PbO.2Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.9H<sub>2</sub>O(?). Hexagonal, botryoidal, gum-like, translucent brownish, with pyromorphite. 6.00
- ditto, grayish-white on schist. 9.00

#### Uranite Group.

Hardness 2-2.5 (Walpurgite 3.5, Rhagite 5)

- 2355+659. Torbernite. CuO.2UO<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O. Tetragonal, thick square tables, minute, sharply defined, pearly emerald-green. 1.50
- 2356 ditto, small, extremely thin, transparent. 2.50
- 2357° ditto, microscopic, yellowish-green. 1.50
- 2358°660. Zeunerite. CuO.2UO<sub>3</sub>.As<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O. Tetragonal, thick square tables, minute but distinct, pearly emerald-green. 3.00
- 2359+661. Autunite. CaO.2UO<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O. Orthorhombic, thin square tables, minute, pearly sulphur-yellow. 1.25
- 2360° foliated aggregate, micaceous. 2.50
  - 662. Uranospinite. CaO.2UO<sub>3</sub>.As<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O(?). Orthorhombic, thin square tables, siskin-green.
- 2361°663. Uranocircite. BaO.2UO<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O. Orthorhombic, very thin square tables, pearly translucent yellow-green, small. 3.00
  - 664. Phosphuranylite. 3UO<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.6H<sub>2</sub>O. Pulverulent incrustation of microscopic rectangular scales, pearly lemon-yellow.
  - 665. Trögerite. 3UO<sub>3</sub>.As<sub>2</sub>O<sub>5</sub>.12H<sub>2</sub>O. Monoclinic, druses of thin crystals, tabular || b, pearly lemon-yellow.
    - Fritzscheite. A mangan-uranite with some V. Squarish tables, pearly red.
- 2362 666. Walpurgite. 5Bi<sub>2</sub>O<sub>3</sub>.3UO<sub>3</sub>.2As<sub>2</sub>O<sub>5</sub>.12H<sub>2</sub>O(?). Triclinic, scale-like crystals, yellow. 2.00

- 186 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species No. No.
- 2363° I. Carnotite. K<sub>2</sub>O.U<sub>2</sub>O<sub>3</sub>.V<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O(?). Highly radio-active. Microscopic crystals, scale-like, bright canary-yellow. 2.00
- 2364 amorphous pulverulent mass. 4.00 2365+ ditto, disseminated in sandstone. 1.50
  - 667. Rhagite. Perhaps  $5Bi_2O_3.2As_2O_5.9H_2O$ . Smooth crystalline aggregates, yellowish.
- 2366°668. Mixite. Perhaps 20CuO.Bi<sub>2</sub>O<sub>3</sub>.5As<sub>2</sub>O<sub>5</sub>.22H<sub>2</sub>O. Minute acicular tufts, bright green. 1.50
- incrustation, dull green. 1.00

#### Antimonates; Also Antimonites, Arsenites

A number of antimonates are included among the phosphates, arsenates, etc. Hardness 6 and 4

- 669. Atopite. Perhaps 2CaO.Sb<sub>2</sub>O<sub>5</sub>. Isometric, octahedrons. Schneebergite. Chiefly Ca and Sb. Isometric, microscopic octahedrons, clear honey-yellow.
- 2368+670. Bindheimite. Hyd. Pb antimonate. Amorphous, minutely curved-lamellar, resinous yellow, with jamesonite.
  - I. Tripuhyite. 2FeO.Sb<sub>2</sub>O<sub>5</sub>. Micro-crystalline aggregates, dull greenish-yellow.
  - Range of Hardness 3-4 (Romeite 5.5)
    - I. Derbylite. 6FeO.5TiO<sub>2</sub>.Sb<sub>2</sub>O<sub>5</sub>(?). Orthorhombic, slender prisms.
    - I. Lewisite. 5CaO.2TiO<sub>2</sub>.3Sb<sub>2</sub>O<sub>5</sub>. Isometric, minute octahedrons.
    - I. Mauzeliite. 4(Ca,Pb)O.TiO<sub>2</sub>.2Sb<sub>2</sub>O<sub>5</sub>. Isometric, octahedrons, dark brown.
  - 671. Romeite. Perhaps CaSb<sub>2</sub>O<sub>4</sub>. Tetragonal, minute octahedrons, yellow.
- 2369\*672. Nadorite.  $PbSb_2O_4.PbCl_2$ . Orthorhombic, very thin tabular ||a|, yellow and brown. 1.50
- 2370°673. Ecdemite. Perhaps Pb<sub>4</sub>As<sub>2</sub>O<sub>7</sub>.2PbCl<sub>2</sub>. Tetragonal(?), incrustation, foliated, pearly yellow. 1.50
  - 674. Ochrolite. Pb<sub>4</sub>Sb<sub>2</sub>O<sub>7</sub>.2PbCl<sub>2</sub>(?). Orthorhombic, small, adamantine sulphur-yellow.
  - 675. Trippkeite. Essentially (nCuO, As<sub>2</sub>O<sub>3</sub>)(?). Tetragonal, small brilliant octahedrons, bluish-green.

II. Ceraleite, Coeruleite. CuO.2Al<sub>2</sub>O<sub>3</sub>. As<sub>2</sub>O<sub>3</sub>. Massive, clay-like, turquois-blue.

#### Antimonates or Antimonites of Doubtful Character

Barcenite. Chiefly Sb and Hg. Massive, gray-black. 2.50 Coronguite. Pb, Ag antimonate(?). Amorphous.

### Phosphates or Arsenates with Carbonates, Sulphates, Borates. Range of Hardness 2.5—5

2372 676. Dahllite. 2Ca<sub>3</sub>P<sub>2</sub>O<sub>8</sub>.CaCO<sub>3</sub>. ½ H<sub>2</sub>O. Fibrous crusts, resinous yellowish-white. 2.50

Ciplyte.  $4CaO.2P_2O_5.SiO_2(?)$ .

- II. Podolite. 3Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>.CaCO<sub>3</sub>. Hexagonal, microscopic crystals, yellow.
- 677. Diadochite. Perhaps 2Fe<sub>2</sub>O<sub>3</sub>.2SO<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.12H<sub>2</sub>O. Monoclinic, microscopic six-sided tables.
- 2373° globular, yellowish-brown. 1.00
- 2374 Destinezite, earthy nodular, yellowish. 1.00
- 2375°678. Pitticite. Hyd. Fe<sup>III</sup> arsenate and sulphate(?). Massive, whitish. 2.00
- 2376°679. Svanbergite. Chiefly hyd. Al and Ca phosphate and sulphate. Rhombohedral, small cuboids, red. 3.00
- 2377 I. Lossenite. 2PbSO<sub>4</sub>.3(FeOH)<sub>3</sub>As<sub>2</sub>O<sub>8</sub>+12H<sub>2</sub>O. Orthorhombic, acute pyramids, brownish-red. 1.50
  - II. Harttite. (Sr,Ca)O.2Al<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.SO<sub>3</sub>.5H<sub>2</sub>O. Hexagonal, pebbles, flesh-red.
- 2378 680. Beudantite. Fe<sup>III</sup> and Pb phosphate or arsenate with sulphate. Rhombohedral, small bright rhombs. 2.50
  - 681. Lindackerite. 3NiO.6CuO.SO<sub>3</sub>.2As<sub>2</sub>O<sub>5</sub>.7H<sub>2</sub>O. Orthorhombic, oblong rhombic tables, vitreous green.
- 2379 682. Lüneburgite. 3MgO.B<sub>2</sub>O<sub>3</sub>.P<sub>2</sub>O<sub>5</sub>.8H<sub>2</sub>O. Flat masses. 2.00

#### Nitrates. Hardness 2

- 2380+683. Soda Niter, Chile Saltpeter. NaNO<sub>3</sub>. Rhombohedral, crystalline mass, translucent white. .40
- 2381°684. Niter, Saltpeter. KNO<sub>3</sub>. Orthorhombic, white crust. .60
  - 685. Nitrocalcite.  $Ca(NO_3)_2 + nH_2O$ . Silky tufts, grayish-white.


686. Nitromagnesite.  $Mg(NO_3)_2 + nH_2O$ . Efflorescences, white.

687. Nitrobarite. Ba(NO<sub>3</sub>)<sub>2</sub>. Isometric, tetartohedral, plus and minus tetrahedrons forming octahedron, colorless.

- 688. Gerhardtite. 4CuO.N<sub>2</sub>O<sub>5</sub>.3H<sub>2</sub>O. Orthorhombic, vitreous deep emerald-green.
- 689. I. Darapskite. NaNO<sub>3</sub>.Na<sub>2</sub>SO<sub>4</sub>+H<sub>2</sub>O. Monoclinic, tabular || a, colorless.
- 690. Nitroglauberite. 6NaNO<sub>3</sub>.2Na<sub>2</sub>SO<sub>4</sub>.3H<sub>2</sub>O. Fibrous crystalline, white.

#### 5. Borates. Range of Hardness 6—8 (Ludwigite 5, Warwickite, Szaibelyite, Howlite 3.5)

- 691. Nordenskiöldine. CaO.SnO<sub>2</sub>. B<sub>2</sub>O<sub>3</sub>. Rhombohedral, tabular, yellow.
  - II. Hulsite (=Pageite?). 10(Fe<sup>11</sup>, Mg)O.2Fe<sub>2</sub>O<sub>3</sub>.SnO<sub>2</sub>.3B<sub>2</sub>O<sub>3</sub>. 2H<sub>2</sub>O(?). Orthorhombic(?), blackish.
- 692. Jeremejevite. Al<sub>2</sub>O<sub>3</sub>. B<sub>2</sub>O<sub>3</sub>. Hexagonal, long prisms, clear.
- 2382 693. Sussexite. 2(Mn,Zn,Mg)O.B<sub>2</sub>O<sub>3</sub>.H<sub>2</sub>O. Orthorhombic(?), fibrous, silky whitish. 3.00
- 2383 694. Ludwigite. Perhaps 3MgO.B<sub>2</sub>O<sub>3</sub> + FeO.Fe<sub>2</sub>O<sub>3</sub>. Orthorhombic, reniform, concentric radio-fibrous, black. 3.00
- 2384\* finely fibrous mass, silky black.
- 2385°695. Pinakiolite. 3MgO.B<sub>2</sub>O<sub>3</sub>+MnO.Mn<sub>2</sub>O<sub>3</sub>. Orthorhombic, thin prisms, tabular ||b|, brilliant black. 1.25
  - 4BeO.B<sub>2</sub>O<sub>3</sub>.H<sub>2</sub>O. Orthorhombic, prisms, 696. Hambergite. vitreous whitish.
- Minute indistinct 2386°697. Szaibelyite.  $5MgO.2B_2O_3.1\frac{1}{2}H_2O.$ needles, whitish. 1.50
- 2387\*698. Boracite, Stassfurtite. 6MgO.Mg Cl<sub>2</sub>.8B<sub>2</sub>O<sub>3</sub>. Isometric tetrahedral externally, orthorhombic and pseudo-isometric in molecular structure. Cube a truncated by dodecahedron d and tetrahedrons (fig.), small, ideal symmetry, bright, translucent pale gray. .50



2387. Boracite

2388

ditto, octahedron (tetrahedrons o and  $o_1$ ) prominent. .75

Boracite—Continued Type Species No. No. ditto, d prominent, pale green, loose. 2389 tetrahedron, truncated by cube a, 23900 minute, ideal symmetry, adamantine, clear pale green, loose (3). .50 massive, white. .40 2391+ **699.** Rhodizite.  $R_2O.2Al_2O_3.3B_2O_3$ , with R= K,Rb,Cs(?). Isometric, tetra-2395. Colemanite hedral, dodecahedrons, vitreous white. 2392°700. Warwickite. Perhaps 6MgO.FeO.2TiO2.3B2O3. rhombic, small slender prisms in limestone, dull black. .50 4CaO.5B<sub>2</sub>O<sub>3</sub>.2SiO<sub>2</sub>.5H<sub>2</sub>O. 2393°701. Howlite. Orthorhombic(?), crystalline nodules, embedded, white. -Range of Hardness 2.5-4 (Ulexite 1) 702. Lagonite. Fe<sub>2</sub>O<sub>3</sub>.3B<sub>2</sub>O<sub>3</sub>.3H<sub>2</sub>O. Earthy, yellow. 2394 703. Larderellite.  $(NH_4)_2O.4B_2O_3.4H_2O.$ clinic, very light mass of microscopic tables, whitish. 2.00 2CaO.3B<sub>2</sub>O<sub>3</sub>.5H<sub>2</sub>O. 2395°704. Colemanite. Monoclinic, highly complex, unit prism m prominent (fig.), perfect, adamantine, transparent pale yellow. 1.50 2397. Colemanite ditto, pyramids prominent, colorless. 1.50 23960 acute pseudo-rhombic, prism m, ortho-2397 dome W rounded (fig.), sharply defined. cleavage, brilliant, white. .50 2398+ Priceite. 5CaO.6B<sub>2</sub>O<sub>3</sub>.9H<sub>2</sub>O. Massive, friable chalky, 2399 snow-white. .75 Pandermite, compáct, porcelain-like. 2400° 2401°705. Pinnoite. MgO.B<sub>2</sub>O<sub>3</sub>.3H<sub>2</sub>O. Tetragonal, pyramidal hemihedrism, minute prisms, vitreous pale yellow. 2.00 Kaliborite. Hyd. Mg, K borate. Massive, resembling pinnoite. 706. Heintzite.  $K_2Mg_4B_{18}O_{32}$ . I 6  $H_2O(?)$ . 2402. Borax

Monoclinic, clear whitish.

190 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species No. No.

I. Ascharite. 3Mg<sub>2</sub>B<sub>2</sub>O<sub>5.2</sub>H<sub>2</sub>O. Amorphous, white.

2402+707. Borax. Na<sub>2</sub>O.2B<sub>2</sub>O<sub>3</sub>.10H<sub>2</sub>O. Monoclinic, unit prism m, ortho- and clinopinacoids a and b, pyramids z and o, base c (fig.), ideal symmetry, white, loose (3). .40

2403\*708. Ulexite. Na<sub>2</sub>O.2CaO.5B<sub>2</sub>O<sub>3</sub>.16H<sub>2</sub>O(?). Very light loose mass of capillary crystals, white. .50

Franklandite. Na<sub>2</sub>CaB<sub>6</sub>O<sub>11</sub>.7½H<sub>2</sub>O. Fine fibrous, white. Cryptomorphite. Hyd. Ca, Na borate. Kernels of microscopic rhombic plates, white.

709. Bechilite. CaO.2B<sub>2</sub>O<sub>3</sub>.4H<sub>2</sub>O. Crusts, white.

710. Hydroboracite. CaO.MgO.3B<sub>2</sub>O<sub>3</sub>.6H<sub>2</sub>O. Monoclinic(?), lamellar-fibrous, white, spotted red with iron oxide.

I. Sulphoborite. 4MgHBO<sub>3</sub>.2MgSO<sub>4</sub>.7H<sub>2</sub>O. Orthorhombic, small prisms, colorless.

#### Uranates. Hardness 5.5, 3 and 2.3

- 711. II. Uraninite. Uranate of uranyl, Pb, usually Th (or Zr), often the La and Y metals and N with He. Highly radio-active. Isometric.
  - 1. Crystallized varieties, black:
  - (a) Uranniobite, chiefly UO2, less UO3, octahedrons.
- 2404+ (b) Bröggerite, O ratio of UO<sub>3</sub>: other bases=1:1, cubooctahedrons, loose. 3.00
- 2405° (c) Cleveite, much UO<sub>3</sub>, with 10 p.c. Y earths, cubooctahedron modified by dodecahedron. 3.00
  - (d) Nivenite, much UO<sub>3</sub>, with 10 p.c. Y earths, massive.
- 2406+ 2. Massive, Pitchblende, no Th, N or rare earths, pitchblack. 3.00
  - II. Rutherfordine, alteration-product of uraninite.
- 2407+ II. Thorianite. Chiefly ThO<sub>2</sub>, U<sub>3</sub>O<sub>8</sub>. Highly radio-active. Isometric, ideal cubes, brilliant black. 2.50
- 2408<sup>®</sup> penetration-twins, fluor type. 2.50
- 2409°712. Gummite. (PbCa) U<sub>3</sub>SiO<sub>12</sub>.6H<sub>2</sub>O(?). Highly radio-active. Alteration-product of uraninite. Nodules, resinous reddish-yellow. 2.00

Yttrogummite. Y and U oxides, hydrous.


Thorogummite. UO<sub>3</sub>.3ThO<sub>2</sub>.3SiO<sub>2</sub>.6H<sub>2</sub>O. Highly radioactive. Tetragonal, prisms, dull yellowish-brown, loose. 2.00

- I. Mackintoshite. UO<sub>2</sub>.3ThO<sub>2</sub>.3SiO<sub>2</sub>.3H<sub>2</sub>O. Tetragonal, black.
- 713. Uranosphærite. Bi<sub>2</sub>O<sub>3</sub>.2UO<sub>3</sub>.3H<sub>2</sub>O. Hemispheres of minute acute crystals, reddish-yellow.

### 6. Sulphates, Chromates, Tellurates.


#### A. Anhydrous Sulphates, Etc. Range of Hardness 2-3

24110714. Mascagnite. (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>. Orthorhombic, mealy, yellowish. 1.50



715. Taylorite. 5K2SO4. (NH4)2SO4. Crystalline concretions, yellowish-white.

24120716. Thenardite. Na<sub>2</sub>SO<sub>4</sub>. Orthorhombic, prism m with macrodome t and base c rounded in combination, very large thick tabular || c (similar to fig.), eroded, translucent vellowish. 1.00



ditto, large, very thin. (3) .50 2413

cruciform-twin, tw.pl. e (fig.). 2414+

2415°717. Aphthitalite. (K,Na)<sub>2</sub>SO<sub>4</sub>. Rhombohedral, very thin hexagonal tables, white. 2.00




II. Palmierite. 3 (K, Na) 2 SO 4.4 Pb SO 4(?). Hexagonal, microscopic plates, colorless.

2416\*718. Glauberite. Na<sub>2</sub>SO<sub>4</sub>. CaSO<sub>4</sub>. Monoclinic, tabular  $\parallel$  base c, symmetrical, loose. .75



prismatic by extenison of unit pyramid s, 2417 distinct, pale yellowish. 1.50

> I. Langbeinite. K<sub>2</sub>SO<sub>4</sub>.2MgSO<sub>4</sub>. Isometric-tetartohedral, highly modified.



II. Vanthoffite. 3Na<sub>2</sub>SO<sub>4</sub>. MgSO<sub>4</sub>. Color-2418 less. 2.50

Orthorhombic. Range of Hardness 2.5—3.5 Barite Group. Type Species 719. Barite, Barytes. BaSO<sub>4</sub>. Orthorhombic. 1. Ordinary varieties:-(a) Crystals, perfectly developed, clear, brilliant:unit prism m, tabular || base c (fig.), 2419\* 2425. Barite large gray. ditto, with macrodome d elongated, prismatic aspect 2420+ (fig.), yellowish. ditto, with macrodome d, brachydome o (fig.), very 24210 large, thick tabular || c, translucent, dull brownish, loose. ditto, thick tabular, colorless, parallel growth pro-24229 ducing serrate-edged group, very large, loose. ditto, very thin tabular, greenish-blue, with calcite. 1.00 24230 m, d, o, c with pyramid z and prismatic by elongation of 2424 brachypinacoid b. .50 m, d, c, with macropinacoid a, prismatic by elongation of 2425 brachydome o, (similar to fig.) blue, large, loose. .30 macrodome d, brachydome o, base c (similar to fig.), 2426+ tabular, colorless. 2427º II. prismatic || axis c, highly complex, small but perfectly defined, adamantine. acicular, reddish. .40 2428 2426. Barite 2429\* (b) crested aggregate, white. .40 (c) columnar. .40 2430 (d) globular, Bologna Stone, grayish. 2431 (e) lamellar, curved, white. 2432+ (f) granular, grayish. 2433° 2434 (g) compact, yellowish. .30 (h) earthy. 2435 (i) stalactitic, polished section, concentric bands. 1.50 2436° 2. fetid, coarse granular, grayish. 2437° 3. Allomorphite, rectangular cleavages (pseudomorphous after anhydrite?). 4. Celestobarite, with much SrSO<sub>4</sub>. 2438 altered to quartz. 1.00


No No.	
	Celestite. SrSO <sub>4</sub> . Orthorhombic.
	I. Ordinary:—
	(a) Crystals brilliant, perfectly
	developed:—
2439+	unit prism m, macrodome d, base
	c, prismatic by elongation of
	brachydome o (similar to
	fig.), subtransparent white. 1.00
2440	ditto, with pyramid y. 1.00
2441*	d, c with m prominent, trans-
	lucent bluish75
2442	tabular    c, clear colorless.
	.50
2443°	rough flat prisms, red50 2451. Anglesite
2444+	cleavage, translucent pale
	sky-blue20
2445° `	stalactitic, radio-columnar structure with drusy crys-
	talline surface, white50
2446*	(b) fibrous, blue50
2447	(c) lamellar, bluish-white75
24480	(d) granular, coarse, pale blue.
	.20
<b>244</b> 9°	(e) concretionary30
2450	(f) earthy30
	2. Calciocelestite, contains m a m
	3. Barytocelestite, contains
	much Ba.
2451+721.	Anglesite. PbSO <sub>4</sub> . Orthorhombic, 2454. Anglesite
	unit prism m, macropinacoid
	a, macrodome $d$ , pyramids and base $c$ , tabular
	(aspect like fig.), ideal symmetry, adamantine
	translucent gray, on galena. 1.50
2452°	prism m prominent, terminated by low brachydomes,
	limpid, small. 1.00
2453	pyramids predominating. 2.50
2454°	highly modified stout crystals (aspect like fig.), trans-
-707	lucent yellowish-white, perfect. 2.00
2455	pale green crystals. 2.50
2456	drusy crystals coating twinned cerussite. 1.50
~430	diady distrib conting thinned cornovice. Tijo

194 C Type Species	OMPLETE TYPE COLLECTION. DANA'S SY	YSTEM
No. No. 2457*	compact massive, concentrically	
<b>24</b> 3/*	banded, grayish. 1.50	7
2458*722.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2458. Anhydrite
	prismatic by elongation of	- <b>10</b>
	several macrodomes, brachydome	s (fig.), bright
		40
2459°	cleavage, rectangular, red75	1
2460	cleavage, rectangular, grayish75	/./.\
2461	fibrous. 1.00	("/")
2462+	fine granular, pale bluish20	M
	scaly granular, Vulpinite.	ارا ا
2463°	compact, banded vein in granular rock	salt,  m m
	grayish40	
	pseudomorphous, in cubes after rock sal	
723.	Zinkosite. ZnSO <sub>4</sub> . Orthorhombic. Need	ds 2465. Crocoite
	confirmation.	•
2464 <b>724</b> .	Hydrocyanite. CuSO <sub>4</sub> . Orthorhombic,	AA
	green, in lava. 5.00	
725.	I. II. Crocoite. PbCrO. Monoclinic,	
	crystals perfectly developed, highly	
	adamantine, translucent brilliant scarlet:—	m   m
2465*	short unit prism $m$ , prism $f$ , pryamid	
2400	t, base $c$ (similar to fig.), small, with	
	vauquelinite. 2.50	
24660	short unit prism m with unit pyramid	
	v (similar to fig.), small, on limonite.	
	1.00	2466. Crocoite
2467°	long unit prism m, clinodome z, loose,	
	large. 2.00	k. 277 2
2468+	ditto, with clinodome $w$ , pyramid $t$ ,	R
	orthodome $k$ , base $c$ (similar to	
	fig.). 2.00	
2469	ditto, with clinodome y and new	$\begin{bmatrix} m & m \end{bmatrix}$
	clinodome $j$ (fig.). 6.00	
2470°	acicular, on limonite. 1.00	
2471	long prism, not terminated, large,	
	loose25	
2472+	ditto, on limonite, large. 1.00	· ·
2473	ditto, hollow, loose. 1.00	2468. Crocoite

Type Species No. No. Crocoite—Continued

2474

- dull etched rounded crystals on white schist. 1.50
- 726. Phœnicochroite. 3PbO.2CrO<sub>3</sub>. Orthorhombic(?), red, yellow on exposure.
- 2475 727. Vauquelinite. Perhaps 2(Pb,Cu)CrO<sub>4</sub>. (Pb,Cu)<sub>3</sub>P<sub>2</sub>O<sub>6</sub>. Monoclinic, druse of microscopic crystals, dark greenish-brown. 5.00
  - Jossaite. Contains Cr<sub>2</sub>O<sub>3</sub>, PbO, ZnO. Orthorhombic, minute orangeyellow crystals on vauquelinite.



2469. Crocoite

- 2476° Tarapacaite. Chiefly K<sub>2</sub>CrO<sub>4</sub>. Minute fragments, canary-yellow, disseminated in soda niter. 1.00
- 2477 I. Euchlorine. Contains SO<sub>3</sub>,CuO,K<sub>2</sub>O,Na<sub>2</sub>O. Orthorhombic, incrustation on lava, emerald-green. 1.50
  - I. Dietzeite. 7Ca(IO<sub>3</sub>)<sub>2</sub>. 8CaCrO<sub>4</sub>. Monoclinic, dark gold-yellow.
  - II. Bellite. PbCrO<sub>4</sub> with As<sub>2</sub>O<sub>3</sub>. Hexagonal, minute tufted needles, bright crimson-red.

# Sulphates with Chlorides, Carbonates, Etc.—In Part Hydrous Compounds. Range of Hardness 2—4.5

- 728. Sulphohalite. 3Na<sub>2</sub>SO<sub>4</sub>.2NaCl. Isometric. transparent greenish-yellow.
- 729. Caracolite. Pb(OH)Cl.Na<sub>2</sub>SO<sub>4</sub>. Orthorhombic(?), pseudohexagonal twins, incrustation.
  - Chlorothionite. K<sub>2</sub>SO<sub>4</sub>.CuCl<sub>2</sub>. Crystalline crusts, bright blue, from lava.
  - II. Arzrunite. (Pb<sub>2</sub>O)SO<sub>4.3</sub>(CuCl<sub>2</sub>.H<sub>2</sub>O).Cu(OH)<sub>2</sub>(?). Orthorhombic, small prisms, bluish-green.
- 730. Kainite. MgSO<sub>4</sub>.KCl+3H<sub>2</sub>O. Monoclinic, tabular || c. 2478° granular massive. .40
- 2479 731. Connellite. Cu<sub>15</sub>(Cl,OH)<sub>4</sub>SO<sub>16</sub>.15H<sub>2</sub>O(?). Hexagonal, small prisms, translucent blue.
  - 732. Spangolite. (AlCl)SO<sub>4</sub>.6Cu(OH)<sub>2</sub>+3 H<sub>2</sub>O. Rhombohedral, hexagonal tables, dark green.



196 COMPLETE TYPE COLLECTION. DANA'S SYSTEM Type Species
2480*733. Hanksite. 4Na <sub>2</sub> SO <sub>4</sub> . Na <sub>2</sub> CO <sub>3</sub> . Hexagonal, short unit prism
m, unit pyramid $o$ , base $c$ prominent (fig.), ideal
symmetry, translucent yellowish-white, loose50
2481 ditto, with prism also prominent. 1.00
2482° ditto, with pyramid alone prominent. 1.00
2483 ditto, with pyramid s, tabular $   c$ 75
2484*734. Leadhillite. 4PbO.SO <sub>3</sub> .2CO <sub>2</sub> .H <sub>2</sub> O(?). Monoclinic, pseudo-
hexagonal twins, tw.pl. prism m, tabular, pearly
straw-yellow. 2.50
2485 ditto, translucent apple-green. 4.00
2486 cleavage. 1.00
Susannite. 4PbO.SO <sub>3</sub> .2CO <sub>2</sub> .H <sub>2</sub> O(?). Formerly regarded as
rhombohedral but very probably monoclinic and
therefore leadhillite, acute rhombic aspect. 8.00
I. Beresowite. 6PbO.3CrO <sub>3</sub> . CO <sub>2</sub> .Crystalline lamellar, red.
1. Deresowite. Of bo. 30103. Co2. Crystamine famenar, red.
D. Asid and Davis Culmbates. Donney of Handana and a
B. Acid and Basic Sulphates. Range of Hardness 2.5—3.5
735. Misenite. K <sub>2</sub> SO <sub>4</sub> . H <sub>2</sub> SO <sub>4</sub> . Fibers, silky-white.
736. Alumian. Al <sub>2</sub> O <sub>3</sub> .2SO <sub>3</sub> (?). Rhombohedral(?), white.
II. Doughtiyite. Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> .5Al <sub>2</sub> (OH) <sub>6</sub> .21H <sub>2</sub> O. Powder, white.
2488°737. Lanarkite. PbSO <sub>4</sub> .PbO. Monoclinic, slender prismatic by
extension of orthopinacoid a, adamantine, trans-
lucent straw-yellow. 5.00
2489 738. Dolerophanite. 2CuO.SO <sub>3</sub> (?). Monoclinic, brown. 4.00
2490°739. Caledonite. 2(Pb,Cu)O.SO <sub>3</sub> .H <sub>2</sub> O(?). Orthorhombic,
microscopic, prismatic    axis a, translucent bluish-
green. 3.00
2491+740. Brochantite. 4CuO.SO <sub>3</sub> .3H <sub>2</sub> O. Orthorhombic, unit prism
m and domes prominent, ideal symmetry, adaman-
tine, translucent dark emerald-green, small. 1.25
2492° acicular, clear emerald-green. 1.25
2493 drusy incrustation, emerald-green. 1.00
2494 massive, dull. 1.00
II. Steltznerite. CuSO <sub>4</sub> .2Cu(OH) <sub>2</sub> . Orthorhombic, green.
Waringtonite, doubly curving wedge-shaped crystals,
pale green.

2497°741. Linarite. PbO.CuO.SO<sub>3</sub>.H<sub>2</sub>O. Monoclinic, flat prismatic || axis b, adamantine, translucent deep azure-blue, small, perfect. 2.00

2498 minute acicular. 2.00

Antlerite. 10CuO.3SO3.7H2O(?). Massive, green.

#### C. Hydrous Sulphates.—Normal Division

Hardness 2 (Kieserite 3, Szmikite 1.5)

742. Lecontite. (Na,NH<sub>4</sub>,K)<sub>2</sub>SO<sub>4</sub>+2H<sub>2</sub>O. Orthorhombic, prisms.


Guanovulite. 7K<sub>2</sub>O.2(NH<sub>4</sub>)<sub>2</sub>O.12SO<sub>3</sub>.11H<sub>2</sub>O. Crystalline, silky yellowish-white, organic origin.

2499\*743. Mirabilite, Glauber Salt. Na<sub>2</sub>SO<sub>4</sub>+10H<sub>2</sub>O. Monoclinic, efflorescent crust, white. .75

Exanthalose. Na<sub>2</sub>SO<sub>4</sub>. + 2H<sub>2</sub>O. Efflorescence, white.

- 2500\*744. Kieserite. MgSO<sub>4</sub>+H<sub>2</sub>O. Monoclinic, granular massive, whitish. .40
  - I. Cubeite (Kubeite). Contains SO<sub>3</sub>, Fe<sub>2</sub>O<sub>3</sub>, MgO,
     H<sub>2</sub>O. Rhombic or monoclinic pyramids.
  - II. Ferrofallidite. FeO.SO<sub>3</sub>+H<sub>2</sub>O. Clear grains.
  - 745. Szmikite. MnSO<sub>4</sub>+ H<sub>2</sub>O. Amorphous, reddishwhite.
  - 746. Gypsum. CaSO<sub>4</sub>+2H<sub>2</sub>O. Monoclinic.
    - Selenite, large crystals of ideal symmetry, transparent colorless:—
- unit prism m, unit pyramid l, clinopinacoid
  b (fig.), phosphoresces green in ultra- 2501. Gypsum
  violet light, loose. .20
- ditto, with "phantom" lines of growth. .30
- 2503 ditto, very large, loose (not phosphorescent). 1.00
- 2504 ditto, with rough orthodome e (fig.), phosphorescent, loose. .20
- 2505° ditto, very large (not phosphorescent). .75
- 2506° wedge-shaped, very large, enclosing sulphur, etc. .50
- 2507\* lenticular, dull yellowish in clay. .30
- 2508° ditto, rosette-like group, red. .50
- 2509 acicular, small, on lava. .75
- 2510 bent crystal, very large. .50





Type Species	COMPLETE TYPE COLLECTION. DANA'S SYSTEM  Gypeum—Continued
2511*	contact-twins, tw.pl. a (fig.), "swallow-tail twins," yellowish, loose20
2512	ditto, aggregate75
2513	cruciform-penetration-twins, tw.pl. a75
2514+	cleavage    pinacoid b perfect,    a imperfect conchoidal,    n imperfect fibrous, clear, rhomboidal outline20
2515	arenaceous, containing sand50
25169	containing moving liquid. 1.25
2517+	2. fibrous, fine, Satin spar, white40
2518	fibrous, fine, Satin spar, flesh-red60
2519°	fibrous, coarse, white30
2520°	fibrous, plumose, lamellar-stellate60
25219	fibrous, curving flower-like forms75
2522+	3. massive, Alabaster, very fine grained, white20
2523*	massive, scaly-granular, reddish20
2524	massive, earthy, "rock-gypsum." .30
25250	altered to quartz50
747.	Hesite. RSO. + 4H.O. with R=Mn.Zn.Fe. Monoclinic

- 747. Resite. RSO<sub>4</sub>+4H<sub>2</sub>O, with R=Mn,Zn,Fe. Monoclinic (?), prismatic, loose aggregates.
- II. Scleropasthite. Hyd. Fe<sup>II</sup>, Crsulphate. Felted mass, white. 2526+748. Epsomite, Epsom Salt. MgSO<sub>4</sub>+7H<sub>2</sub>O. Orthorhombic, fibrous, white. .75

Tauriscite. FeSO<sub>4</sub>+7H<sub>2</sub>O(?). Orthorhombic, acicular, greenish.

- 2527 749. Goslarite, Zinc Vitriol. ZnSO<sub>4</sub>+7H<sub>2</sub>O. Orthorhombic, acicular. 2.00
- 2528\* massive, white. 1.00
  - II. Cuprogoslarite, contains Cu.
  - 750. Morenosite. NiSO<sub>4</sub>+7H<sub>2</sub>O. Orthorhombic, acicular, greenish.
    - II. Boothite. CuO.SO<sub>3</sub>.7H<sub>2</sub>O. Monoclinic, blue. Fauserite. (MnMg)SO<sub>4</sub>+6H<sub>2</sub>O(?). Orthorhombic, white.

#### Melanterite or Copperas Group

Monoclinic. Soft

These species are the ordinary vitriols, being identical in general formula with the members of the Epsomite group, and regarded as essentially the same compound under oblique crystallization.

Type Species No. No.

- 2529+751. Melanterite, Copperas or Green Vitriol. FeSO<sub>4</sub>+7H<sub>2</sub>O.

  Monoclinic, fibrous. .75
- 2530 pulverulent coating. .40
  - 752. Mallardite. MnSO<sub>4</sub>+7H<sub>2</sub>O. Monoclinic, fibrous crystalline masses, colorless.
- 2531 753. Pisanite. (Fe,Cu)SO<sub>4</sub>+7H<sub>2</sub>O. Monoclinic, concretionary, bright blue. 1.25
  - I. Salvadorite. (Cu,Fe)SO<sub>4</sub>+7H<sub>2</sub>O. Monoclinic, aggregates of rough prisms, bluish-green.
  - 754. Bieberite. CoSO<sub>4</sub>+7H<sub>2</sub>O. Monoclinic, crusts, red. Cupromagnesite. (CuMg)SO<sub>4</sub>+7H<sub>2</sub>O. Monoclinic, crusts on lava, bluish-green.
- 2532°755. Chalcanthite, Blue Vitriol. CuSO<sub>4</sub>+5H<sub>2</sub>O. Triclinic, flattened || p. 2.00
- 2533 fibrous, translucent. 1.50
- 2534+ massive, fine prussian-blue. .50
  - I. Siderotil. FeSO<sub>4</sub>+5H<sub>2</sub>O. Divergent needles.
- 2535 756. Syngenite. CaSO<sub>4</sub>·K<sub>2</sub>SO<sub>4</sub>+H<sub>2</sub>O. Monoclinic, prisms flattened || a, clear colorless. 3.00
  - 757. Löweite. MgSO<sub>4</sub>. Na<sub>2</sub>SO<sub>4</sub> + 2 ½ H<sub>2</sub>O. Tetragonal, cleavable, whitish.
- 2536°758. Blödite. MgSO<sub>4</sub>. Na<sub>2</sub>SO<sub>4</sub>+4H<sub>2</sub>O. Monoclinic, highly modified short prism, large, clear colorless. 1.50
- 2537 ditto, small, with kröhnkite. .75
- 2538\* massive. .50
  - I. Leonite. K<sub>2</sub>SO<sub>4</sub>.MgSO<sub>4</sub>+4H<sub>2</sub>O. Monoclinic, tabular.
  - 759. Boussingaultite.  $(NH_4)_2SO_4$ .  $MgSO_4 + 6H_2O$ . Monoclinic, prismatic with c prominent.
  - 760. Picromerite. MgSO<sub>4</sub>.K<sub>2</sub>SO<sub>4</sub>+6H<sub>2</sub>O. Monoclinic, crystalline incrustation, white.
  - 761. Cyanochroite. CuSO<sub>4</sub>. K<sub>2</sub>SO<sub>4</sub>+6H<sub>2</sub>O. Monoclinic, crystalline crust, clear blue.
- Hardness 4.5 and 3
  2539\* II. Natrochalcite. Na<sub>2</sub>SO<sub>4</sub>.Cu<sub>4</sub>(OH)<sub>2</sub>(SO<sub>4</sub>)<sub>2</sub>+2H<sub>2</sub>O. Monoclinic, sharply developed acute pyramids (fig.), brilliant and translucent, fine emerald-green. 3.00
- 2540 762. Polyhalite. 2CaSO<sub>4</sub>.MgSO<sub>4</sub>.K<sub>2</sub>SO<sub>4</sub>+2H<sub>2</sub>O. Monoclinic(?), cleavage, red. .75

200 COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type Species Polyhalite—Continued
No. No.

2541° fibrous, translucent. .75

2542+ granular-cleavable, reddish. .30

2543 Krugite. 4CaSO<sub>4</sub>.MgSO<sub>4</sub>.K<sub>2</sub>SO<sub>4</sub>+2H<sub>2</sub>O. Crystalline massive, gray. .50

Mamanite. Like polyhalite but CaO: MgO: K<sub>2</sub>O=3:2:1. Fibro-foliated, silky white.

763. Wattevillite. CaSO<sub>4</sub>.Na<sub>2</sub>SO<sub>4</sub>+4H<sub>2</sub>O(?).
Orthorhombic or monoclinic, microscopic needles, silky snow-white.

#### Alum Group

Isometric. Range of Hardness 1-3

Hydrous sulphates of aluminium with an alkali metal and 24 molecules of water.

2544°764. Kalinite.  $K_2SO_4$ .Al<sub>2</sub>( $SO_4$ )<sub>3</sub>+24H<sub>2</sub>O. Isometric, crusts, white. .75

765. Tschermigite.  $(NH_4)_2SO_4.Al_2(SO_4)_3 + 24$  $H_2O$ . Octahedrons.

2545° fibrous, subtransparent white. 1.25

2546 766. Mendozite. Na<sub>2</sub>SO<sub>4</sub>.Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> + 24H<sub>2</sub>O. <sup>2539. Natrochalcite</sup> Fibrous mass, white. 4.00

I. Kauaiite. Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> 7·18, Al<sub>2</sub>O<sub>3</sub> 33·40, K<sub>2</sub>SO<sub>4</sub> 17·00, Na<sub>2</sub> SO<sub>4</sub> 4·91, H<sub>2</sub>O 31·57. Chalk-like.

2547°767. Tamarugite.  $Na_2SO_4.Al_2(SO_4)_3 + 12H_2O$ . Massive, white. 1.00

#### Halotrichite Group. Monoclinic. Soft

Hydrous sulphates of aluminium with magnesium, manganese, etc. and 22 to 24 molecules of water.

2548°768. Pickeringite. MgSO<sub>4</sub>.Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>+22H<sub>2</sub>O. Monoclinic(?), silky fibrous mass, white. 1.00

Seelandite, MgAl<sub>2</sub>(SO<sub>4</sub>)<sub>4</sub>+27H<sub>2</sub>O.
 Stüvenite. (Na<sub>2</sub>Mg)SO<sub>4</sub>Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>+24H<sub>2</sub>O(?). Needles.

2549 Picroallumogene. 2MgSO<sub>4</sub>.Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>+28H<sub>2</sub>O(?). Massive, whitish. 1.25

Type Species No. No.

Sonomaite. 3MgSO<sub>4</sub>.Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>+33 H<sub>2</sub>O. Crystalline, silky colorless.

Dumreicherite. 4MgSO<sub>4</sub>.Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>+ 36H<sub>2</sub>O. Monoclinic(?), columnar crusts.

Aromite. 6MgSO<sub>4</sub>.Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>.54H<sub>2</sub>O. Crystalline.



2550\*769. Halotrichite. FeSO<sub>4</sub>.Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>+24H<sub>2</sub>O. Monoclinic or triclinic, silky fibers, white. 1.50


2551


pulverulent incrustation. .75

770. Apjohnite. MnSO<sub>4</sub>. Al<sub>2</sub>
(SO<sub>4</sub>)<sub>3</sub>+24 H<sub>2</sub>O.
Monoclinic (?), fibrous mass, whitish.

Bushmanite, (Mn.Mg) $SO_4.Al_2(SO_4)_3 + 22$  $(or 24)H_2O.$ 


771. Dietrichite.  $(Z_n, F_e, M_n)$   $SO_4.Al_2(SO_4)_3 + 22$   $H_2O$ . Monoclinic (?), silky fibers.





2552 772. Coquimbite. Fe<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>+
9H<sub>2</sub>O Rhombohedral, octahedroid,
rhombohedron o
and base c predominating, truncated by diagonal
prism a and rhombohedron r (fig.),
distinct. 2.50

2553° granular massive. 1.00 773. Quenstedtite. Fe<sub>2</sub>O<sub>3</sub>.3 SO<sub>3</sub>.10H<sub>2</sub>O. Monoclinic, transparent reddish-violet.



202 ( Type Species No. No.	COMPLETE TYPE COLLECTION. DANA'S SYSTEM
	Ihlëite. Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> +12H <sub>2</sub> O. Efflor-
	escence, yellow.
	Kornelite. $Fe_2(SO_4)_3 + 7\frac{1}{2}H_2O$ .
2554+775	Alunogen. $Al_2(SO_4)_3 + 18H_2O$ .
-554	Managlinia for moder
	forming surface of silky fi-
	- 1 1 / / 1
	brous mass, yellowish-white.
	.75
2555*776	II. Kröhnkite. CuSO <sub>4</sub> .Na <sub>2</sub> SO <sub>4</sub> +2
2555****	H <sub>2</sub> O. Monoclinic, octahe- 2558. Kröhnkite
	1120. Monochime, octane-
	droid, unit prism m, unit
	pyramid $q$ (fig.), ideal symmetry, bluish-green. 1.50
2556°	slender prisms $m$ , with pyramids $q$ and $s$ prominent
	(similar to figs. a and b), ideal symmetry, ada-
	mantine, fine clear blue. 2.00
2557	ditto, but short prisms forming solid crusts. 3.00
2558	contact-twins, tw.pl. base c, imperfect (fig.), fine clear
	blue. 2.00
2559°	penetration-twins, tw.pl. base c, ideal symmetry with
007	aspect of parallel growth, bluish-green. 1.50
	Phillipite. $CuSO_4$ . $Fe_2(SO_4)_3 + nH_2O_1$ , approximately. Mass-
	ive, azure-blue.
777	Ferronatrite. 3Na <sub>2</sub> O.Fe <sub>2</sub> O <sub>3</sub> .6SO <sub>3</sub> .6H <sub>2</sub> O. Rhombohedral,
,,,,	spherical, lamellar-stellate, whitish.
770	•
778.	Römerite. Perhaps FeSO <sub>4</sub> .Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> +12H <sub>2</sub> O. Triclinic,

# C. Hydrous Sulphates.—Basic Division Hardness 2.5

2561 779. Langite. 4CuO.SO<sub>3</sub>.4H<sub>2</sub>O. Orthorhombic, minute twins, tw.pl. prism m, pseudo-hexagonal.

2562° concretionary crust, green. 1.50

tabular || c.

2560°

granular massive, brown. 1.25

2563°780. Herrengrundite. CaO.4CuO.2SO<sub>3</sub>.6H<sub>2</sub>O. Monoclinic, small scale-like hexagons in spherical groups, pearly, bright emerald-green. 1.25

780A. Arnimite. 5CuO.2SO<sub>3</sub>.6H<sub>2</sub>O. Crystalline incrustation, bright green.

- Type Species
  No. No.
  I. Kamarezite. (CuOH)<sub>2</sub>SO<sub>4</sub>.Cu(OH)<sub>2</sub>.6H<sub>2</sub>O. Orthorhom-
- 2564 781. Cyanotrichite, Lettsomite. 4CuO.Al<sub>2</sub>O<sub>3</sub>.SO<sub>3</sub>.8H<sub>2</sub>O. Orthorhombic, capillary, fine blue. 3.00
- 2565°782. Serpierite. Basic Cu and Zn sulphate. Orthorhombic, tufts of microscopic crystals, tabular ||c|, blue. 2.00
  - -Range of Hardness 1.5—3 (Carphosiderite 4)
  - 783. Castanite. Fe<sub>2</sub>O<sub>3</sub>.2SO<sub>3</sub>.8H<sub>2</sub>O. Monoclinic, minute prisms, brilliant, chestnut-brown.

Rubrite. Fe<sub>2</sub>O<sub>3</sub>.2SO<sub>3</sub>.3H<sub>2</sub>O.

- 784. Copiapite. Perhaps 2Fe<sub>2</sub>O<sub>3</sub>.5SO<sub>3</sub>.18H<sub>2</sub>O. Monoclinic. tabular  $\parallel b$ .
- granular massive, brownish-vellow. 1.00 2566+
  - 785. Knoxvillite. Hyd. Cr, Fe<sup>III</sup> and Al sulphate. rhombic(?), rhombic plates, greenish-yellow.
    - Redingtonite. Hyd. Cr sulphate. Finely fibrous mass silky pale purple.
- 2567 786. Utahite. 3Fe<sub>2</sub>O<sub>3</sub>.3SO<sub>3</sub>.4H<sub>2</sub>O. Rhombohedral, microscopic hexagonal scale-like tables, silky orangeyellow. 1.50
- 787. Amarantite. Fe<sub>2</sub>O<sub>3</sub>.2SO<sub>3</sub>.7H<sub>2</sub>O. Triclinic, slender prisms. columnar mass, red. 2.00 2568°
- 2569°788. Fibroferrite. Fe<sub>2</sub>O<sub>3</sub>.2SO<sub>3</sub>.10H<sub>2</sub>O. Monoclinic(?), radiofibrous silky mass, pearly pale yellow. 1.50
  - 789. Raimondite. 2Fe<sub>2</sub>O<sub>3</sub>.3SO<sub>3</sub>.7H<sub>2</sub>O. Hexagonal or rhombohedral, scale-like hexagons, pearly yellow.
- Apatelite. 4Fe<sub>2</sub>O<sub>3</sub>.6SO<sub>3</sub>.3H<sub>2</sub>O. Nodular, clear yellow. .75 2570 2571°790. Carphosiderite. Perhaps 3Fe,O<sub>3</sub>.4SO<sub>3</sub>.10H<sub>2</sub>O. Rhombohedral(?). Submicaceous mass, straw-yellow. 1.50
- 2572\*791. Aluminite. Al<sub>2</sub>O<sub>3</sub>.SO<sub>3</sub>.9H<sub>2</sub>O. Monoclinic, chalky reniform nodules. .50

oölitic earthy, whitish.

Werthemanite. Al<sub>2</sub>O<sub>3</sub>.SO<sub>3</sub>.3H<sub>2</sub>O. Massive, white.

Winebergite. Al Basic sulphate.

- I. Planoferrite. Fe<sub>2</sub>O<sub>3</sub>.SO<sub>3</sub>.15H<sub>2</sub>O. Orthorhombic(?), tabular, greenish.
- 792. Glockerite. 2Fe<sub>2</sub>O<sub>3</sub>.SO<sub>3</sub>.6H<sub>2</sub>O. Massive.
- 793. Felsöbanyite. 2Al<sub>2</sub>O<sub>3</sub>.SO<sub>3</sub>.10H<sub>2</sub>O. Orthorhombic, mass of minute hexagonal scales, pearly white.
- 794. Paraluminite. 2Al<sub>2</sub>O<sub>3</sub>.SO<sub>3</sub>.15H<sub>2</sub>O(?). Massive, whitish.

- 795. Cyprusite. Perhaps 7Fe<sub>2</sub>O<sub>3</sub>.Al<sub>2</sub>O<sub>3</sub>.10SO<sub>3</sub>.14H<sub>2</sub>O. Hexagonal(?), chalky mass of microscopic hexagons, yellow.
  - Range of Hardness 2.5—3.5
- 796. Voltaite. (Fe<sup>II</sup>,Mg)<sub>5</sub>(Fe<sup>III</sup>,Al)<sub>4</sub>S<sub>10</sub>O<sub>41</sub>.15H<sub>2</sub>O(?). Isometric (?), cubo-octahedrons, resinous dark-greenish.
- 797. Metavoltine. Perhaps 5(K<sub>2</sub>,Na<sub>2</sub>,Fe)O.3Fe<sub>2</sub>O<sub>3</sub>.12SO<sub>3</sub>.18 H<sub>2</sub>O. Hexagonal, mass of minute scales, yellow.
- 2573°798. Botryogen. Perhaps MgO.FeO.Fe<sub>2</sub>O<sub>3</sub>.4SO<sub>3</sub>.18H<sub>2</sub>O. Monoclinic, small prisms, deep red, 1.00
  - II. Palacheite, 2MgO.Fe<sub>2</sub>O<sub>3</sub>.4SO<sub>3</sub>.15H<sub>2</sub>O, prisms.
  - I. Idrizite. (Mg,Fe) (Fe,Al)<sub>2</sub>S<sub>3</sub>O<sub>13</sub> + 16H<sub>2</sub>O. Crystalline, yellowish-gray.
- 2574°799. Sideronatrite. 2Na<sub>2</sub>O.Fe<sub>2</sub>O<sub>3</sub>.4SO<sub>3</sub>.7H<sub>2</sub>O. Orthorhombic, fibro-crystalline mass, yellow. 2.00
- 2575°800. Alunite. K<sub>2</sub>O.3Al<sub>2</sub>O<sub>3</sub>.4SO<sub>3</sub>.6H<sub>2</sub>O. Rhombohedral, minute cuboid rhombs, brightly defined. .50
- 2576+ granular massive, white. .30
- 2577 compact massive. .30
- 801. Jarosite. K<sub>2</sub>O.<sub>3</sub>Fe<sub>2</sub>O<sub>3</sub>.<sub>4</sub>SO<sub>3</sub>.<sub>6</sub>H<sub>2</sub>O. Rhombohedral.
- 2578+ I. Crystallized, minute cuboid rhombs, sharp and symmetrical, brown. 1.00
- 2579° tabular || c, yellow-brown, small, sharply defined. 1.00 2580 modified, translucent brown. 1.50
- 2. Concretionary, tuberose incrustation. 3.00
  - II. Natrojarosite, Na replaces K.
    - II. Plumbojarosite, Pb replaces K.
      - Decomposition products of pyrite:—
      - Plagiocitrite. (K,Na)<sub>2</sub>O.2FeO.3(Al,Fe)<sub>2</sub>O<sub>3</sub>.6SO<sub>3</sub>.27H<sub>2</sub>O(?). Monoclinic or triclinic, microscopic prisms, yellow.
      - Clinophæite. 4(K,Na)<sub>2</sub>O.FeO.(Fe,Al)<sub>2</sub>O<sub>3</sub>.5SO<sub>3</sub>.8H<sub>2</sub>O(?).
  - Monoclinic(?), microscopic crystals, blackish-green.

    802. Löwigite. Perhaps K<sub>2</sub>O.3Al<sub>2</sub>O<sub>3</sub>.4SO<sub>3</sub>.9H<sub>2</sub>O. Rounded masses, pale straw-yellow.
  - 803. I. Ettringite. Perhaps 10CaO.2Al<sub>2</sub>O<sub>3</sub>.5SO<sub>3</sub>.54H<sub>2</sub>O. Hexagonal, minute needles, clear colorless.
  - 804. Quetenite. MgO.Fe<sub>2</sub>O<sub>3</sub>.3SO<sub>3</sub>.13H<sub>2</sub>O. Monoclinic or triclinic(?), indistinct prisms, reddish-brown.
  - 805. Zincaluminite. 6ZnO.3Al<sub>2</sub>O<sub>3</sub>.2SO<sub>3</sub>.18H<sub>2</sub>O. Hexagonal (?), minute hexagonal scales, bluish-white.

Type Species No. No.

- Lamprophanite. Hyd. Pb, Mn, Ca, Mg, Na, K sulphate. Cleavable folia, pearly white.
- 806. Johannite. Hyd. U,Cu sulphate. Monoclinic, masses of microscopic tables, fine emerald-green.
- 807. Uranopilite. Perhaps CaO.8UO<sub>3</sub>.2SO<sub>3</sub>.25H<sub>2</sub>O. Incrustation of minute needles, yellow.

Tellurates; Also Tellurites, Selenites. Soft

- 808. Montanite. Bi<sub>2</sub>O<sub>3</sub>.TeO<sub>3</sub>.2H<sub>2</sub>O. Incrusting, earthy, whitish.


  Hardness 5 and 2
- 809. Emmonsite. Hyd. Fe<sup>III</sup> tellurite(?). Monoclinic(?), thin cleavage scales, clear yellowish-green.
- 810. Durdenite. Fe<sub>2</sub>O<sub>3</sub>.<sub>3</sub>TeO<sub>2</sub>.<sub>4</sub>H<sub>2</sub>O. Massive, small mammillary forms, greenish-yellow.
  - Magnolite. Hg<sub>2</sub>TeO<sub>4</sub>(?). Microscopic needles, silky white.

    Hardness 3
- 811. Chalcomenite. CuO.SeO<sub>2</sub>.2H<sub>2</sub>O. Monoclinic, small short prisms, bright blue.
  - Molybdomenite. Pb selenite(?). Orthorhombic, very thin scales, pearly white.
  - Kerstenite. Chiefly SeO<sub>2</sub>,PbO. Botryoidal, sulphur-yellow.
- 7. Tungstates, Molybdates. Hardness 5 (Raspite 2.5)
  - 812. Wolframite. (Fe, Mn) WO4. Monoclinic.
- 2582° I. I. Normal Wolframite, Fe WO<sub>4</sub>, square prismatic || axis b, orthopinacoid a and base c both prominent, highly modified (?), small, brilliantly distinct, black. 1.00
- 2583 ditto, long lenticular. 1.00
- 2. Ordinary, ratio Fe: Mn=9: I
  to 2: 3, tabular || orthopinacoid
  a, unit prism m, orthodome t,
  clinodome f all prominent (similar to fig.), large, sharply defined, brilliant black, loose. 1.00

2585+ bladed basal cleavage. .75

2586 lamellar massive. 1.00

2587° granular massive. .75



Type Species No. No.	LETE TIPE COLLECTION. DANAS SISTEM
No. No. 2588 <b>0813. Hü</b> l	onerite. MnWO4. Monoclinic, distinct bladed crys
	tals, dark reddish-brown. 2.00
2589+ di	tto, large, embedded in quartz. 1.00
2590° I. Ras	pite. PbWO4. Monoclinic, minute tables, adamantin
	clear brownish-yellow. 6.00
Scheelite	Group. Tetragonal. Range of Hardness 3-4.5
2591 <b>9814.</b> Sch	eelite. CaWO4. Tetragonal, pyramidal hemihedrisn
	octahedroid, unit pyramid p prominent, smal
	perfect, brownish. 1.00
2592 00	tahedroid, diametral pyramid e, ideal symmetry
	translucent straw-yellow. 3.00
2593* ui	nit pyramid p, truncated by diametral
	pyramid e (similar to fig.), small, brightly defined. 1.00
2594° ol	otuse diametral ovramid a prominent / 1/1/
<b>2</b> 394° OL	small but distinct, grayish. 1.25
2595° ta	hular II hase c dull hounded by pyra- \ \ \ \
	mids e and p adamantine, minute,
	perfect75
2596 di	tto, globular groups, greenish75 2593. Scheelite
	assive cleavable-granular, whitish75
	tered to wolframite. 2.00
815. Cup	rotungstite. CuWO4, also (Ca,Cu)WO4. Crystalline
	granular, glassy green.
816. Pow	ellite: CaMoO <sub>4</sub> . Tetragonal, minute modified octa
	hedroids, subtransparent, resinous greenish-yellow
2599 <b>0817</b> . Stol	zite. PbWO4. Tetragonal, pyramidal hemihedrism
	very acute pyramid and base, nearly opaque
	resinous brownish, small. 3.00
2600 ur	iit pyramids $n$ , $v$ , unit prism $m$ , base $c$ , ideal sym
	metry, small, red. 6.00
2601* th	ick tabular    base c, unit pyramid n, diametral pyra
	mid e, minute, ideal symmetry, adamantine, clea
C 040 TIT 1	reddish-brown. 3.00
2002 818. Wuli	fenite. PbMoO4. Tetragonal, pyramidal hemihedrism
	minute ideal octahedroid, unit pyramid e, red. 1.2
2603 ta	bular    base c with unit pyramid u, ideal symmetry
	adamantine, translucent fine orange-red. 1.50

SCHEELITE GROUP. IODATES. OXALATES AND MELLATES 207

Type Species No. No.
2604° ditto, with diametral pyramid s
(similar to fig.). 1.50

2605+ ditto, less symmetrical, paler red,
cavernous aggregate of brilliant crystals. 1.00

2606\* prism m rounded, with base c, small, ideal symmetry, resinous yellowish-brown. 1.00

ditto, thick tabular, resinous brownish, large. 1.50 ditto, very thin tabular, clear lemon-yellow. 2.00

2609 819. Reinite. FeWO<sub>4</sub>. Tetragonal, pyramidal, blackish-brown. 8.00

Achrematite. 3[3Pb<sub>3</sub>As<sub>2</sub>O<sub>8</sub>.PbCl<sub>2</sub>].4[Pb<sub>2</sub>MoO<sub>5</sub>]. Massive, cryptocrystalline, yellowish-red.

820. Belonesite. MgMoO<sub>4</sub>(?). Tetragonal, minute needles, clear white.

#### VIII. Indates

S.,I. Lautarite. Ca(IO<sub>3</sub>)<sub>2</sub>. Monoclinic, radiately arranged prisms.

# VII. Salts of Organic Acids

Oxalates, Mellates, Etc. Range of Hardness 2-2.5

- 2610 821. Whewellite. CaC<sub>2</sub>O<sub>4</sub>+H<sub>2</sub>O. Monoclinic, twins, tw.pl. e, small heart-shaped, sharp and perfect, glassy colorless. 9.00
  - 822. Oxammite.  $(NH_4)_2C_2O_4 + 2H_2O$ . Orthorhombic, prismatic, silky clear whitish.
- 2611 823. Humboldtine. 2FeC<sub>2</sub>O<sub>4</sub>+3H<sub>2</sub>O. Plates, yellowish. 4.00 2612\*824. Mellite. Al<sub>2</sub>C<sub>12</sub>O<sub>12</sub>+18H<sub>2</sub>O. Tetragonal, obtuse unit pyramid o, sharply defined, resinous translucent honeyyellow. 1.00
- 2613° Pigotite.  $4Al_2O_3.C_{12}H_{10}O_8 + 27H_2O$ . Massive, brown. 1.50
  - II. Moissanite. CSi. Hexagonal plates in meteoric iron.

# VIII. Hydrocarbon Compounds

The hydrocarbon compounds in general, with perhaps a few exceptions, are not homogeneous substances, but mixtures, which by the action of solvents or by fractional distillation may be separated into two or more component parts. Hence, they are not definite mineral species. Those printed in capitals are indicated by Dana as leading compounds. A very large number of names of related but less important substances are here omitted.

## Simple Hydrocarbons

Chiefly Members of the Paraffin Series.  $CnH_{2n+2}$ . Type No.

SCHEERERITE. C 73 p.c., H 24 p.c. A polymer of marsh gas(?). Monoclinic, thin tables, translucent.

2614 HATCHETTITE. C 85.55 p.c., H 14.45 p.c. Massive, translucent yellowish. .50

2615\* OZOCERITE. C 84.43 p.c., H 13.69 p.c. Foliated wax, dark brownish. .20

2616 FICHTELITE. C<sub>15</sub>H<sub>26—28</sub>(?). Monoclinic, translucent white.. 75

2617 HARTITE. Ratio of C to H=12:20. Triclinic or monoclinic, translucent white. .75

KÖNLITE. Ratio of C to H=1: 1.  $n(C_6H_6)$ . A polymer of benzene. Amorphous, brownish.

### 2 Oxygenated Hydrocarbons

Comprise chiefly the numerous kinds of native fossil resins often designated by the generic term, amber.

2618+ Succinite, True Amber. Ratio of C,H,O=40:64:4.

Irregular mass, translucent yellow, clouded. .50

2619 transparent, precious. .50

2620° ditto, containing insects. .75

RETINITE. A generic name including Copalite and many other amber-like resins. They contain little or no succinic acid.

2621° Copalite. Ratio for C,H,O=40:64:1. Amorphous, resinous clear pale yellow. .40

Type No.

ditto, containing insects. .60

2623° I. Allingite. A fossil resin, in shale. .50

BATHVILLITE. Ratio of C,H,O=40:68:4. Amorphous, like completely decayed wood, dull fawn-brown.

2624° TASMANITE. Ratio of C,H,O,S=40:62:2:1. Minute scales in shale, resinous brown. .40

Dysodile. C 69 p.c., H 10 p.c., O 16.9 p.c., S 2.35 p.c., N 1.7 p.c. Thin folia, yellowish.

Pyroretinite. Ratio of C, H, O=40:56:4. Resin-like. Leucopetrite. C<sub>50</sub>H<sub>84</sub>O<sub>3</sub>. Between resin and wax in characters.

GEOMYRICITE. C<sub>34</sub>H<sub>68</sub>O<sub>2</sub>. Wax-like.

GEOCERITE. C<sub>28</sub>H<sub>56</sub>O<sub>2</sub>. Wax-like, white.

BOMBICCITE. C<sub>7</sub>HO<sub>13</sub>. Triclinic, clear colorless.

2625° IDRIALITE. C<sub>80</sub>H<sub>56</sub>O<sub>2</sub>. Massive, white, mixed with cinnabar, clay, pyrite, gypsum and a solid brownish-black earth. 1.00

ROCHLEDERITE. Ratio of C,H,O=40:56:6. Resin-like, transulcent reddish-brown.

DOPPLERITE. C<sub>12</sub>H<sub>14</sub>O<sub>6</sub>. Amorphous, jelly-like, brownish.

### Appendix to Hydrocarbons

In general the following more complex substances are less definite than those described in the preceding groups.

2626+ Petroleum. Chiefly consists of members of the paraffin and asphaltum series, C , H<sub>2 m+2</sub>, varying from Marsh Gas, CH<sub>4</sub>, to the solid forms. Oily liquid. .20

2627+ ASPHALTUM, Mineral Pitch. A mixture of different hydrocarbons, part of which are oxygenated. Amorphous, blackish, solid. .20

2628 ditto, viscous. .20

2629<sup>®</sup> ELATERITE, Elastic Bitumen. Approximately C 85 p.c., H 12—13 p.c. Massive, soft, very elastic, dark brown. .40

26300 Wurtzilite. Compact, brilliant black. .20

2631<sup>o</sup> Uintahite, Gilsonite. An asphaltum. Compact, brilliant black. .20

210	COMPLETE TYPE COLLECTION. DANA'S SYSTEM
Type No.	
140.	MINERAL COAL. Mainly oxygenated hydrocarbons of
	vegetable origin. Massive:—
2632+	1. Anthracite, Hard Coal, 85-93 p.c. C, compact,
	brilliant black20
2633	ditto, in limestone20
	2. Bituminous, Soft Coal, 5—15 p.c. O:—
2634°	(a) Caking or coking coal, fragile, greasy pitch-black.
	.20
2635*	(b) Non-caking coal, fragile, greasy pitch-black20
26369	ditto, breaking in layers, iridescent20
2637*	(c) Cannel coal, compact, dull grayish-black20
26380	(d) Brown coal, Lignite, friable, dark brownish20
2639	Brown coal, Jet, compact, brilliant black, polished40
26400	Peat, partly carbonized vegetable fibres, loosely matted
•	mass30

# Supplement

#### Minerals Measured but not Analyzed

The angles of the following very rare minerals have been measured and the forms so determined, indicate that future chemical analyses may prove them to be distinct species.

- S. Hessenbergite. A silicate. Monoclinic, tabular || c, adamantine, clear colorless with bluish tinge.
- S. Mursinskite. Tetragonal, clear yellow.

# PART III

Index to

Complete Type Collection, Dana's System

Price List

of Hand Size Specimens

## Index

#### Complete Type Collection—Dana's System

# Price List

#### Hand Size Specimens

ABBREVIATIONS. The species numbers preceding the names are those in Dana's "System of Mineralogy," 6th Ed. "r" or "n" follows the name of a mineral, it is related to or near the species, the number of which precedes the indexed name; when followed by "s," it is a synonym; followed by "s. v.," "s.n." or "s. r." it is a synonym of a variety of the species, or a synonym of a mineral near or related to it; if followed by "ap." it will be found in the first appendix following the species number given. "H" designates Hydrocarbons, described at the end of the "System."

Approximate Prices are quoted on good typical specimens of minerals ordinarily in stock. A range of price indicates different types or varieties. Hence selections are best made from the preceding systematic list, giving composition, crystallization, structure and color, with separate price for each type.

HAND SIZE SPECIMENS, averaging 10 x 7 cm. (4 x 23/4 in.),

furnished at approximately the *list prices*.

Museum Size Specimens, averaging 12x9 cm. (43/4 x 3 ½ in.), sold at double the list prices.

Printed Labels attached give name, composition and locality.

PASTEBOARD TRAYS are included (or blocks with museum

specimens if requested).

Free Transportation to any address in the world. Any

or all specimens may be returned at our expense.

OUR APPROVAL SYSTEM with its risk of double transportation cost assumed by us, gives assurance that purchasers will be pleased.

A DEDUCTION of 10 per cent. is made on \$20.00 worth of

hand or museum specimens without chest, if all are kept.

DETAILED INFORMATION as to sizes, labels, trays, transportation, terms, chests, cabinets, etc. is given in Part I.

OTHER PRICE LISTS. Besides the preceding complete descriptive list of over 2600 minerals, price lists will be found in: Part IV., 180 of the Common Minerals arranged according to metallic constituents; Part V., 400 Economic Minerals, similarly arranged; Part VI., 300 crystals, classified under their system of crystallization; Part VIII., Laboratory List (alphabetical) of pure minerals sold by weight for chemical purposes.

## Index and Price List

447. Acadialite\$ .50—\$ .75	483. Alipite, A. r
57. Acanthite 2.00	325. Alkali-augite\$ .75
819. Achrematite, r	426. Alkali Tourmaline
426. Achroite	578. Allactite 2.00
366. Achtaragdite, r	409. Allanite
326. Acmite	9. Allemontite 3.00
338. Actinolite	H. Allingite, r
210. Actinolite in Quartz 1.00	370. Allochroite, s.v
210. Actinolitic Quartz	102. Alloclasite 2.50
563. Adamite 1.00— 2.00	719. Allomorphite
556. Adelite, n	24. Allopalladium
524. Adelpholite, r	498. Allophane
313. Adularia	544. Alluaudite, r
326. Æegirite 1.∞	370. Almandite30— 2.00
343. Ænigmatite 2.50	509. Aloisiite, r
532. Æschynite 1.50— 2.50	510. Alshedite
458. Agalmatolite, r 1.00	278. Alstonite, s 2.00
270. Agaric mineral	46. Altaite 2.00
210. Agate	Alums, 764-767
210. Agate-Jasper 1.00	736. Alumian
210. Agatized Wood 1.00	791. Aluminite
327. Aglaite, s.r	370. Aluminium Garnet30— 3.00
434. Agnolite, n	212. Alumocalcite
373. Agricolite 5.00	800. Alunite
47. Aguilarite, n 6.00	775. Alunogen
138. Aikinite 3.00	462. Alurgite, B., r
248. Ainalite, r 4.∞	394. Alvite, r 1.50
63. Alabandite 1.00- 3.00	17. Amalgam 3.00— 9.00
746. Alabaster	13. Amalgam, Gold, r
325. Alalite 2.00	787. Amarantite 2.00
118. Alaskaite, s 3.00	315. Amazonite, s.v50— 2.00
435. Albine	315. Amazonstone50— 2.00
316. Albite	H. Amber, Succinite5075
210. Albite in Quartz 1.00	H. Amber, a generic term
242. Alexandrite 2.50	559. Amblygonite
389. Algerite, n	324. Amblystegite
38. Algodonite 3.00	470. Amesite, r
30	414

210.	Amethyst\$ .50—\$3.00	789. Apatelite, r.,\$ .75
338.	Amianthus, s.v	549. Apatite
338.	Amphibole20— 1.50	270. Aphrite
	Amphodelite	477. Aphrosiderite
450.	Analcite	717. Aphthitalite 2.00
	Anapāite, n	770. Apjohnite
	Anatase, s	370. Aplome 1.00
	Ancylite, r	435. Apophyllite
	Andalusite	344. Aquamarine 1.25— 5.00
	Andesine	223. Aqueous Vapor
	Andesite, s	277. Aragonite
•	Andesite, a rock	418. Ardennite
114.	Andorite, n 4.00— 7.00	210. Arenaceous Quartz
	Andradite40— 3.00	342. Arfvedsonite 1.25
	Andrewsite, r	78. Argentiferous Bornite
	Anglesite 1.00— 2.50	45. Argentiferous Galena
	Anhydrite20— 1.00	118. Argentiferous Galeno-
	Ankerite, A30— 1.00	bismutite 3.00
	Annabergite 1.00	87. Argentiferous Smaltite 2.00
	Annerödite 4.00	148. Argentiferous Tetra-
	Annivite	hedrite, s 1.00
	Anomalite 1.50	270. Argentine
	Anomite, see note	42. Argentite 1.25— 2.50
	Anorthite	56. Argentopyrite, r
315.	Anorthoclase, A75— 1.00	232. Argillaceous Hematite, s.v20
	Anthophyllite40— .60	163. Argyrodite 2.50— 9.00
	Anthosiderite, r	253. Arkansite
	Anthracite	780. Arnimite, A
	Anthracite in Quartz50	768. Aromite, r
	Anthraconite,	17. Arquerite 3.00— 9.00
	Antigorite	Arsenates, etc., 536-690
<b>4</b>	Antimonates, etc., 669-675	8. Arsenic
	ANTIMONIDES, ETC., 35-108	9. Arsenic, Antimonial, r
	Antimonites, etc., 669-675	87. Arsenical Cobalt, s 1.00— 2.50
٥.	Antimonial Arsenic, r.	71. Arsenical Nickel, s . 1.00— 3.00
-	Antimonial Native Silver 3.00	98. Arsenical Pyrites, s25— 1.25
•	Antimonial Niccolite 1.00	145. Arsenical Red Sil-
	Antimonial Red Sil-	ver, s 1.25— 3.00
	ver, s 1.00— 2.50	Arsenides, etc., 35-108
TO.	Antimony 2.50— 3.00	582. Arseniopleite 2.00
	Antimony Glance, s35— 2.00	577. Arseniosiderite 1.50
	Antimony Ocher, s	Arsenites, etc., 669-675
	Antlerite, r	8. Arsenolamprite, r
	Antozonite	213. Arsenolite 3.00
-12.		

98.	Arsenopyrite\$ .25— \$1.25	336. Babingtonite \$5.00
4.	Arsenschwefel, r	458. Baddeckite, r
3.	Arsensulfurite, r	254. Baddeleyite, n 4.00
	Artinite, n	99. Badenite, r
729.	Arzrunite, r	409. Bagrationite
338.	Asbeferrite	325. Baikalite
338.	Asbestus	401. Bakerite, n
	Asbestus, s.v	234. Balas Ruby
	Asbestus in Quartz 1.00	399. Bamlite
	Asbolite, r	210. Banded Agate
	Ascharite, n	675. Barcenite, n 2.50
549.	Asparagus-stone 1.50	270. Bardiglio Marble30
	Aspasiolite, r	270. Baricalcite
H.	Asphaltum	719. Barite
	Asteriated Quartz, s.v 1.00	342. Barkevikite, A
231.	Asteriated Sapphire, s.v.40— 1.50	610. Barrandite
325.	Asteroite	320. Barsowite, r
338.	Astochite 1.25	430. Barylite, ap
435.	Astrolite, r	354. Barysilite 1.50
514.	Astrophyllite	462. Barythiotite 1.25
	Atacamite 1.00— 2.50	719. Barytes, s
193.	Atelite, r	282. Barytocalcite 1.00— 1.50
584.	Atelestite 2.00	720. Barytocelestite
	Atheriastite, n	210. Basanite
669.	Atopite	232. Basanomelan 1.50
645.	Attacolite, r	579. Basiliite, r
394.	Auerbachite, r	324. Bastite, r
395.	Auerlite, r	285. Bastnäsite 4.00
645.	Augelite, r	462. Bastonite, r
325.	Augite	487. Batavite, r
	Auralite, r	H. Bathvillite
	Aurichalcite	374. Batrachite 1.50
85.	Auriferous Pyrite	122. Baumhauerite, n 4.00
	Auripigment, s., 1.00— 3.00	261. Bauxite
	Automolite 1.50— 2.50	444. Bavenite, r 4.∞
	Autunite 1.25— 2.50	629. Bayldonite 3.∞
	Avalite 1.50	394. Beccarite
	Aventurine	709. Bechilite
	Aventurine4075	425. Beckelite, n
	Aventurine Quartz 1.00— 1.25	155. Beegerite
	Awaruite 3.00	210. Beekite
	Axinite	727. Bellite, r
289.	Azurite	820. Belonesite
210.	Babel-quartz	507. Bementite 2.00

514. Benitoite, n\$7.00	210. Black Tourmaline in
648. Beraunite 1.00— 1.50	Quartz\$ .50—\$2.00
734. Beresowite, r	58. Blende, s
338. Bergamaskite	758. Blödite
479. Berlauite, r	535. Blomstrandine, n
645. Berlinite, r	535. Blomstrandite, ap
473. Berthierine, s.r	210. Blood-stone50
119. Berthierite 1.00	344. Blue Aquamarine 2.00
422. Bertrandite 2.00	755. Blue Vitriol, s50— 2.00
344. Beryl	599. Bobierrite
546. Beryllonite 1.00— 4.00	409. Bodenite, r
49. Berzelianite 1.50	269. Bog Manganese, r20— .40
538. Berzeliite 2.00	259. Bog Ore
680. Beudantite 2.50	493. Bole
407. Beustite 1.50	192. Boléite, r
76. Beyrichite	108. Bolivianite, ap
426. Bi-colored Tourma-	719. Bologna Stone, s.v
line 1.50— 2.00	375. Boltonite50
754. Bieberite	H. Bombiccite
497. Biharite, r	353. Bonsdorfite, r
670. Bindheimite 1.∞	750. Boothite, n
149. Binnite, formerly 123 3.00	698. Boracite
462. Biotite	BORATES, 691-713
270. Bird's-eye Marble	707. Borax
197. Bischofite	653. Borickite
217. Bismite 1.50— 2.00	78. Bornite
11. Bismuth 1.00— 5.00	1. Bort
13. Bismuth-gold	798. Botryogen 1.00
29. Bismuthinite 1.00— 3.00	401. Botryolite
29. Bismuth Glance, s 1.00— 3.00	139. Boulangerite75— 1.00
306. Bismutite 1.00	136. Bournonite 1.25— 2.00
95. Bismutosmaltite	759. Boussingaultite
283. Bismutosphärite	481. Bowenite
H. Bitumen, Elastic, s.	566. Brackebuschite 2.50
Elaterite	465. Brandisite 1.25
H. Bituminous Coal20— .40	591. Brandtite 1.50
509. Bityite, r	247. Braunite 1.00— 1.50
247. Bixbyite, n 2.00	85. Bravoite
338. Black Hornblende20— 1.50	426. Brazilian Emerald50— 2.00
210. Black Hornblende in	426. Brazilian Peridot, s.v50— 2.00
Quartz	426. Brazilian Sapphire 3.00
58. Black Jack, s. v	254. Brazilite, s 4.00
2. Black Lead, s30— .75	270. Breccia Marble30
426. Black Tourmaline20— 1.00	270. Brecciated Onyx 1.00
,	

338. Breislakite\$1.00	395. Calciothorite, r
72. Breithauptite 1.50— 2.50	565. Calciovolborthite\$4.00
272. Breunnerite 1.00	270. Calcite
439. Brewsterite 2.00	370. Calcium-iron Garnet .50— 3.00
537. Britholite, n	270. Calc Spar, s20— 2.50
153. Brittle Silver, s 2.00— 3.00	270. Calc Spar, s20— 2.50 270. Calc Tufa20— .60
	370. Calc I tha 1.00
740. Brochantite 1.00— 1.50	739. Caldente 3.00
711. Bröggerite 3.00	,
Bromides, etc., 164-209	612. Callainite
278. Bromlite 2.00	164. Calomel 4.00
171. Bromyrite 5.00	551. Campylite 2.00
132. Brongniardite	325. Canaanite
323. Bronzite	360. Cancrinite
253. Brookite	162. Canfieldite, n
269. Brostenite, r 2.50	H. Cannel Coal
259. Brown Clay-iron-	175. Capped Fluor 1.50
stone	210. Capped Quartz 1.25
H. Brown Coal	347. Cappelenite
259. Brown Iron-ore, s20— 1.00	729. Caracolite
426. Brown Tourmaline40— 2.00	1. Carbonado 5.00
262. Brucite 1.00— 2.50	Carbonates, 270-309
471. Brunsvigite, r	541. Carminite
618. Brushite	201. Carnallite
407. Bucklandite	210. Carnelian
210. Buhrstone	666. Carnotite, n 1.00— 4.00
227. Bunsenite	498. Carolathine, r 1.00
770. Bushmanite	424. Carpholite 1.00
335. Bustamite 2.00	790. Carphosiderite 1.50
338. Byssolite	270. Carrara Marble
603. Cabrerite 4.00	82. Carrollite
212. Cacholong 1.00	540. Caryinite 2.00
392. Cacoclasite, r	349. Caryocerite 5.00
647. Cacoxenite 1.00	508. Caryopilite 1.00
58. Cadmiferous Blende75	248. Cassiterite30 — 2.50
275. Cadmiferous Smithsonite 1.50	783. Castanite
344. Caesium Beryl 4.00	310. Castorite
210. Cairngorn Stone, s.v25— 2.00	342. Cataphorite, r
H. Caking (coking) Coal	346. Catapleiite 1.00
423. Calamine	458. Cataspilite, n
105. Calaverite, r 4.00	210. Cat's-Eye50
270. Calcareous Marl	242. Cat's-Eye 5.00
720. Calciocelestite	210. Cavernous Quartz50
652. Calcioferrite	489. Celadonite
280. Calciostrontianite	720. Celestite
	,

71Q.	Celestobarite\$1.00	111. Chiviatite
	Celsian, n	88. Chloanthite\$1.25—\$2.50
	Cement Rock, s.v20	179. Chloralluminite, r
	Cenosite 4.00	549. Chlor-apatite
	Ceraleite, r	457. Chlorastrolite, ap50— 1.00
	Cerargyrite 1.25— 3.00	CHLORIDES, ETC., 164-209
	Cerasite	469. Chlorite, s
	Cerite 1.50	210. Chloritic Quartz75— 1.00
	Cerolite, r 1.50	466. Chloritoid35— 1.00
	Cerussite 50— 2.00	176. Chloromagnesite
	Cervantite	167. Chloromanganokalite, r.
	Ceylonite40— 3.00	328. Chloromelanite
	Ceylon Peridot 3.00	505. Chloropal
	Chabazite	175. Chlorophane 1.50
	Chalcanthite50— 2.00	479. Chlorophæite, n
	Chalcedony40— 1.50	353. Chlorophyllite, r50
	Chalcocite 1.00— 3.00	234. Chlorospinel 2.00
474.	Chalcodite	729. Chlorothionite, r
	Chalcolamfrite, n 1.25	596. Chlorotile, r
811.	Chalcomenite	572. Chondrarsenite
435.	Chalcomorphite, n	415. Chondrodite40— 2.00
268.	Chalcophanite	58. Christophite
636.	Chalcophyllite 2.∞— 3.∞	CHROMATES, ETC., 714-811
83.	Chalcopyrite35— 2.∞	325. Chrome-diopside
	Chalcopyrrhotite, r	234. Chrome-spinel, s.v.
	Chalcosiderite 1.25	462. Chromglimmer
117.	Chalcostibite 9.00	241. Chromic Iron, s20— .50
224.	Chalcotrichite	241. Chromite
	Chalilite	370. Chromium Garnet 1.00— 3.00
	Chalk	241. Chrompicotite
	Chalk, French	242. Chrysoberyl 1.00— 5.00
	Chalmersite, n 6.∞	504. Chrysocolla50— 4.00
	Chalybite, s20— 3.00	376. Chrysolite30— 6.00
	Chalypite, r	210. Chrysoprase 1.00— 1.50
	Chamosite, r	481. Chrysotile50
	Chenevixite	606. Churchite
	Chert, s.v	495. Cimolite
	Chessylite, s75— 4.00	66. Cinnabar
	Chesterlite	370. Cinnamon-stone
	Chiastolite	676. Ciplyte, r
	Childrenite 1.50	270. Cipolin Marble
	Chilenite	576. Cirrolite
	Chile Saltpeter, s	210. Citrine, s.v
184.	Chiolite 2.00	158. Clarite, r

215. Claudetite\$6.00	784. Copiapite\$1.∞
47. Clausthalite 2.∞	15. Copper
232. Clay Iron-stone	751. Copperas, s
316. Cleavelandite	54. Copper Glance, s 1.00— 3.00
58. Cleiophane 1.∞	83. Copper Pyrites, s35— 2.00
711. Cleveite 3.∞	772. Coquimbite 1.00— 2.50
1. Cliftonite, r	353. Cordierite, s50— 2.00
468. Clinochlore50— 2.50	284. Cordylite, n 5.00
571. Clinoclasite 2.50	634. Cornwallite
423. Clinohedrite, n 9.∞	675. Coronguite, n
416. Clinohumite 4.∞	470. Corundophilite75— 2.00
801. Clinophæite, r	231. Corundum20— 4.00
465. Clintonite	91. Corynite 9.00
210. Clouded Agate	128. Cosalite 2.00
270. Clouded Onyx	467. Cosmochlore, r
H. Coal, Anthracite	459. Cossaite
H. Coal, Bituminous20— .40	343. Cossyrite
H. Coal, Brown20— .40	180. Cotunnite 1.50
H. Coal, Caking (coking)20	67. Covellite 1.00— 3.00
H. Coal, Cannel	245. Crednerite
H. Coal, Mineral20— .40	233. Crichtonite
H. Coal, Non-Caking	211. Cristobalite, r
601. Cobalt Bloom, s.v 1.00	341. Crocidolite
89. Cobalt Glance, s	725. Crocoite
89. Cobaltite	472. Cronstedtite 2.00
325. Coccolite	53. Crookesite 7.00
96. Cockscomb Pyrites	339. Crossite, r
675. Coeruleite, s.r	183. Cryolite
645. Cœruleolactite, r	183. Cryolithionite, r
25. Cohenite, r	461. Cryophyllite
704. Colemanite50— 4.00	553. Cryphiolite, r
586. Collophanite	185. Cryptohalite, r
499. Collyrite	708. Cryptomorphite, r
370. Colophonite	81. Cubanite 6.00
COLUMBATES, ETC., 520-535	744. Cubeïte, r 1.∞
525. Columbite 1.00— 2.00	
212. Common Opal40— 1.50	338. Cummingtonite
628. Conichalcite 1.00	549. Cupro-apatite
486. Connarite	112. Cuprobismutite
731. Connellite 4.00	564. Cuprodescloizite 1.00
460. Cookeite, r50	749. Cuprogoslarite
H. Copalite, a kind of	173. Cuproiodargyrite, r
retinite	754. Cupromagnesite, r
	104. out.oumBurgeo, r

45.	Cuproplumbite, r \$2.50	677. Diadochite \$1.00
815.	Cuprotungstite	325. Diallage
413.	Cuspidine 6.00	274. Dialogite, s75— 4.00
400.	Cyanite	1. Diamond 1.00— 6.00
<b>761.</b>	Cyanochroite	134. Diaphorite 2.50
	Cyanotrichite 3.00	256. Diaspore 1.50— 3.00
320.	Cyclopite 1.00	212. Diatomaceous Earth, s.v30
162.	Cylindrite, n 1.25	588. Dickinsonite
	Cymatolite, r	771. Dietrichite
	Cyprine 1.00	727. Dietzeite, n
	Cyprusite	569. Dihydrite 3.00
	Cyrtolite, r 1.00	325. Diopside
	Dahllite 2.50	383. Dioptase 1.00— 5.00
	Daleminzite, r	388. Dipyre
	Damourite	400. Disthene, s
98.	Danaite	585. Dittmarite, r
	Danalite 4.00	112. Dognacskaite, r
396.	Danburite	270. Dog-tooth Spar50
	Dannemorite	738. Dolerophanite 4.00
	Daphnite	271. Dolomite
	Darapskite	270. Dolomitic Calcite
	Dark Ruby Silver, s 1.00— 2.50	37. Domeykite 1.25— 2.50
	Datolite	H. Dopplerite
	Daubréeite	270. Doubly Refracting
	Daubreelite 8.00	Spar, s 1.00— 2.00
_	Davidite, r	736. Doughtiyite, r
-	Daviesite	200. Douglasite
	Davyne, r 2.50	426. Dravite
	Dawsonite 1.00	210. Drusy Quartz
	Dechenite, r	275. Dry-bone, s.v
	Delafossite, ap	480. Dudleyite, r
	Delessite	573. Dufrenite
	Delorenzite, n	127. Dufrenoysite 2.00
	Delvauxite, r	427. Dumortierite 1.00
	Demantoid 3.∞	768. Dumreicherite, r
	Dendritic Agate 1.∞	296. Dundasite, n
	Derbylite, n	558. Durangite 1.∞
	De Saulesite, r, A 2.00	810. Durdenite
	Descloizite 1.00— 1.50	141. Dürfeldtite, r
	Desmine, 840— 1.00	519. Dysanalyte
	Destinezite 1.00	35. Dyscrasite 2.50— 8.00
	Deweylite	236. Dysluite 2.00
	Diabantite	H. Dysodile
324.	Diaclasite, r	335. Dyssnite, r

458.	Dysyntribite, r\$ .40	402. Erdmannite, r
673.	Ecdemite 1.50	535. Erikite, n
338.	Edenite	568. Erinite\$2.00
452.	Edingtonite 4.00	437. Erionite, n
25.	Edmonsonite, r	78. Erubescite, s75— 2.50
188.	Egglestonite, n 8.00	601. Erythrite 1.00— 2.00
	Egyptian Jasper	193. Erythrocalcite, r 5.00
270.	Egyptian Marble, s.v	199. Erythrosiderite
570.	Ehlite, r 1.50	69. Erythrozincite, r
	Eisenbrucite, r	353. Esmarkite, r
232.	Eisenrosen 1.50	259. Esmeraldaite, r
479.	Ekmannite, n	370. Essonite, s.v
357.	Elæolite	803. Ettringite
H.	Elastic Bitumen, s	51. Eucairite 9.∞
	Elaterite	727. Euchlorine, r 1.50
H.	Elaterite	632. Euchroite 2.00
13.	Electrum 2.00	403. Euclase 3.00— 9.00
648.	Eleonorite 1.50	345. Eucolite 1.25
453.	Ellagite, r	510. Eucolite-titanite
183.	Elpasolite, r	395. Eucrasite, r
346.	Elpidite, n 2.50	358. Eucryptite
170.	Embolite 1.25— 3.00	345. Eudialyte 1.25
	Emerald	312. Eudidymite
	Emery	450. Eudnophite
809.	Emmonsite	462. Eukamptite, r
116.	Emplectite 1.25	368. Eulytite 2.00— 3.00
158.	Enargite 1.00— 2.00	459. Euphyllite, r 1.25
270.	Encrinal Marble	479. Euralite, n
535.	Endeiolite, n	564. Eusynchite, r
551.	Endlichite, r50— 1.50	450. Euthallite
	Enstatite30— 8.00	534. Euxenite 1.50
650.	Eosphorite 5.00	645. Evansite 1.00
	Epiboulangerite 2.00	743. Exanthalose, r
	Epichlorite, n	210. Eye-agate 1.50
	Epididymite, n 1.00	148. Fahlerz, s 1.00- 2.00
407.	Epidote40— 3.00	353. Fahlunite, r
	Epidote in Quartz	592. Fairfieldite
	Epigenite	141. Falkenhaynite, r
	Epigenite, r	159. Famatinite 2.50
	Epiphanite, n	453. Fargite
	Epistilbite 1.50	456. Faröelite, s.v 1.00
	Epistolite, n 2.00	325. Fassaite 1.00
	Epsomite	451. Faujasite 1.00
748.	Epsom Salt, s	750. Fauserite, r

377. Fayalite\$1.00—\$2.50	625. Forbesite\$4.00
130. Feather Ore, s 1.00	212. Forcherite 1.50
FELDSPAR GROUP, 313-320	443. Foresite, r 2.00
316. Feldspar, Soda, s20— 1.00	375. Forsterite50
793. Felsöbanyite	210. Fortification-Agate 1.25
523. Fergusonite 1.50	232. Fossil Ore, s.v
FERRATES, ETC., 234-247	407. Fouquéite
376. Ferrite, r	335. Fowlerite
270. Ferrocalcite	549. Francolite 1.50
89. Ferrocobaltite	162. Franckeite, n 1.50
744. Ferrofallidite, r	708. Franklandite, r
777. Ferronatrite	239. Franklinite
583. Ferrostibian, r	149. Fredricite
210. Ferruginous Quartz75— 1.50	148. Freibergite 1.00
719. Fetid Barite	135. Freieslebenite 2.50
270. Fetid Calcite	484. French Chalk
788. Fibroferrite 1.50	395. Freyalite, r
399. Fibrolite	384. Friedelite 3.00
210. Fibrous Quartz	56. Frieseite
H. Fichtelite	665. Fritzscheite, r
191. Fiedlerite	458. Fuchsite
589. Fillowite	391. Fuggerite, n 2.00
212. Fiorite	325. Funkite
270. Fire-marble, s.v 1.00	389. Gabbronite, n
212. Fire-opal	404. Gadolinite 2.50— 5.00
640. Fischerite	236. Gahnite 1.50— 2.50
250. Flêches d'Amour, s 2.50	45. Galena
210. Flexible Sandstone, s.v20	45. Galenite, s
580. Flinkite	118. Galenobismutite 2.50— 3.00
210. Flint	338. Gamsigradite
212. Float-stone 1.00	355. Ganomalite 1.50
548. Florencite, n	432. Ganophyllite 5.00
277. Flos-ferri 1.00	370. Garnet
203. Fluellite	483. Garnierite, A
196. Fluorerite 1.50	297. Gay-lussite
556. Fluor-Adelite, s.n	207. Gearksutite 1.00
549. Fluor-apatite20— 2.00	337. Gedrite
FLUORIDES, ETC., 164-209	392. Gehlenite
175. Fluorite	518. Geikielite, n
175. Fluor Spar, s20— 2.00 106. Foliated Tellurium 3.00	483. Genthite50 H. Geocerite
270. Fontainebleau Lime-	152. Geocronite 1.50
stone	
193. Footeite, n	H. Geomyricite
193. 1 Willie, II	II. GOMIJIIGIE

552.	Georgiadésite, n	2. Graphitoid, r
	Gerhardtite	148. Gray Copper, s\$1.00—\$2.00
574	Gersbyite, r	210. Greasy Quartz30
	Gersdorffite\$1.50—\$4.00	68. Greenockite 1.50— 4.00
	Geyerite	212. Green-opal
	Geyserite	510. Greenovite, 1.50
	Giallo Antico Marble75	751. Green Vitriol, s40— .75
264.	Gibbsite	270. Griotte Marble
458.	Gieseckite, n	555. Griphite, r40
458.	Gigantolite, n	469. Grochauite, r
458.	Gilbertite	448. Groddeckite, r
506.	Gillingite, r 2.00	370. Grossularite50— 3.00
H.	Gilsonite, s., Uintahite20	510. Grothite
	Giorgissite, r	338. Grünerite
212.	Girasol 1.00	31. Grünlingite, r
444.	Gismondite 2.00	59. Guadalcazarite, r
313.	Glassy Feldspar, s.v	30. Guanajuatite 2.00
	Glauberite	549. Guano, r
	Glauber Salt, s	742. Guanovulite, r
374.	Glaucochroite, n 9.00	512. Guarinite 3.00
	Glaucodot 1.00— 1.25	110. Guejarite 2.50
	Glaucolite 2.00	142. Guitermanite 1.00
	Glauconite	712. Gummite 2.00
	Glaucophane	65. Gunnarite, r
	Glaucopyrite	271. Gurhofite, s.v
648.	Globosite, r	746. Gypsum
	Glockerite	434. Gyrolite 3.00
	Gmelinite 1.50	365. Hackmanite, n
	Gold 1.50— 7.00	206. Hagemannite, r
	Gold Amalgam, r	616. Haidingerite 3.00
	Golden Beryl 2.50	333. Hainite, n
	Goldschmidtite, s 2.00— 3.00	166. Halite
	Gonnardite, n 1.00	480. Hallite, r
	Gorceixite, r	493. Halloysite
	Goslarite 1.00— 2.00	HALOIDS, 164-209
	Göthite 1.00— 2.00	769. Halotrichite
	Göthite in Quartz 1.50	696. Hambergite
	Goyazite	548. Hamlinite 8.00
	Graftonite, n 8.∞	409. Hancockite, n 1.00— 1.50
	Grandidierite, n	733. Hanksite
	Granular Quartz	623. Hannayite
	Granuline, r	H. Hard Coal, s. Anthracite 20
	Graphic Tellurium, s 2.00	355. Hardystonite50— 3.00
2.	Graphite	212. Harlequin Opal 2.00

442.	Harmotome\$1.00	338. Hillängsite
	Harstigite	411. Hillebrandite, n
H.	Hartite	334. Hiortdahlite\$3.00
679.	Harttite, r	506. Hisingerite
338.	Hastingsite	116. Histrixite, r
H.	Hatchettite50	270. Hislopite
521.	Hatchettolite	505. Hoeferite, r
75.	Hauchecornite, n	600. Hærnesite
86.	Hauerite 1.00	402. Homilite 2.00
462.	Haughtonite	587. Hopeite
243.	Hausmannite75— 3.00	338. Hornblende20— 1.50
599.	Hautefeuillite, r	210. Hornblende in Quartz
<b>3</b> 63.	Hatiyne, s	169. Horn Silver, s 1.25— 3.00
363.	Hatiynite	210. Horn Stone
447.	Haydenite 1.00	36. Horsfordite
401.	Haytorite	376. Hortonolite, A 6.00
325.	Hectorite, r	266. Houghite, r
325.	Hedenbergite50— 1.50	701. Howlite 1.00
552.	Hedyphane, r 1.50	166. Huantajayite, r
706.	Heintzite	45. Huascolite, r
210.	Heliotrope, s.v	813. Hübnerite 1.00— 2.00
417.	Hellendite, n	691. Hulsite, r
366.	Helvite 1.00— 1.50	391. Humboldtilite 2.00— 4.00
627.	Hemafibrite	823. Humboldtine 4.00
232.	Hematite20— 2.00	414. Humite 2.00— 8.00
581.	Hematolite	624. Hureaulite
655.	Henwoodite, r 2.00	536. Hussakite
66.	Hepatic Cinnabar 1.50	115. Hutchinsonite, n 9.00
	Hercynite	394. Hyacinth
	Herderite 4.∞	212. Hyalite
	Herrengrundite 1.25	314. Hyalophane 1.50
	Herrerite 1.50	376. Hyalosiderite 1.00
447.	Herschelite, s.v 1.00— 2.00	356. Hyalotekite
	Hessenbergite, Supplement	264. Hydrargillite, s40— 2.00
43.	Hessite 2.50	270. Hydraulic Limestone
269.	Heterogenite, ap	549. Hydroapatite, r
544.	Heterosite, r	462. Hydrobiotite, r
	Heubachite, ap	710. Hydroboracite
	Heulandite75— 1.50	294. Hydrocalcite, r
	Hexagonite	Hydrocarbons—Described
	Hibschite, n	at end of Dana's Sys-
	Hiddenite 2.50	tem, following No. 824.
531.	Hielmite 2.00	310. Hydrocastorite, r
185.	Hieratite	292. Hydrocerussite 2.00— 3.00

208 I	Hydroconite, r	T 77.0	Iodobromite
			Iodyrite\$1.00—\$8.00
			Iolite
	Hydrodolomite, r75— 2.00		Iridium 2.00— 2.50
	Hydrofluorite, r		Iridosmine 2.00— 3.00
-	Hydrogiobertite		Iron 1.00— 3.00
			Iron, Chromic, s20— .50
			Iron Garnet40— 3.00
		234.	Iron-magnesia Spinel
	Hydrophane 1.50		s.v40— 3.00
-	· ·		Iron, Magnetic, s20— 1.25
-	Hydrophite, r		Iron, Meteoric 1.00— 3.00
			Iron-natrolite
	Hydrosamarskite		Iron Pyrites, s20— 5.00
	Hydrotalcite		Iron, Terrestrial 1.00— 3.00
			Iron, Titaniferous, s25— 3.00
			Iron Tourmaline
			Irvingite
291. I			Iserine, r
324. I	Hypersthene 1.00— 3.00	250.	Iserite, r
316. I	Hyposclerite	626.	Isoclasite
233. I	Hystatite	210.	Itacolumite
223. I	[ce	458.	Ivigtite
270. I	[celand Spar 1.00— 2.00	526.	Ixiolite, A. r
376. I	Iddingsite, B. r	240.	Jacobsite 1.00— 1.50
393. I	[docrase, s40— 8.00	328.	Jade, s, also s. of Ne-
H. I	[drialite 1.00		phrite 1.50 3.00
798. I	[drizite, r	328.	Jadeite 1.50— 3.00
378. I	[gelströmite 2.00		Jalpaite, r
	Chlëite	130.	Jamesonite 1.00— 1.50
747. I	[lesite		Jargon 1.00
			Jarosite 1.00— 3.00
			Jasper
			Jasperized Wood
			Jasponyx
			Jasp-opal
			Jaspery Clay Iron-stone20
			Jefferisite
	Indiferous Blende 1.00		Jeffersonite 1.50
-	Indigo Copper, s 1.00— 3.00		Jelletite
			Jeremejevite
	Inesite 1.25— 2.50		Jet, Mineral Coal
			Johannite
	IODIDES, ETC., 164-209		Johnstrupite 1.50
•		2-2.	J

506.	Jollyte, r	H. Könlite
	Jordanite\$2.50	520. Koppite, A\$1.00
	Josëite	774. Kornelite, r
	Josephinite 1.∞	429. Kornerupine 2.00
	Jossaite, r	468. Kotschubeite
	Kaersutite	604. Köttigite
	Kainite	233. Kragerö Hematite
	Kalgoorlite, r	236. Kreittonnite
	Kaliborite, r	198. Kremersite
	Kalicine, r	105. Krennerite 5.00
	Kalinite	776. Kröhnkite 1.50— 5.00
	Kaliophilite	762. Krugite, r50
	Kalk-cancrinite, r	429. Kryptotil, r
	Kallilite, n	270. Ktypeite, r50
	Kamarezite, A. r	744. Kubeite, s.r
	Kämmererite, A75— 2.∞	327. Kunzite 2.00
	Kaolinite20— .50	337. Kupfferite
	Kauaiite, r	14. Küstelite
	Kehoeite, r	162. Kylindrite, s.n 1.25
	Keilhauite 1.00— 2.00	319. Labradorite30— 2.00
	Kentrolite 2.∞	702. Lagonite
	Kermesite 2.50— 3.00	269. Lampadite, r
	Kerrite, r	805. Lamprophanite, r
	Kerstenite, r	514. Lamprophyllite, r
	Kertschenite, r	737. Lanarkite 5.00
	Keweenawite, n 5.00	270. Landscape Marble 1.25
	Kibdelophane	419. Långbanite 2.00
232.	Kidney Ore	718. Langbeinite, n
	Kieserite	779. Langite 1.50
	Kilbrickenite 1.50	302. Lansfordite
327.	Killinite, r	298. Lanthanite 2.50
338.	Kirwanite, r	365. Lapis-Lazuli, s75— 3.00
284.	Kischtimite, r	703. Larderellite 2.00
553-	Kjerulfine 2.00— 3.00	457. Lasallite, r
	Klaprotholite	549. Lasurapatite
164.	Kleinite, n	320. Latrobite
	Klementite, r 1.00	446. Laubanite
	Knebelite 1.25— 2.00	445. Laumontite40— 1.00
	Knopite, n 1.50	189. Laurionite 1.00
	Knoxvillite	94. Laurite
-	Kobellite 8.00	820. Lautarite, n
	Koenenite, n	158. Lautite, r
	Kongsbergite	596. Lavendulan, r 2.00
	Koninckite 1.50	332. Låvenite 4.00

325.	Lavrovite	259. Limonite\$ .20—\$1.00
178.	Lawrencite\$2.00	741. Linarite 2.00
423.	Lawsonite, n75— 1.50	681. Lindackerite
	Lazulite	79. Linnæite 1.50— 2.00
	Lazurfeldspar	456. Lintonite
	Lazurite	654. Liroconite 2.00
	Lead 1.00	644. Liskeardite 2.00
	Leadhillite 1.00— 4.00	460. Lithia Mica, s20— 1.00
	Lecontite	544. Lithiophilite
	Ledouxite, r	269. Lithiophorite, r 1.00
	Lehrbachite 3.00	461. Lithium-iron Mica, s .40— 1.50
142.	Lengenbachite, r 4.00	270. Lithographic Stone20
	Lennilite, r	492. Lithomarge50
	Lenticular Iron Ore20	122. Liveingite, r
	Lenzinite	109. Livingstonite 2.50
	Leonhardite 1.00	237. Lodestone
	Leonite, n	338. Loganite, r
	Lepidocrocite 2.00	468. Loganite, A
	Lepidolite20— 1.00	97. Löllingite35— 1.50
462.	Lepidomelane, B60	120. Lorandite, n 2.50
	Lettsomite, s 3.00	529. Loranskite, r
325.	Leucaugite	514. Lorenzenite, n
468.	Leuchtenbergite 2.00	679. Lossenite, n 1.50
	Leucite	325. Lotalite
	Leucochalcite 1.00	457. Lotrite, r
435.	Leucocyclite 2.00	757. Löweite
H.	Leucopetrite	802. Löwigite
351.	Leucophanite 1.50— 3.00	313. Loxoclase
416.	Leucophoenicite, n 1.00	480. Lucasite, r
97.	Leucopyrite	638. Ludlamite 3.00
312.	Leucosphenite, n	694. Ludwigite 1.00— 3.00
492.	Leverrierite, r	270. Lumachelle 1.00
449.	Levynite	682. Lüneburgite 2.00
670.	Lewisite, n	212. Lussatite, r 1.50
562.	Libethenite 2.00	458. Lythrodes, r
458.	Liebenerite, r	712. Mackintoshite, n
308.	Liebigite	480. Maconite, r
417.	Lievrite, s	270. Madreporic Marble
145.	Light Ruby Silver, s 1.25— 3.00	462. Magnesia Mica, A.s20— .75
	Lignite, s., Brown Coal20	237. Magnesian Magnetite
140.	Lillianite	233. Magnesian Menaccanite
<b>481</b> .	Limbachite, r	234. Magnesia Spinel, s.v40— 1.00
230.	Lime, r	234. Magnesia-Iron Spinel, s.v 2.00
270.	Limestone, s20— 1.25	238. Magnesioferrite 2.00

272.	Magnesite\$ .20—\$1.00	490. Marl\$ .20
426.	Magnesium Tourmaline 40	338. Marmairolite
462.	Magnesium-iron	58. Marmatite 1.50
	Mica, s	481. Marmolite
237.	Magnetic Iron, s20— 1.25	165. Marshite, n 6.00
20.	Magnetic Platinum . 1.50— 4.00	620. Martinite
74.	Magnetic Pyrites, s .20— 1.50	232. Martite, r
237.	Magnetite20— 1.25	714. Mascagnite 1.50
241.	Magnochromite	319. Maskelynite, r
810.	Magnolite, r	466. Masonite
288.	Malachite	229. Massicot 2.00
	Malacolite	120. Matildite
394.	Malacon, r	186. Matlockite 2.00— 3.00
752.	Mallardite	376. Matricite, r
762.	Mamanite, r	670. Mauzeliite, n
270.	Mandelato Marble75	651. Mazapilite 4.00
<b>398</b> .	Manganandalusite, r	485. Meerschaum, s
549.	Manganapatite	386. Meionite 1.50
	MANGANATES, ETC., 234-247	230. Melaconite 1.00
262.	Manganbrucite 1.50	544. Melanchlor, r
510.	Manganesian Titanite 1.50	370. Melanite50
370.	Manganesian Calcium-	348. Melanocerite
	iron Garnet40— .150	230. Melanochalcite, r
325.	Manganhedenbergite 1.50	479. Melanolite, n
	Manganite 1.00— .300	211. Melanophlogite, r
237.	Manganmagnetite	506. Melanosiderite, r
270.	Manganocalcite	421. Melanotekite 1.00
434-	Manganocalcite, s.r	193. Melanothallite, r
	Manganocalcite 1.50	751. Melanterite
462.	Manganophyllite75— 1.25	391. Melilite
	Manganosite 2.00	352. Meliphanite 1.00
	Manganostibiite	457. Melite, r
	Manganotantalite 1.50— 3.00	824. Mellite 1.∞
	Manganpectolite 1.00	77. Melonite 9.∞
	Manganspherite	233. Menaccanite25— 3.00
	Mangan-vesuvianite 2.∞	187. Mendipite 3.∞
	Marble	766. Mendozite 4.00
	Marble of Languedoc	151. Meneghinite 1.00
	Marcasite	212. Menilite
	Marceline, r	58. Mercurial Blende
	Margarite	148. Mercurial Tetrahedrite, s 2.00
	Margarodite	164. Mercuric Chloride, r
	Marialite 5.00	16. Mercury 1.00
520.	Marignacite	462. Meroxene, see note

272.	Mesitite, A\$ 50—\$2.50	811. Molybdomenite, r
456.	Mesole 1.00	420. Molybdophyllite, n
455.	Mesolite 1.50— 2.50	181. Molysite
593-	Messelite	537. Monazite\$ .40—\$2.00
619.	Metabrushite	560. Monetite 1.00
471.	Metachlorite, r 1.25	539. Monimolite
59.	Metacinnabarite 1.00— 2.50	325. Monradite, r
	Metastibnite, r	808. Montanite
797.	Metavoltine	374. Monticellite50— 2.00
481.	Metaxoite, r	496. Montmorillonite
	Meteoric Iron 1.00— 3.00	229. Montroydite, n 8.00
	Meteoric Stone (Aerolite) 1.50	430. Monzonite, ap
	Mexican Onyx	313. Moonstone
	Meymacite, r 4.00	316. Moonstone
121.	Miargyrite 6.00	463. Moravite, r
	Micaceous Iron Ore40	437. Mordenite
	Micaceous Quartz	506. Morencite, r
212.	Michaelite	750. Morenosite
315.	Microcline20— 2.00	423. Moresnetite, r75— 1.∞
	Microlite 1.50	559. Morinite, r
361.	Microsommite 4.00	516. Mosandrite 1.00
	Miemite	210. Moss-agate75— 1.50
165.	Miersite, n	527. Mossite, n
311.	Milarite 3.00	277. Mossottite 1.50
212.	Milk-opal	567. Mottramite, r
210.	Milky Quartz20— .50	338. Mountain Cork
	Millerite 1.00— 2.00	338. Mountain Leather50
551.	Mimetite 1.50— 2.00	338. Mountain Wood50
Η.	Mineral Coal20— .40	505. Müllerite, r
614.	Minervite, r	313. Murchisonite
244.	Minium 4.∞	409. Muromontite, r
743.	Mirabilite	Mursinskite, supplement.
735.	Misenite	458. Muscovite ,20— 1.50
	Mispickel, s25— 1.25	325. Mussite 1.00
241.	Mitchellite	672. Nadorite 1.50
668.	Mixite 1.00— 1.50	394. Naëgite 5.00
388.	Mizzonite	106. Nagyagite 3.00— 4.00
210.	Mocha-stone, s.v 1.50	270. Nail-head Spar
	Mohawkite, r 6.00	269. Namaqualite, ap
	Moissanite, r	514. Narsarsukite, n
	Molybdenite40— 1.00	165. Nantokite 2.00
	MOLYBDATES, ETC., 812-820	361. Nasonite, n 2.00
219.	Molybdic Ocher, s 1.00— 2.00	NATIVE ELEMENTS, 1-25
219.	Molybdite 1.00— 2.00	761. Natrochalcite, n 3.00
		<del>-</del>

B01.	Natrojarosite	691. Nordenskiöldine
453.	Natrolite\$ .75—\$3.00	428. Nordmarkite
	Natron	286. Northupite, n\$ .75—\$1.00
	Natron-catapleiite	364. Nosean, s 1.25— 3.00
	Natrophilite	364. Noselite 1.25— 3.00
560.	Natrophite, r	674. Ochrolite
	Naumannite 4.00	252. Octahedrite
	Necronite	458. Oellacherite
	Needle Ore 2.∞	394. Œrstedite, r
	Nemalite 1.25	441. Offrétite, r 1.00
	Neociano, ap 2.00	407. Oisanite 1.50
	Neotantalite, A.n	433. Okenite 2.00
	Neotocite 2.50	316. Olafite
	Nephelite	64. Oldhamite
	Nephrite	317. Oligoclase
	Nepouite, A.n	273. Oligonite, manganiferous
	Neptunite, n 2.00	561. Olivenite 1.50— 2.00
	Nesquehonite	376. Olivine
	Neurolite, r	325. Omphacite
	Nevyanskite 3.00	458. Oncosine 1.00
	Newberyite	210. Onegite, s.v 1.50
	Newtonite	61. Onofrite
71.	Niccolite 1.00— 3.00	210. Onyx
	Nickel Glance, s 1.50— 4.00	270. Onyx, Mexican
98.	Nickeliferous Arsenopyrite	270. Oölite
	Nickeliferous Iron	458. Oosite, n
	(Awaruite) 3.∞	212. Opal
237.	Nickeliferous Magnetite	212. Opal-agate 1.50
85.	Nickeliferous Pyrite, s	212. Opalized Wood
· 74·	Nickeliferous Pyrrhotite20	481. Ophicalcite
237.	Nickel Oxide, r	395. Orangite 4.00— 6.00
95.	Nickel-skutterudite	231. Oriental Amethyst 3.00
250.	Nigrine, ferriferous rutile50	231. Oriental Emerald 3.00
684.	Niter	231. Oriental Ruby 1.00— 4.00
	NITRATES, 683-690	231. Oriental Topaz 2.00
687.	Nitrobarite	37. Orileyite, r
685.	Nitrocalcite	27. Orpiment 1.00— 3.00
690.	Nitroglauberite	409. Orthite, s50— 1.50
	Nitromagnesite	313. Orthoclase
	Nivenite	330. Osmelite
	Nocerite 1.00	549. Osteolite, s.v
	Nohlite, r	291. Otavite, r
	Non-caking Coal,20	467. Ottrelite
	Noralite	270 Ouverovite s v Too— 200

822	Oxammite	509. Penwithite, r
022.	Oxides, 210-269	353. Peplolite, r
	Oxychlorides, 186-194	192. Percylite\$4.00
456	Ozarkite\$ .75	225. Periclase 1.25— 2.00
450.	Oxyfluorides, 195-196	316. Pericline
	OXYSULPHIDES, 107-108	
ш		376. Peridot, s
	Ozocerite	426. Peridot, Ceylon 3.00
-		426. Peridot, Brazilian50— 2.00
	Pacite, r. (	316. Peristerite 1.00
	Pageite, r (=Hulsite?)	518. Perovskite
	Pagodite, s.n 1.00	313. Perthite, r
	Paisbergite 1.00	310. Petalite
	Palacheite	210. Petrified Wood, Jasperized30
	Paligorskite, r	212. Petrified Wood, Opalized40
	Palladium	H. Petroleum
	Palladium Gold, s.v 5.00	44. Petzite 3.00
	Palmerite, r	338. Phäactinite, r
	Palmierite, n	447. Phacolite 1.00— 2.00
	Pandermite, r	617. Pharmacolite 1.25
	Panno-di-Morte Marble	646. Pharmacosiderite 1.50— 4.00
	Papierspath	324. Phästine, r
	Paragonite	382. Phenacite 1.00— 4.00
	Parahopeite, n	480. Philadelphite, r
	Paralaurionite, n	776. Phillipite, r
	Paralogite, n	441. Phillipsite 1.00— 1.50
	Paraluminite	462. Phlogopite, A20— .75
	Paramelaconite, r	726. Phœnicochroite
	Paratacamite, n	491. Pholidolite
	Pargasite	286. Phosgenite
	Parisite 6.00	549. Phosphate Rock
	Partschinite	PHOSPHATES, ETC., 536-690
	Partzite, r 2.00	549. Phosphatic Nodules, r20
	Patronite, r	549. Phosphorite, s.v
	Pattersonite, n	609. Phosphosiderite
	Pearceite, n	664. Phosphuranylite
	Pearl Sinter 1.50	335. Photicite, r
271.	Pearl Spar 1.00	467. Phyllite
H.	Peat, related to mineral coal .30	648. Picite, r
	Pectolite	768. Pickeringite 1.00
	Peganite	234. Picotite
232.	Pencil Ore	768. Picroallumogene, r 1.25
	Penfieldite, n	407. Picroepidote, r
	Penninite, A50— 2.00	481. Picrolite
65.	Pentlandite 2.00	760. Picromerite

595. Picropharmacolite	458. Polyargite, n
325. Picrophyll, r\$ .75	157. Polyargyrite
456. Picrothomsonite, r	156. Polybasite\$2.00—\$2.50
233. Picrotitanite, s.v	535. Polycrase 3.00
337. Piddingtonite, r	353. Polychroilite, r
408. Piedmontite75— 2.00	75. Polydymite 3.00
824. Pigotite, r 1.50	762. Polyhalite
483. Pimelite, A.r	461. Polylithionite
695. Pinakiolite 1.25	533. Polymignite 6.00
458. Pinite, n	550. Polysphærite
458. Pinitoid, n	13. Porpezite 5.00
705. Pinnoite 2.00	270. Portor Marble
272. Pinolite	313. Potash Feldspar, s20— 1.50
296. Pirssonite, n 2.00	458. Potash Mica, s20— 1.50
753. Pisanite 1.25	816. Powellite
270. Pisolite, s.r	210. Prase
272. Pistomesite, A50	353. Praseolite, r
711. Pitchblende 3.00	212. Precious Opal 1.00— 2.00
325. Pitkärantite, r	411. Prehnite
678. Pitticite 2.00	388. Prehnitoid
801. Plagiocitrite, r	58. Pribramite
122. Plagionite 2.00— 3.00	704. Priceite, r
504. Plancheite, n	429. Prismatine 2.00
611. Planerite, r	469. Prochlorite
791. Planoferrite, r	185. Proidonite, r
210. Plasma	413. Prolectite, n
67. Platiniferous Covellite 2.∞	204. Prosopite 5.00
20. Platinum 1.50— 4.00	325. Protheite
251. Plattnerite 7.∞	461. Protolithionite, r
120. Plenargyrite, r	480. Protovermiculite, r
234. Pleonaste, s.v 40.— 3.00	145. Proustite 1.25— 3.00
435. Plombierite, n	549. Pseudoapatite
2. Plumbago, s	538. Pseudoberzeliite, r
498. Plumballophane	462. Pseudobiotite, r
229. Plumbic Ocher, s 2.00	246. Pseudobrookite 2.00
270. Plumbocalcite 1.25	180. Pseudocotunnite, r
241. Plumboferrite, r	570. Pseudomalachite 1.50
658. Plumbogummite 6.00— 9.00	437. Pseudonatrolite, r
801. Plumbojarosite	468. Pseudophite, A
108. Plumbostannite, ap	389. Pseudo-Scapolite, n
676. Podolite, r	344. Pseudosmaragd, r
249. Polianite 1.00— 2.00	210. Pseudomorphous
322. Pollucite 3.00— 5.00	Quartz
370. Polyadelphite40— 1.50	269. Psilomelane

567. Psittacinite\$6.00	804. Quetenite
462. Pterolite, B.r	16. Quicksilver, s\$1.00
The state of the s	
436. Ptilolite 1.50	3. Quisqueite, r
542. Pucherite 2.50	269. Rabdionite, ap
270. Pudding-stone	461. Rabenglimmer
210. Pudding-stone	210. Radiated Quartz
608. Purpurite, n	481. Radiotite
397. Pycnite	789. Raimondite
469. Pycnochlorite	208. Ralstonite 2.50
458. Pycnophyllite	100. Rammelsbergite 1.00
325. Pyrallolite, r	212. Randannite
484. Pyrallolite,	309. Randite, r
353. Pyrargillite, r	457. Ranite
144. Pyrargyrite 1.00— 2.50	232. Raphisiderite, r
85. Pyrite	813. Raspite, n 6.00
98. Pyrites, Arsenical, s25— 1.25	462. Rastolyte, r
96. Pyrites, Cockscomb	127. Rathite, n 7.00
83. Pyrites, Copper, s35— 2.00	353. Raumite, r
85. Pyrites, Iron, s30— 5.00	496. Razoumovskyn, r
74. Pyrites, Magnetic, s 20.— 1.50	26. Realgar
96. Pyrites, Spear 1.00	492. Rectorite, r 1.00
84. Pyrites, Tin, s75— 3.00	232. Red Chalk, s.v
267. Pyroaurite 2.00	594. Reddingite
520. Pyrochlore 1.25— 2.00	232. Reddle
263. Pyrochroite 1.50— 2.50	785. Redingtonite, r
254. Pyrolusite	232. Red Ocher
550. Pyromorphite30— 2.50	54. Redruthite, s 1.50— 2:00
370. Pyrope	483. Refdanskite, A, r
233. Pyrophanite, n	162. Regnolite, r
497. Pyrophyllite	270. Reichite
H. Pyroretinite	819. Reinite 8.00
409. Pyrorthite	304. Remingtonite
480. Pyrosclerite, r 1.25	484. Rensselaerite
385. Pyrosmalite 2.50	212. Resin-opal
146. Pyrostilpnite 4.00	481. Retinalite
325. Pyroxene	H. Retinite (amber-like
522. Pyrrhite, r	resins)
74. Pyrrhotite	577. Retzian, n
210. Quartz	113. Rezbanyite
210. Quartz Breccia	25. Rhabdite, r
210. Quartz Conglomerate20	605. Rhabdophanite
210. Quartz Inclusions50— 2.00	667. Rhagite
210. Quartzose Sandstone .20— .40	13. Rhodite
773. Quenstedtite	699. Rhodizite
110. €	-77.

274. Rhodochrosite\$ .75-\$4.00	58. Ruby Blende \$ .75
335. Rhodonite	224. Ruby Copper, s50— 3.00
343. Rhönite, n	145. Ruby Silver, s, 144
313. Rhyacolite 1.00	and 145 1.00— 3.00
210. Riband Jasper 1.00	234. Ruby Spinel40— 1.00
653. Richellite, r 1.00	270. Ruin Marble 1.00
155. Richmondite, r	479. Rumpfite
264. Richmondite, r	711. Rutherfordine
338. Richterite 1.00	250. Rutile
57. Rickardite, n	99. Safflorite 2.00
70.1 1.5	210. Sagenitic Quartz 2.50
340. Riebeckite,	168. Sal-ammoniac75— 1.00
149. Rionite	325. Salite
	466. Salmite
468. Ripidolite, s50— 2.50 147. Rittingerite	44 0 1
222. Rivotite, r	100. Salt, s
H. Rochlederite	753. Salvadorite, r
210. Rock Crystal	529. Samarskite 2.50
, <b>.</b> .	257. Sammetblende 1.25
	498. Samoite, r
	149. Sandbergerite 3.00
430. Roeblingite, n 4.00 379. Roepperite, A 1.50— 4.00	210. Sandstone, Flexible
535. Rogersite, ap 1.50	<b>~</b>
671. Romeite	313. Sanidine
778. Römerite 1.25	231. Sapphire40— 1.00
290. Rosasite, r	210. Sapphire-quartz
463. Roscoelite 3.00	430. Sapphirine 2.00
480. Roseite, r	390. Sarcolite 2.50
590. Roselite 2.50	555. Sarcopside, r
331. Rosenbuschite 2.50	210. Sard
210. Rose Quartz30— 2.00	
458. Rosite, n	270. Sarencolin Marble
622. Rösslerite, r	557. Sarkinite 2.50
344. Rosterite, r	115. Sartorite 5.00
370. Rothoffite 1.00	457. Sasbachite, ap
405. Rowlandite, r 5.00	265. Sassolite
462. Rubellan, r	270. Satin Spar 1.00
426. Rubellite	746. Satin Spar, s.v40— .60
257. Rubinglimmer, s.v 1.00	179. Scacchite
783. Rubrite, r	387. Scapolite, s20— 2.00
231. Ruby, Oriental 1.00— 4.00	58. Schalenblende
234. Ruby, Balas	69. Schalenblende 1.00— 1.25

129.	Schapbachite	465. Seybertite\$ .75—\$1.25
814.	Scheelite\$ .75—\$3.00	270. Shell-Marble
H.	Scheererite	25. Siderazot, r
325.	Schefferite	273. Siderite
585.	Schertelite, r	210. Siderite
324.	Schiller Spar, s.r	25. Siderite, s 3.00
125.	Schirmerite	273. Siderodot
336.	Schizolite, n	25. Siderolite, s 2.50
	Schneebergite, r	799. Sideronatrite 2.00
371.	Schorlomite 1.00	462. Siderophyllite
25.	Schreibersite, r 6.00	273. Sideroplesite 60
309.	Schröckinergite, r	755. Siderotil, r
500.	Schrötterite 1.25	79. Siegenite 2.∞
	Schungite, s.r	270. Siena Marble
	Schwartzembergite 4.00	338. Silfbergite
148.	Schwatzite 2.00	Silicates, 310-519
747.	Scleropasthite, r	210. Siliceous Sinter 1.25
454.	Scolecite 1.25— 2.00	212. Siliceous Sinter,s.v75— 1.50
607.	Scorodite 1.00— 3.00	210. Silicified Shells50
407.	Scorza	210. Silicified Wood30
506.	Scotiolite	212. Silicified Wood
447-	Seebachite, s.v 1.00— 2.00	430. Silicomagnesiofluorite, r
<b>768</b> .	Seelandite	399. Sillimanite30— .40
	SELINIDES, ETC., 35-108	14. Silver 1.00— 7.00
	Seleniferous Bismuthinite	153. Silver, Brittle, s 2.00— 3.00
118.	Seleniferous Galeno-	144. Silver, Dark Ruby, s. 1.00— 2.50
	bismutite 3.00	42. Silver Glance, s 1.25— 2.50
746.	Selenite	169. Silver, Horn, s 1.25— 3.00
	SELENITES, ETC., 808-811	145. Silver, Light Ruby, s. 1.25— 3.00
	Selenium	524. Sipylite 3.∞
	Selensulphur	22. Siserskite
	Selen-Tellurium	466. Sismondine 1.00
	Seligmannite, r	526. Skogbölite, A 2.00
	Sellaite 6.00	95. Skutterudite 8.00
	Semiopal	457. Sloanite, ap
	Semseyite 9.00	87. Smaltite 1.00— 2.50
233.	Senaite, n	338. Smaragdite
214.	Senarmontite	493. Smectite
485.	Sepiolite	119. Smithite, n
458.	Sericite	275. Smithsonite
481.	Serpentine20— 1.50	210. Smoky Quartz25— 2.00
481.	Serpentine Marble75— 1.00	484. Soapstone, s.v
782.	Serpierite 2.00	538. Soda-berzeliite
430.	Sevendibite, r	316. Soda Feldspar, s20— 1.00

<b>362</b> .	Sodalite\$ .75—\$2.50	56. Sternbergite\$3.00
	Soda Niter	222. Stetefeldtite, r
459.	Sodium Mica, s	222. Stibianite, r
H.	Soft Coal, s. Bituminous Coal .20	583. Stibiatil, r
	Sonomaite, r	222. Stibiconite
338.	Soretite	37. Stibiodomeykite
487.	Spadaite	222. Stibioferrite, r
441.	Spangite, r	526. Stibiotantalite, A.r 2.50— 9.00
732.	Spangolite	28. Stibnite
273.	Spathic Iron, s20— 3.00	210. Stibnite in Quartz 2.00
	Spear Pyrites 1.00	443. Stilbite
232.	Specular Iron, s.v20— 2.00	473. Stilpnochloran, r
93.	Sperrylite 3.00	474. Stilpnomelane
	Spessartite	270. Stinkstone, s
	Sphærite	618. Stofferite
	Sphærocobaltite 4.00	422. Stokesite, n
273.	Sphærosiderite	817. Stolzite 3.00— 6.00
	Sphalerite20— 1.50	325. Strakonitzite, r
	Sphene, s	335. Stratopeite, r
	Sphenoclase, ap	248. Stream Tin
	Spinel	608. Strengite2.00
	Spodiophyllite, r	475. Strigovite
	Spodiosite	389. Stroganovite, n
	Spodumene20— 2.50	55. Stromeyerite 2.50— 8.00
	Sprudelstein	280. Strontianite
	Spurrite, n	270. Strontianocalcite 2.00
	Staffelite50	527. Strüverite, r
	Stalactite	585. Struvite
270.	Stalagmite	41. Stützite
	Stanniferous Blende.	768. Stüvenite, r
	Stannite	141. Stylotypite
	Star Quartz 1.00	H. Succinite
	Star Sapphire 40— 1.50	710. Sulfoborite, n
	Stassfurtite, s	211. Sulfuricin, r
	Statuary Marble	Sulphantimonates, etc., 158-163
	Staurolite	Sulphantimonites, etc., 109-157
	Steatargillite, n	Sulpharsenates, etc., 158-163
	Steatite, s	SULPHARSENITES, ETC., 109-157
	Steeleite, r	SULPHATES, ETC., 714-807
	Steenstrupine, r 3.00	SULPHIDES, ETC., 35-108
	Steinmannite 1.50	Sulphobismuthites, etc., 109-
	Steltznerite, n	157
	Stephanite 2.00— 3.00	728. Sulphohalite
515.	Stercorite	3. Sulphur

31. Sulphurous Tetra-	84. Teallite, n
dymite\$1.50—\$2.00	Tellurates, etc., 808-811
159. Sulvanite, n 1.00	Tellurides, etc., 35-108
317. Sunstone, s.v40— .75	218. Tellurite\$6.∞
316. Sunstone, s.v	TELLURITES, ETC., 808-811
210. Sunstone, s.v 1.00— 1.25	7. Tellurium 1.00
734. Susannite, r 8.00	305. Tengerite 2.50
693. Sussexite 3.00	149. Tennantite 2.50— 3.00
550. Svabite, n 2.50	230. Tenorite 1.00— 2.50
679. Svanbergite 3.00	379. Tephroite 1.25— 2.00
75. Sychnodymite, n	389. Terenite, n
104. Sylvanite 2.00— 3.00	188. Terlinguaite, n 9.00
167. Sylvite	493. Termierite, r
598. Symplesite 2.00	25. Terrestrial Iron 1.00— 3.00
579. Synadelphite 4.00	287. Teschemacherite
284. Synchisite, r	435. Tesselite
756. Syngenite 3.00	31. Tetradymite 1.50— 2.00
324. Szaboite	148. Tetrahedrite 1.00— 2.00
697. Szaibelyite 1.50	337. Thalackerite
338. Szichenyite	
745. Szmikite	405. Thalénite, n 3.00
	502. Thaumasite
212. Tabasheer, r	716. Thenardite
	294. Thermonatrite 1.50
202. Tachhydrite	481. Thermophyllite
394. Tachyaphaltite, r	270. Thinolite, r 1.00 273. Thomäite, r
480. Tænislite, n	206. Thomsenolite 1.00— 1.50
630. Tagilite	456. Thomsonite
484. Talc	711. Thorianite, r 2.50
555. Talktriplite	395. Thorite 2.50— 6.00
193. Tallingite, r	712. Thorogummite, r 2.00
767. Tamarugite 1.00	406. Thulite
320. Tankite	473. Thuringite
TANTALATES, ETC., 520-535	60. Tiemannite 2.50— 6.00
526. Tantalite 1.50— 3.00	210. Tiger-eye
143. Tapalpite	556. Tilasite, n
527. Tapiolite 4.00— 8.00	224. Tile Ore
352. Taramellite, n 645. Taranakite, r	47. Tilkerodite
727. Tarapacaite, r 1.00	84 Tin Puritee s re- soo
563. Tarbuttite, n	84. Tin Pyrites, s75— 3.00 248. Tinstone, s30— 2.50
277. Tarnowitzite 1.50	Titanates, etc., 510-519
H. Tasmanite	233. Titanic Iron, s.r
748. Tauriscite, r	325. Titaniferous Augite
575. Tavistockite	370. Titaniferous Calcium-
715. Taylorite	iron Garnet

237. Titaniferous Magnetite	635. Tyrolite\$1.00
510. Titanite	
376. Titan-olivine, B	
510. Titanomorphite	411. Uigite, r
435. Tobermorite, n	H. Uintahite, related to elaterite .20
173. Tocornalite, r	708. Ulexite
397. Topaz	
370. Topazolite 1.00-	
659. Torbernite 1.50-	
481. Totaigite, r	409. Uralorthite
210. Touchstone, s.v	
426. Tourmaline20-	-
210. Tourmaline in Quartz .50-	
269. Transvaalite, ap	663. Uranocircite 3.00
325. Traversellite	503. Uranophane 2.00
270. Travertine	
119. Trechmannite, n	713. Uranosphærite
	- 1.50   662. Uranospinite
596. Trichalcite	307. Uranothallite
211. Tridymite 1.∞-	- 1.50   395. Uranothorite
380. Trimerite	326. Urbanite, n
543. Triphylite	786. Utahite
555. Triplite	
556. Triploidite	
212. Tripoli Slate	
212. Tripolite	
675. Trippkeite	337. Valléite, r
670. Tripuhyite, n	1 33
350. Tritomite	Vanadates, etc., 536-690
665. Trögerite	552. Vanadinite 1.00— 1.50
73. Troilite	1.50 718. Vanthoffite, n 2.50
645. Trolleite, r	210. Variegated Jasper 1.00
299. Trona	40 611. Variscite
381. Troostite	269. Varvicite, r
513. Tscheffkinite	3.00 727. Vauquelinite 5.00
316. Tschermakite	467. Venasquite
765. Tschermigite	1.25 481. Verde-antique
270. Tufa, Calc	60 270. Verde-antique Marble 30
TUNGSTATES, ETC., 812-82	480. Vermiculite, r
220. Tungstite	Vermiculites, 480
255. Turgite	50   393. Vesuvianite40- 8.00
275. Turkey-fat ore	
642. Turquois	75   529. Vietinghofite, r
286. Tychite, n	376. Villarsite, r

325. Violan\$1.50	370. Wiluite\$ .50
755. Vitriol, Blue, s50— 2.00	791. Winebergite, r
597. Vivianite	407. Withamite 1.50
309. Voglite 4.00	279. Witherite
633. Volborthite 3.00	137. Wittichenite 3.00
222. Volgerite, r	333. Wöhlerite 1.∞
796. Voltaite	103. Wolfachite
108. Voltzite	812. Wolframite
104. Von Diestite, r	329. Wollastonite
722. Vulpinite	212. Wood Opal
269. Wad, r	210. Wood, Silicified (Petrified)30
553. Wagnerite 2.00— 4.00	212. Wood, Silicified (Petrified)40
338. Waldheimite, r	248. Wood Tin 1.50
330. Walkerite	399. Wörthite
666. Walpurgite 2.00	818. Wulfenite 1.00— 2.00
306. Walthérite, r	H. Wurtzilite, related to elaterite .20
465. Waluewite, A 1.25	69. Wurtzite 1.00— 2.00
622. Wapplerite 1.50	572. Xantharsenite, r
642. Wardite, n 1.25	160. Xanthoconite 2.50
740. Waringtonite	465. Xanthophyllite, A 1.25
126. Warrenite	409. Xanthorite
700. Warwickite	260. Xanthosiderite
233. Washingtonite	399. Xenolite
409. Wasite, r	536. Xenotime 1.50— 3.∞
223. Water	338. Xiphonite
763. Wattevillite	435. Xonotlite, n
639. Wavellite	435. Xylochlore
212. Wax-opal, s.v	259. Yellow Ocher
33. Wehrlite	210. Yellow Quartz50
285. Weibyeite, r	370. Yttergarnet, s.v 2.00
352. Weinbergerite, r	405. Yttrialite 8.∞
313. Weissigite	370. Yttriferous Calcium-
442. Wellsite, n	iron Garnet 2.00
387. Wernerite	209. Yttrocerite
791. Werthemanite, r	519. Yttrocrasite, n
399. Westanite, r	712. Yttrogummite, r
136. Wheel Ore, s	528. Yttrotantalite 3.00
	511. Yttrotitanite, s 1.00— 2.00 303. Zaratite
39. Whitneyite 5.00 480. Willcoxite, r	Zeolites, 436-457
381. Willemite	434. Zeophyllite, s 3.00
481. Williamsite	613. Zepharovichite
92. Willyamite, n 8.00	660. Zeunerite 3.00
389. Wilsonite, n	277. Zeyringite
30y. 11 months, 11	all. Entingio

12. Zinc	114. Zinkenite\$2.00—\$2.50
805. Zincaluminite	723. Zinkosite
58. Zinc Blende, s \$.20—\$1.50	461. Zinnwaldite40— 1.50
271. Zinciferous Dolomite	394. Zircon
274. Zinciferous Rhodochrosite	518. Zirkelite, n
335. Zinciferous Rhodonite .75— 3.00	264. Zirlite, r
228. Zincite	406. Zoisite
270. Zincocalcite	457. Zonochlorite, ap 3.00
236. Zinc-Spinel, s 1.50— 2.50	52. Zorgite 3.00
749. Zinc Vitriol, s 1.00— 2.00	369. Zunyite 1.00
289. Zinkazurite, r	555. Zwieselite

# PART IV

Elementary Systematic Collections

The arrangement, apart from the silicates, is according to the metallic constituents. Adapted for a short course in any popular book for beginners.

## No. 14A. Normal or High School Collection

One hundred and eighty museum size specimens, averaging 12 x 9 cm. (4¾ x 3½ in.). Prepared especially to meet the demand among Normal and High Schools and private Academies for a practical reference collection, embracing only the common or most important species and varieties. The striking colors and choice crystallizations, in which the collection abounds, make it an attractive and essential feature in the class-room or school museum. The list includes every name in the summary of species as given in Prof. E. S. Dana's "Minerals and How to Study Them."

Indiviudal museum specimens may be purchased at double the (hand size) prices given after each name in the High School List. The sum of such individual values in the museum size is \$228.10. The "collection price" for all the specimens is \$180.00, delivered to any address. This price includes pasteboard trays (or blocks if requested) and three No. 3 Oak Chests, as shown in Plate IX. Without chests, 10 per cent. less. Mahogany 10-drawer cabinet \$45.00. See Plate.

Purchase in Parts. Free delivery with trays and No. 3

chest. Without chest, 10 per cent. less.

PART I. (School Collection No. 21A) 60 names marked with +, totaling \$58.40 ......\$50.00 PART II. 60 names marked with \*, totaling \$67.80 ... 50.00 PART III. 60 remaining names, totaling \$101.90 .... 80.00

#### No. 14. Student's Normal or High School Collection

One hundred and eighty hand size specimens averaging 10 x 7 cm. (4 x 2¾ in.). Like the preceding, but smaller size. Individual specimens sold at listed prices. These total \$114.05. The "collection price" for all the specimens is \$90.00, delivered to any address with pasteboard trays and three No. 2 Oak Chests, as shown in Plate X. Without chests, 10 per cent. less. Mahogany 6-drawer cabinet, \$30.00.

PURCHASE IN PARTS. Free delivery with trays and No. 2

chest. Without chest, 10 per cent. less.

PART I. (Student's School Collection No. 21) 60
names marked with +, totaling \$29.20 ... \$25.00
PART II. 60 names marked with \*, totaling \$33.90 .. 25.00
PART III. 60 remaining names, totaling \$50.95 .... 40.00

## No. 18A. Secondary School Collection

One hundred and twenty museum size specimens, averaging 12 x 9 cm.  $(4\frac{3}{4} \times 3\frac{1}{4})$  in.).

An abridgment of No. 14A. arranged for schools desiring to cut down the specimens to the minimum number required in a brief course. Except in point of numbers, it presents the same attractive and showy appearance as the foregoing, and forms an excellent nucleus about which may be conveniently gathered other important minerals. The Secondary School List is exactly as recommended by Prof. E. S. Dana, and comprises the names marked with + or \* in the following High School List.

Individual museum specimens may be purchased at double the prices listed (for the hand size). The sum of such individual values in the museum size is \$126.20. The "collection price" for all the specimens is \$100.00, delivered to any address. This price includes pasteboard trays (or blocks if requested), and two No. 3 Oak Chests, as shown in Plate X. Without chests, 10 per cent. less.

Purchase in Parts. Free delivery with trays and No. 3 Chest. Without chest, 10 per cent. less.

PART I. (School Collection No. 21A) 60 names marked with +, totaling \$58.40 ......\$50.00 PART II. 60 names marked with \*, totaling \$67.80 ... 50.00

#### No. 18. Student's Secondary School Collection

One hundred and twenty hand size specimens, averaging 10 x 7 cm. (4 x 2¾ in.). Like the preceding, but smaller. Individual specimens sold at listed prices. These total \$63.10. The "collection price" for all the specimens is \$50.00, delivered to any address. This includes pasteboard trays with one No. 3 Oak Chest, as shown in Plate X. Without chest, 10 per cent. less.

Purchase in Parts. Free delivery with trays and No. 2 Chest. Without chest, 10 per cent. less.

PART I. (Student's School Collection No. 21.) 60
names marked with + in High School
List, totaling \$29.20 .....\$25.00

PART II. 60 names marked with \*, totaling \$33.90 . . 25.00

#### No. 21A. School Collection

Sixty museum size specimens, averaging 12 x 9 cm. (4¾ x 3½ in.). This limited selection is not intended for serious study, but more to interest beginners by the beauty of form and color of the specimens and the utility of a few of the popularly known kinds. Excellent for illustrating nature-study talks and elementary work generally. It will also serve as the smallest practicable nucleus essential in beginning a more extensive collection, these first specimens being always worthy of a place beside the later and rarer additions. According to the following "School List," comprising the minerals marked +.

Individual museum specimens may be purchased at double the listed prices (for the hand size). The sum of such individual values in the museum size is \$58.40. The "collection price" for all the specimens, delivered to any address, is \$50.00. This price includes pasteboard trays (or blocks if requested) and one No. 3 Oak Chest, shown in Plate X. Without chest, 10 per cent. less.

#### No. 21. Student's School Collection

Sixty hand size specimens averaging 10 x 7 cm.  $(4 \times 2\frac{3}{4})$  in.). Like the preceding, but smaller. Individual specimens sold at listed prices. These total \$29.20. The "collection price" for all the specimens is \$25.00, delivered to any address. This includes pasteboard trays and one No. 2 Oak Chest, shown in Plate XI. Without chest, 10 per cent. less.

## The High School List

Entire 180 names. Collections 14A and 14.

## The Secondary School List

120 names marked + or \*. Collections 18A and 18.

#### The School List

. 60 names marked +. Collections 21A and 21.

	Carbon. C	
I	DIAMOND. Small octahedral crystal	\$1.50
2+	GRAPHITE, Plumbago or Black Lead. Foliated mass	.40
	Sulphur. S	
3+	SULPHUR. Native, group of brilliant perfect crystals, translucent bright yellow	.75
	Arsenic. As	
4	ARSENIC. Native, fine granular, gray	.75
5	REALGAR. As monosulphide, red	1.00
	ORPIMENT. As trisulphide, foliated, fine yellow ARSENOPYRITE, Mispickel. Fe sulph-arsenide, granular,	1.00
	tin-white	. 20
	Antimony. Sb	
8	Antimony. Native, crystalline, gray	2.50
9+	STIBNITE, Antimony Glance. Sb trisulphide, crystalline,	
	bladed-columnar, steel-gray	.35
	Bismuth. Bi	
10	BISMUTH. Native, crystalline foliated	1.00
	Molybdenum. Mo	
11*	MOLYBDENITE. Mo disulphide, crystallized, tabular, lead-	

### Gold. Au

	GOLD. Native, free grains disseminated in quartz	\$2.00
13	SYLVANITE. Au and Ag telluride, crystallized	2.00
	Platinum. Pt	
14	PLATINUM. Native, grains, steel-gray	1.50
	Silver. Ag	
15+	SILVER, "Leaf Silver." Native, plate	1.50
	ARGENTITE, Silver Glance. Ag sulphide, massive, black.	1.25
17	PYRARGYRITE, Dark Ruby Silver. Ag sulphantimonite	1.00
18	PROUSTITE, Light Ruby Silver. Ag sulpharsenite, dissem-	
	inated	1.25
19		1.25
	Mercury. Hg	
	MERCURY, Quicksilver. Native, globules on matrix	1.00
21+	CINNABAR. Hg sulphide, crystalline, crimson	1.25
	Copper. Cu	
	COPPER. Native, in "Calumet Conglomerate."	. 20
23*	CHALCOCITE, Copper Glance. Cu sulphide, dark steel-	
	gray	1.00
24*	BORNITE, Peacock Ore. Cu and Fe sulphide, coppery bluish-brown, tarnishing iridescent	75
25+	CHALCOPYRITE, Copper Pyrites. Cu and Fe sulphide,	.75
-5	brass-yellow	.35
26+	TETRAHEDRITE, Fahlerz or Gray Copper. Cu sulphan-	
	timonite, massive	1.00
27+	CUPRITE, Chalcotrichite, Ruby Copper. Cu oxide,	
~ O.T	crystallized, capillary	.75
	AZURITE. Cu basic carbonate, crystallized, blue	·75 1.00
	DIOPTASE. Cu basic ortho-silicate, loose crystal, brilliant	2.00
,	emerald-green	1.00
3 I	CHRYSOCOLLA. Cu hydrous silicate, turquois-blue,	
	amorphous	. 50
32	BROCHANTITE. Basic Cu sulphate, brilliant crystals,	
	dark green	.75
	Lead. Pb	
33	LEAD. Native, on matrix	1.00

	ELEMENTARY SYSTEMATIC COLLECTIONS	247
	GALENA. Pb sulphide, cubic cleavage, shining lead-gray.  JAMESONITE, Feather Ore. Pb sulphantimonite, crystal-	\$ .40
36	line granular, steel-gray	1.00
<b>J</b> -	crystallized, splendent blackish-gray	I.25
37+	Pyromorphite. Pb chloride and phosphate, crystals, green.	.75
	MIMETITE. Pb arsenate and chloride, crystals, yellow.	1.50
39*	VANADINITE. Pb vanadate and chloride, crystals, red.	1.00
40	CROCOITE. Pb chromate, prisms, brilliant fine red	1.00
41*	WULFENITE. Pb molybdate, perfect tabular crystals,	
	brilliant orange-red	1.00
	CERUSSITE. Pb carbonate, compact, gray	. 50
43*	Anglesite. Pb sulphate, crystallized, brilliant	1.50
	Tin. Sn	
44+	CASSITERITE, Stream Tin. Sn dioxide, grains, iron-black.	. 50
	Titanium. Ti	
45	ILMENITE. Fe and Ti oxides, iron-black	.25
	RUTILE. Ti dioxide, sharp prismatic crystals, red	.50
47	OCTAHEDRITE, Anatase. Ti dioxide, crystallized, small.	1.50
48	BROOKITE, Arkansite. Ti dioxide, bright black crystals.	.75
	TITANITE, Sphene. Ti calcium titano-silicate, crystal	.50
.,	Radium and Uranium. Ra, U	·
	Highly radio-active minerals.	
50	URANINITE, Pitchblende. Contains Ra, U etc., black	3.00
51	TORBERNITE. U and Cu hydrous phosphate with Ra,	0
	small crystals, green	1.50
52	AUTUNITE. U and Ca hydrous phosphate with Ra,	·
_	small crystals, yellow	1.25
	Iron. Fe	
53	IRON. Native Fe with Ni and Co, Meteoric, etched to	
00	show Widmannstätten crystalline figures	2.00
54+	PYRRHOTITE, Magnetic Pyrites. Fe and Ni sulphide	. 20
55+	PYRITE, Iron Pyrites. Fe sulphide, crystallized, isometric,	
	splendent yellow	. 50
56	massive	.20
	MARCASITE. Fe sulphide, crystallized, orthorhombic	.75
58		
	splendent black	.60
59+	massive granular, red	.20

60 MAGNETITE. Fe protoxide and sesquioxide, octahedral	_
crystals, iron-black	\$ .40
61+ Lodestone. Compact	. 50
62* FRANKLINITE. Fe, Zn and Mn ferrate and manganate	. 40
63* CHROMITE, Chromic Iron. Fe chromate, granular	. 20
64+ LIMONITE, Brown Iron Ore. Fe hydrous sesquioxide	. 20
65+ SIDERITE, Chalybite or Spathic Iron. Fe protocarbonate,	
rhombic cleavage, brown	.20
Nickel. Ni (See also No. 54).	
66 GENTHITE. Hydrous Ni and Mg basic silicate, green	. 50
67* GARNIERITE. Hydrated Ni and Mg silicate, green	. 50
68+ MILLERITE. Ni sulphide, fibrous crystalline, brass-	
yellow	1.00
69* NICCOLITE, Arsenical Nickel. Ni arsenide, reddish-gray.	1.00
Cobalt. Co	
70 LINNÆITE. Co sulphide, small octahedral crystals, gray.	2.00
71 SMALTITE. Co arsenide, compact, gray	1.00
72 COBALTITE, Cobalt Glance. Co sulph-arsenide, crystals	.60
73 ERYTHRITE, Cobalt Bloom. Co hydrous arsenate, red	1.00
Columbium and Tantalum. Cb, Ta	
74* COLUMBITE. Ferrous Fe and Mn columbate and tanta-	
late, crystalline, disseminated in greisen, iron-black.	1.00
Tungsten. Wo	
	75
	·75
,	· <b>7</b> 5
Lithium. Li	
77* SPODUMENE. Li and Al Metasilicate, cleavage, whitish.	.20
78 TRIPHYLITE. Li, Fe and Mn phosphate, brown	.60
79 AMBLYGONITE. Li and Al fluo-phosphate, white	. 50
80* LEPIDOLITE, Lithia Mica. Basic Li, Al and K fluo-silicate,	
micaceous granular, lilac	.20
Manganese. Mn	
81* Pyrolusite. Mn dioxide, crystalline, black	.20
82* MANGANITE. Hydrous Mn sesquioxide, fibrous crystal-	
line, black	1.00
83+ RHODONITE, Fowlerite. Mn and Zn metasilicate, pink,	
crystalline	∙35
84* Rhodochrosite, Dialogite. Mn protocarbonate, cleav-	
able, pink	.75

ELEMENTARY SYSTEMATIC COLLECTIONS	249
Zinc. Zn	
85+ SPHALERITE, Zinc Blende. Zn sulphide, crystallized, resinous	\$ .50
86* ZINCITE. Zn oxide, granular, red	₽ .50 ·75
87* WILLEMITE. Zn orthosilicate, massive, green	.60
88* CALAMINE. Basic Zn silicate, crystallized drusy	. 50
89+ SMITHSONITE. Zn carbonate	.40
Aluminium. Al	•
90+ CORUNDUM. Al sesquioxide, crystallized, gray	. 50
91* BAUXITE. Hydrous Al sesquioxide, yellowish-white	.20
92* SPINEL. Mg aluminate, octahedral crystal	.75
93* CRYOLITE. Al and Na fluoride, semitranslucent white	. 30
94 Turquois. Hydrous basic Al phosphate, blue	· 75
95* WAVELLITE. Hydrous basic Al phosphate, radiated,	40
green	.40
Calcium. Ca	
96 FLUORITE, Fluor Spar. Ca fluoride, cubic crystals, blue	. 50
97+ cleavable-granular, greenish-white	.20
98+ CALCITE, Calc Spar. Ca carbonate, crystallized, scaleno-hedral	. 50
99* Iceland Spar. Doubly refracting rhombic cleavage	1.00
100+ Marble. Crystalline, white	.20
101* Chalk. Amorphous, white	.20
102* Travertine. Columnar-crystalline, indistinctly banded	.60
103+ Aragonite. Ca carbonate, pseudo-hexagonal twin crys-	
tals	. 50
104+ APATITE. Ca phosphate, with Ca fluoride, crystalline,	
green	. 20
105+ Gypsum, Selenite. Hyd. Ca sulphate, cleavage, clear	. 20
106 Alabaster. Compact, translucent white	.20
107* ANHYDRITE. Ca anhydrous sulphate, bluish-gray	. 20
Magnesium. Mg	
108* BRUCITE. Mg hydrate, cleavage, pearly whitish	1.00
109 Magnesite. Mg carbonate, porcelain-like, white	.20
110+ DOLOMITE, Pearl Spar. Ca and Mg carbonate, curved	
rhombs	.30
Boron. B	
111 COLEMANITE. Hydrous Ca borate, crystalline, white	. 50

250	ELEMENTARY SYSTEMATIC COLLECTIONS	
112	BORAX. Hydrous Na borate, crystals	\$ .40
113	BORACITE, Stassfurtite. Mg chloroborate	.30
	Barium. Ba	
114+	BARITE, Barytes or Heavy Spar. Ba sulphate, lamellar,	20
	white	.20
115*	WITHERITE. Ba carbonate, crystalline	. 20
116+	CELESTITE. Sr sulphate, blue cleavage	.20
117*	STRONTIANITE. Sr carbonate, crystalline columnar	. 20
	Sodium. Na	
118+	HALITE, Rock Salt. Na chloride, cleavage, clear	.20
119	SODA NITER. Na nitrate, crystalline, white	.40
	Potassium. K	
120	Sylvite. K chloride, cleavage	.30
	Rare Elements	· ·
121*	ZIRCON. Zr silicate, loose perfect crystals, brown	.40
122		.40
	Silicon. Si	.40
122+	QUARTZ, Rock Crystal. Si dioxide, prism, clear glassy	. 50
124*		.25
125*		.50
126+		.40
127*	Agate. Banded, polished	· 75
128*	Flint. Nodule, gray	. 20
129+	Jasper. Red	.30
130+	OPAL, Precious. Si dioxide with water, play of colors	1.00
131	Fire. Translucent fiery red	·75
132		
	structure well marked, yellowish-brown	.40
	Silicates—The Feldspars	
133+	ORTHOCLASE. Al and K polysilicate, crystals, grayish	. 50
134		
	large crystal, green	. 50
135+	ALBITE, Cleavelandite. Al and Na polysilicate, lamel-	
	lar, white	. 20
	Anorthite. Al and Ca polysilicate, crystallized	1.00
137*	OLIGOCLASE, Sunstone, Aventurine. Al, Na and Ca	
	polysilicate, cleavage, with twinning striæ and internal fiery reflections	
	ternal nery renections	·75

	ELEMENTARY SYSTEMATIC COLLECTIONS	251
138*	LABRADORITE. Al, Na and Ca polysilicate, cleavage, with twinning striæ, chatoyant, bluish-gray  Silcates—Various	\$ .30
120*	Pyroxene, Diopside. Ca and Mg metasilicate, crys-	
-39.	talline, green	. 50
T 40*	Diallage. Ca, Fe and Mg metasilicate, lamellar	.30
141		.50
	Pyroxene, Augite. Ca, Mg, Fe and Al metasilicate,	
	Crystals Enstatite, Bronzite. Mg metasilicate, sublamellar	.40
143 144*	Amphibole, Tremolite. Ca and Mg metasilicate,	.30
	crystalline, whitish	. 40
145*		
_	crystals in talc, green	. 30
	Asbestus. Ca, Fe and Mg metasilicate, fibrous, white	. 20
147+	Hornblende. Ca, Al and Mg metasilicate, cleavable,	
_	black	. 20
-	BERYL. Be and Al metasilicate, green	⋅35
149	GARNET, Grossularite, Cinnamon Stone. Ca and Al ortho- silicate, dodecahedron truncated by trapezohedron,	
150+	bright, brown	.60
130	rical dodecahedron	.30
151+	Muscovite, Potash or Common Mica. Hydrous K	_
	and Al metasilicate, cleavage sheet, gray	. 20
152+	BIOTITE, Magnesium-iron Mica. Mg, Fe, K and Al or-	
152	thosilicate, cleavage sheet, black	. 20
- 33	cate, cleavage sheet, bronze, asteriated	. 20
T = A *	CLINOCHLORE. A hydromica, basic Mg and Al silicate,	.20
<b>-34</b> **	cleavage, green	. 50
155*	CHRYSOLITE, Olivine. Mg and Fe orthosilicate, granu-	. 30
-33.	lar, green	.30
_	WERNERITE, Scapolite. Ca, Al and Na chloro-silicate,	.30
	coarse crystalline granular, pinkish	. 20
	VESUVIANITE. Basic Al and Ca silicate, crystalline	. 40
158+	EPIDOTE. Basic Fe, Al and Ca silicate, crystalline col-	
	umnar, green	. 40
	Zoisite, Thulite. Basic Al and Ca silicate, fine pink	. 40
160*	TOURMALINE. Complex Al, B silicate, black crystals	.40

161 Rubellite. Slender delicate pink crystals in pale lila	С
lepidolite	
162* TOPAZ. Al fluo-silicate, perfect crystals, clear, precious	
163* Andalusite. Al silicate, grayish	
164* CYANITE. Al silicate, crystalline bladed, blue	
165 SILLIMANITE. Al silicate, embedded prisms, gray	•
166 PYROPHYLLITE. Basic Al silicate, radio-fibrous, pearly	•
167+ STAUROLITE. Basic Fe, Al and Mg silicate, twin crystal	
168+ TALC, Steatite. Acid Mg metasilicate, schistose, gray.	
169+ SERPENTINE. Basic Mg silicate, green	
170 KAOLINITE. Basic Al silicate, earthy, white	
171* DATOLITE. Ca and B orthosilicate, glassy crystals	
172+ PREHNITE. Acid Ca and Al orthosilicate, drusy globu	
lar, green	50
173+ APOPHYLLITE. Ca and K silicate, crystallized, pearly	•
transparent whitish	
174* PECTOLITE. Ca and Na metasilicate, radiated, white.	
Silicates—The Zeolites	
175 THOMSONITE. Hydrous Na, Ca and Al silicate	50
176+ NATROLITE. Hydrous Na and Al silicate, radio-fibrous	
white	•
177* ANALCITE. Hydrous Na and Al silicate, crystals, white	
178+ CHABAZITE. Hydrous Na, Ca and Al silicate, cuboic	
rhombs, whitish	
179+ STILBITE. Hydrous Na, Ca and Al silicate, crystallized	
pearly	, 40
180* HEULANDITE. Hydrous Na, Ca and Al silicate, crystal:	
io. Incomment in the state of the call the sincator of state	• • • • •

# PART V

Economic Mineralogy

Mining Sets of Industrial Minerals and Ores

# Economic Mineralogy

## Mining Sets of Industrial Minerals and Ores

#### No. 24A. School of Mines Collection

Four hundred museum size specimens, averaging  $12 \times 9$  cm. (4¾ x 3½ in.). Designed to illustrate as fully as possible the occurrence of the useful minerals. The more striking differences of form are included, as well as important variations in quality of ore, structure, color and mode of occurrence.

The School of Mines List, on the following pages, includes all the common economic minerals, while a few which are rarer and of less immediate commercial interest, are added because valuable if found in marketable quantity.

The commoner species are shown in much wider variety than is possible in shorter collections. Additional varieties and types of the commoner and more important species, will be found in the list of the Complete Type Collection in Part II. The School of Mines Collection will serve every purpose of a high grade reference or working collection for the mining man or prospector, or for the use of mining schools or other institutions offering advanced courses in economic mineralogy. Apart from its utility, it makes an attractive and imposing display, when properly cased, whether in the mining office or public museum.

Individual museum specimens may be purchased at double the hand size prices in the School of Mines List. The sum of these individual values in the museum size is \$1009.20. The "collection price" for all the specimens is \$800.00, delivered to any address. This includes pasteboard trays (or blocks if requested) and two mahogany 10-drawer cabinets. Without cabinets, 10 per cent. less.

Purchase in Parts. Free delivery, with trays (or blocks if requested) and one 10-drawer cabinet with each part. Without cabinets, 10 per cent. less.

#### No. 24. Expert's or Prospector's School of Mines Collection

Four hundred hand size specimens, averaging 10 x 7 cm. (4 x 2¾ in.). Same as preceding, but smaller size. Individual hand specimens may be purchased at the prices in the School of Mines List. These total \$504.60. The "collection price" is \$400.00, delivered to any address. This includes trays and mahogany 10-drawer cabinet. Without cabinet, 10 per cent. less.

Purchase in Parts. Free delivery, with trays.

PART I. 200 specimens in Mining List (Expert's or Prosspector's Mining Collection No. 27), totaling \$202.30, in 10-drawer cabinet.......\$190.00

Without cabinet, \$153.00

Total..... 400.00

Purchase in Sections. Listed as collections Nos. 51 to 65.

## No. 27A. Mining Collection

Two hundred museum size specimens, averaging 12 x 9 cm. (4¾ x 3½ in.). The demand for a reasonably complete series of economic minerals is met by this well arranged collection. As will be seen in the following "Mining List," which is one of the most popular we publish, no attempt is made to represent two varieties of the same mineral, except with very important species. It contains a large proportion of valuable ores, including practically all of those mentioned in the principal mining hand-books. For the work of the prospector or practical man seeking acquaintance with the actual ores themselves, this collection meets every requirement possible within the

limitations of two hundred specimens. Furthermore it makes a very satisfactory showing in the office, laboratory, classroom or public museum.

Individual museum size specimens may be purchased at double the hand size prices given in the Mining List. The sum of such individual values in the museum size is \$404.60. The "collection price" for all the specimens is \$340.00, delivered to any address. This price includes pasteboard trays (or blocks if requested) and 10-drawer cabinet, or four No. 3 oak chests. Without cabinet or chests, 10 per cent. less.

Purchase in Sections. Listed as collections Nos. 52A, 54A, 56A, 58A, 60A, 62A, 64A, and 66A.

#### No. 27. Expert's or Prospector's Mining Collection

Two hundred hand size specimens, averaging 10 x 7 cm. (4 x 2¾ in.). Same as preceding but smaller size. Individual specimens, totaling \$202.30 sold as listed. The "collection price" for all the specimens, with pasteboard trays and 6-drawer cabinet or two No. 3 oak chests, is \$170.00. Without cabinet or chests, 10 per cent. less.

PURCHASE IN SECTIONS. Listed as collections Nos. 52, 54, 56, 58, 60, 62, 64 and 66.

### Sectional Series

### of the School of Mines and Mining Lists

The following collections form successive sections of Nos. 24A, 24, 27A and 27. When sections valued at \$20.00 or over are purchased they are accompanied by the chests mentioned. Without the chests they are 10 per cent. less. If a sufficient number of sections are purchased to fill a drawer cabinet, the latter will be delivered, if requested, instead of chests.

Any fifty-specimen section of the School of Mines List may be purchased in separate parts as shown under Nos. 51A and 51, by first getting a twenty-five specimen section and later completing it by paying the difference between the collection prices of the twenty-five and fifty-specimen sections.

Prices include delivery to any address.

#### Ores of Gold, Silver, Platinum, etc.

No. 51A. Fifty specimens, mostly small, but quality corresponding to the museum size. Total, \$214.00. "Collection price," delivered with trays and No. 3 chest, \$180.00.

Purchase in Parts. Trays and No. 2 chest with each part.

PART I. 25 specimens marked + (No. 52A), \$80.00.

Part II. 25 remaining specimens, \$100.00.

No. 51. Fifty specimens, mostly small, but quality corresponding to the hand size. Total, \$107.00. "Collection price," delivered with trays and No. 2 chest, \$90.00.

Purchase in Parts, at half the price of above 51A parts.

No. 52A. Twenty-five specimens marked +, mostly small, but quality corresponding to the museum size. Total, \$91.50. "Collection price," delivered with trays and No. 2 chest, \$80.00.

No. 52. Twenty-five specimens marked +, mostly small, but quality corresponding to the hand size. Total, \$45.75. "Collection price," delivered with trays and No. 1 chest, \$40.00.

#### Ores of Iron

No. 53A. Fifty museum specimens, averaging 12 x 9 cm. (4¾ x 3½ in.), totaling \$55.40. "Collection price," delivered with trays and No. 3 chest, \$40.00

No. 53. Fifty hand specimens, averaging 10 x 7 cm. (4 x 23/4 in.), totaling \$27.70. "Collection price," delivered with trays

and No. 2 chest, \$20.00.

No. 54A. Twenty-five museum specimens marked +, averaging 12\* x 9 cm. (43/4 x 31/2 in.), totaling \$23.50. "Collection price," delivered with trays and No. 2 chest, \$20.00.

No. 54. Twenty-five hand specimens marked +, averaging 10 x 7 cm. (4 x 23/4 in.), totaling \$11.75. "Collection price,"

delivered with trays, \$10.00.

# Lead, Antimony, Zinc and Cadmium Minerals

No. 55A. Fifty museum specimens, averaging 12 x 9 cm. (4¾ x 3½ in.), totaling \$106.60. "Collection price," delivered with trays and No. 3 chest, \$90.00.

No. 55. Fifty hand specimens, averaging 10 x 7 cm. (4 x 234 in.), totaling \$53.30. "Collection price," delivered with

trays and No. 2 chest, \$45.00.

No. 56A. Twenty-five museum specimens marked +, averaging 12 x 9 cm. (4¾ x 3½ in.), totaling \$49.80. "Collection price," delivered with trays and No. 2 chest, \$40.00.

No. 56. Twenty-five hand specimens marked +, averaging 10 x 7 cm. (4 x 2¾ in.), totaling \$24.90. "Collection price," delivered with trays and No. 1 chest, \$20.00.

## Copper Minerals

No. 57A. Fifty museum specimens, averaging 12 x 9 cm. (43/4 x 3½ in.), totaling \$134.30. "Collection price," delivered with trays and No. 3 chest, \$110.00.

No. 57. Fifty hand specimens, averaging 10 x 7 cm. (4 x 23/4 in.), totaling \$67.15. "Collection price," delivered with

trays and No. 2 chest, \$55.00.

No. 58A. Twenty-five museum specimens marked +, averaging 12 x 9 cm. (4¾ x 3½ in.), totaling \$45.60. "Collection price," delivered with trays and No. 2 chest, \$40.00.

No. 58. Twenty-five hand specimens marked +, averaging 10 x 7 cm. (4 x 23/4 in.), totaling \$22.80. "Collection price," delivered with trays and No. 1 chest, \$20.00.

# Lithium, Barium, Strontium, Sodium, Potassium, Magnesium, Calcium, Boron and Carbon Minerals

No. 59A. Fifty museum specimens, averaging  $12 \times 9$  cm.  $(4\frac{3}{4} \times 3\frac{1}{4})$  in.), totaling \$50.50. "Collection price," delivered with trays and No. 3 chest, \$40.00.

No. 59. Fifty hand specimens, averaging 10 x 7 cm. (4 x 23/4 in.), totaling \$25.25. "Collection price," delivered with

trays and No. 2 chest, \$20.00.

No. 60A. Twenty-five museum specimens marked +, averaging 12 x 9 cm. (4¾ x 3½ in.), totaling \$17.80. "Collection price," delivered with trays, \$15.00.

No. 60. Twenty-five hand specimens marked +, averaging 10 x 7 cm. (4 x 2¾ in.), totaling \$8.90. "Collection price,"

delivered with trays, \$7.50.

## Nickel, Cobalt, Chromium, Manganese and Aluminium Minerals

No. 61A. Fifty museum specimens, averaging 12 x 9 cm. (4¾ x 3½ in.), totaling \$87.20. "Collection price," delivered with trays and No. 3 chest, \$70.00.

No. 61. Fifty hand specimens, averaging 10 x 7 cm. (4 x 2¾ in.), totaling \$43.60. "Collection price," delivered with

trays and No. 2 chest, \$35.00.

No. 62A. Twenty-five museum specimens marked +, averaging 12 x 9 cm. (4¾ x 3½ in.), totaling \$32.80. "Collection price," delivered with trays and No. 2 chest, \$25.00.

No. 62. Twenty-five hand specimens marked +, averaging

No. 62. Twenty-five hand specimens marked +, averaging 10 x 7 cm. (4 x 2¾ in.), totaling \$16.40. "Collection price," delivered with trays, \$12.50.

# Radio-active and Other Rare Element Minerals, including Uranium, Thorium, Yttrium, the Cerium Metals, Zirconium, Germanium and Caesium

No. 63A. Fifty museum specimens, standard of size 12 x 9 cm. (43/4 x 3½ in.), but many are smaller. Total, \$246.60. "Collection price," delivered with trays and No. 3 chest, \$180.00.

No. 63. Fifty hand specimens, standard of size 10 x 7 cm. (4 x 2¾ in.), but many are smaller. Total, \$123.30. "Collection price," delivered with trays and No. 2 chest, \$90.00.

No. 64A. Twenty-five museum specimens marked +, standard of size 12 x 9 cm. (4¾ x 3½ in.), but many are smaller. Total, \$95.80. "Collection price," delivered with trays and No. 2 chest, \$80.00.

No. 64. Twenty-five hand specimens marked +, standard of size 10 x 7 cm. (4 x 2¾ in.), but many are smaller. Total, \$47.90. "Collection price," delivered with trays and No. 1 chest, \$40.00.

# Tin, Tungsten, Titanium, Molybdenum, Vanadium, Tantalum, Columbium, Arsenic, Mercury, Bismuth, Selenium, Tellurium and Sulphur Minerals

No. 65A. Fifty museum specimens averaging 12 x 9 cm. (43/4 x 3½ in.), totaling \$117.60. "Collection price," delivered with trays and No. 3 chest, \$90.00.

No. 65. Fifty hand specimens averaging 10 x 7 cm. (4 x 23/4 in.), totaling \$58.80. "Collection price," delivered with trays and No. 2 chest, \$45.00.

No. 66A. Twenty-five museum specimens marked +, averaging 12 x 9 cm. (43/4 x 3½ in.), totaling \$47.80. "Collection price," delivered with trays and No. 2 chest, \$40.00.

No. 66. Twenty-five hand specimens marked +, averaging

No. 66. Twenty-five hand specimens marked +, averaging 10 x 7 cm. (4 x 2¾ in.), totaling \$23.90. "Collection price," delivered with trays and No. 1 chest, \$20.00.

#### School of Mines List

#### 400 KINDS FORMING ENTIRE LIST.

# Mining List

#### 200 KINDS MARKED +.

The theoretical percentage of valuable element contained is given. Actually, it is often less. Where the amount is not stated, it is a relatively unimportant factor in the commercial value.

#### Nos. 51 and 52. Gold, Silver and Platinum Minerals

#### Gold, Au

- 1+ Gold. Native, crystallized, gold-yellow. 2.00
- 2 arborescent, crystallized. 7.00
- 3 spongiform. 3.00
- 4 filiform, "wire gold." 2.00
- 5 masses or "stringers," disseminated. 2.50
- 6+ grains disseminated in Quartz. 2.00
- 7 ditto, in altered pyrite crystals. 2.00
- 8+ "dust," grains. 1.50
- 9+ nugget. 1.50
- 10+ Electrum. Alloyed with much silver, crystallized, pale yellow. 2.00
- 11 ditto, "leaf gold," crystallized plate, pale yellow. 2.00
- 12 Petzite. Au 25.5, Ag 42., telluride, iron-gray. 3.00
- 13+ Sylvanite. Au 24.5, Ag 13.4, telluride, crystals, silver-white. 2.00
- "Graphic Tellurium," arborescent twinning. 2.00
- 15+ Calaverite. Au 39.5, Ag 3.1, telluride, pale bronze-yellow. 2.50
- 16+ Nagyagite, Foliated Tellurium. Au 8·1, Pb, sulphotelluride, crystalline plates, blackish lead-gray. 3.00

#### Silver, Ag

- 17 Silver. Native, crystallized, silver-white, tarnishing. 3.00
- 18+ "Leaf Silver." Bright crystalline plate. 1.50
- 19+ filiform, wire silver. 1.50
- 20 grains disseminated in matrix. 1.00

- 21 ditto, scales. 2.00
- 22+ Dyscrasite. Ag 78-6, antimonide, crystalline. 2.50
- 23 Argentite, Silver Glance. Ag 87-1, sulphide, crystallized. 2.00
- 24+ massive, sectile, blackish lead-gray. 1.25
- 25 Amalgam. Ag 27.—86. Hg 72.—13. Crystal, silver-white. 4.00
- 26+ Hessite. Ag 63.3, telluride, small crystals, dark-gray. 2.50
- 27+ Galena. 35. to 354. Troy oz. Ag to the ton, argentiferous, Pb sulphide, granular. .75
- 28 Acanthite. Ag 87·1, sulphide, acicular, iron-black. 2.00
- 29 Stromeyerite. Ag 53·1, Cu, sulphide, massive, steel-gray. 2.50
- 30+ Bornite. Argentiferous, granular bluish-brown, tarnishing. .75
- 31 Andorite. Ag 22.5 Sb 41.6 Pb 23.1, sulphantimonite, massive, steel-gray. 4.00
- 32 Pyrargyrite, Dark Ruby Silver. Ag 59.9, sulphantimonite, crystallized, reddish-black. 2.00
- 33+ massive, compact. 1.00
- 34 Proustite, Light Ruby Silver. Ag 65.4, sulpharsenite, crystallized, vermillion. 3.00
- 35+ massive compact, dark red. 1.25
- 36+ Tetrahedrite, Freibergite. 3.—31. Ag, Cu sulphantimonite, granular. 1.00
- 37+ Stephanite, Brittle Silver. Ag 68.5, sulphantimonite, crystallized. 2.00
- 38+ Polybasite. Ag 75.6, Cu sulphantimonite, iron-black. 2.00
- 39 Cerargyrite, Horn Silver. Ag 75.3, chloride, crystallized. 3.00
- 40 massive, highly sectile, grayish. 1.25
- 41+ coating on rock. 1.25
- 42 Embolite. Ag 64.3, chlorobromide, crystallized. 2.00
- 43+ massive, olive-green, darkening on exposure. 1.25
- 44 Iodyrite. Ag 46., iodide, crystals. 1.00
- 45+ massive, sulphur-yellow. 2.00
- 46 Boleite. Ag .15, Pb and Cu oxychloride, cubic crystals, deep blue. .75

## Platinum, Pt; Iridium, Ir and Osmium, Os

- 47 Platinum. Nugget, steel-gray. 4.00
- 48+ minute grains and scales. 1.50
- 49 Sperrylite. Pt 56.5, arsenide minute crystals, tin-white. 2.00
- 50+ Iridosmine. Native Ir 59.83, Os 32.4, Pt .76, grains, tin-white.

#### Nos. 53 and 54. Iron Minerals

- 51 Iron. Meteoric, etched plate, crystalline, steel-gray. 2.00
- 52+ Native, Terrestrial, dark steel-gray, oxidizing. 1.00
- 53+ Pyrite. S 53.4, Fe 46.6, sulphide, cubic crystals. .50
- 54 octahedral crystals. .50
- 55+ pyritohedral crystals, splendent brass-yellow. .50
- 56 penetration or "iron-cross" twins. .75
- 57+ massive. .20
- 58+ ditto, auriferous, 2.5 oz Au per ton. .50
- 59 altered to Limonite, brownish. .50
- 60 Pyrrhotite. Magnetic Pyrites. Fe 61.6, S 38.4, sulphide, granular. .20
- 61+ Marcasite. S 53.4, Fe 46.6, sulphide, crystallized, orthorhombic, Cockscomb Pyrites. .75
- 62 globular. .50
- 63+ Hematite, Specular Iron. Fe 70-, sesquioxide, rhombic crystals splendent black. .60
- 64 Specular Iron. Tabular Crystals. .60
- 65+ Pencil Ore. Columnar diverging. .60
- 66+ compact, red. .20
- 67 ditto with red jasper, Jaspilite. .40
- 68 parting, thick lamellar. .50
- 69+ micaceous, thin foliated. .40
- 70 Kidney Ore, short fibrous, reniform. .60
- 71+ red ocherous, Lenticular Fossil Ore (oölitic). .20
- 72 Martite. Fe 69.9, sesquioxide, octahedral crystals. .75
- 73+ dodecahedral crystals. 1.00
- 74+ Ilmenite. Menaccanite. Fe Ti oxide, compact, iron-black. .25
- 75+ Magnetite. Fe 72.4, protoxide and sesquioxide, octahedral crystals. .40
- 76 dodecahedral crystals, striated, splendent-black. 1.00
- 77+ granular massive, iron-black. .20
- 78 sand. .20
- 79+ Lodestone. Compact. .50
- 80+ Turgite. Fe 66.2, sesquioxide, earthy, red. .20
- 81 Göthite. Fe 62.9, sesquioxide, acicular crystals. 1.00
- 82+ fibrous, concentric radiated reniform. 1.00
- 83 Sammetblende, velvety druse, yellowish-brown. 1.25
- 84+ Limonite, Brown Iron Ore. Fe 59.8, hydrous sesquioxide, compactly fibro-columnar. .40

- 85 globular crust, iridescent bronze. .50
- 86+ mammillary subfibrous, shining black surface. .30
- 87 stalactitic, radio-fibrous. .40
- 88 pisolitic. .40
- 89+ Yellow Ochre. .20
- 90+ Bog Ore, porous. .50
- 91 Brown-clay-iron-stone, massive. .20
- 92+ Xanthosiderite. Fe 57.1, sesquioxide, long fibrous, brown. .75
- 93+ Siderite. Fe 62·1, carbonate, crystallized, obtuse rhombs. .50
- 94 acute rhombs, brown. .75
- 95 Black-band ore, highly carbonaceous. .40
- 96+ cleavable. .20
- 97 granular. .20
- 98 Dufrenite. Fe 45., phosphate, crystalline fibro-columnar. .75
- 99 Melanterite, Green Vitriol. Fe 21.7, sulphate, fibrous, green. .50
- 100 Coquimbite. Fe 19.9, Al, sulphate, bluish-violet. 1.00

# Nos. 55 and 56. Lead, Antimony, Zinc and Cadmium Minerals

#### Lead. Pb

- 101+ Galena, Lead Glance. Pb 86.6, sulphide, cubic crystal, leadgray. .75
- 102 octahedral crystal. 1.00
- 103+ cubic cleavage, bright. .40
- 104 fine granular. .40
- 105+ Jamesonite, Feather Ore. Pb 50.8, Sb 29.5, sulphantimonite, crystalline granular, steel-gray. 1.00
- 106 capillary, matted. 1.25
- 107+ Massicot, Yellow Plumbic Ochre. Pb 92.8, oxide, earthy, orpiment-yellow. 2.00
- 108+ Cerussite. Pb 76.8, carbonate, crystallized aggregate, satiny white. 1.25
- 109 reticulated twinning. 2.00
- 110+ massive, gray. .50
- 111 Phosgenite. Pb 76., chlorocarbonate, prismatic crystal. 1.25
- 112+ Pyromorphite. Pb 78.4, chlorophosphate, crystallized, green. .75
- 113 brown crystals. 1.00
- 114+ Anglesite. Pb 68-3, sulphate brilliant crystals. 1.50

- 115 compact massive, dull gray. 1.50
- 116 Crocoite. Pb 64., Cr 16.1, chromate, prismatic crystals, brilliant crimson. 1.00

#### Antimony, Sb

- 117 Allemontite. Sb 34.8, As 65.2, alloy, crystalline, tin-white, tarnishing. 3.00
- 118+ Antimony. Native, granular crystalline, tin-white. 2.00
- 119+ Stibnite, Antimony Glance. Sb 71.4, sulphide, crystals. 1.00
- 120+ crystalline, columnar bladed, lead-gray. .35
- 121 crystalline granular. .35
- partially oxidized crystals, yellow. 1.00
- 123+ Zinkenite. Sb 41-8, Pb 35-9, sulphantimonite, fibrous. 2.00
- 124+ Berthierite. Sb 56.6, Fe sulphantimonite, crystalline fibrous, steel-gray. 1.00
- 125 Senarmontite. Sb 83.3, trioxide, small octahedrons. .75
- 126+ Cervantite, Antimony Ochre. Sb 78.9, oxide, massive, yellowish-white. .40
- 127 Bindheimite. Sb 22.6, Pb 58.5, lead antimonate, pulverulent coating, yellow. 1.00
- 128 Nadorite. Sb 30·5, Pb 52·4, chlorantimonate, yellow and brown.

#### Zinc, Zn

- 129+ Sphalerite, Zinc Blende. Zn 67., sulphide, crystallized, brownish. .50
- "Ruby Blende," crystals, bright, transparent. .75
- 131+ "Black Jack," crystals, glistening. .50
- 132 dodecahedral cleavage. .75
- 133+ granular cleavable, resinous. .20
- 134 Christophite, granular cleavable, black. .40
- 135+ Wurtzite, Schalenblende. Zn 67, sulphide, fibrous, brown. 1.00
- 136+ Zincite. Zn 80·3, oxide, crystalline, red, with franklinite. .75
- 137 Franklinite. Zn 11-9, Fe 30-8 and Mn oxide, octahedral crystal, black. 1.25
- 138+ massive granular, coarse. .40
- 139 Chalcophanite. Zn 21·1, Mn 46·2, oxide, botryoidal subfibrous. .75
- 140+ Smithsonite. Zn 52, carbonate, botryoidal, massive. .40
- earthy, impure, "dry-bone," grayish. .40
- 142+ Aurichalcite. Zn 42.6, Cu. 16.8, carbonate, microscopic crystals forming capillary velvety crust, turquois-blue. .75

- 143 Hydrozincite. Zn 60-1, carbonate, reniform fibrous crust. 1.50
- 144 Willemite. Zn 42., orthosilicate, crystallized, flesh-red. 1.50
- 145+ massive, apple-green, with franklinite. .60
- 146 Calamine. Zn 54·1, silicate, tabular crystals, grouped. 1.00
- 147+ crystalline mass. .50
- 148 Adamite. Zn 45.3, arsenate, crystallized, light yellow. 1.00

#### Cadmium, Cd

- 149+ Greenockite, Cadmium Blende. Cd 77.7, sulphide, coating on ore. 1.50
- 150 Smithsonite. Cadmiferous, "turkey-fat ore," yellow. 1.50

#### Nos. 57 and 58. Copper Minerals

- 151 Copper. Native, tetrahexahedral crystals. 1.00
- 152+ crystallized, arborescent. .50
- 153 plates or "leaf copper." .50
- 154+ massive. .75
- 155+ disseminated in conglomerate. .20
- 156+ Domeykite. Cu 76-1, arsenide, compact, iridescent-bronze. 1.25
- 157 argentiferous, granular. 2.50
- 158 Algodonite. Cu 83.5, arsenide, silver-white, tarnishing. 3.00
- 159 Whitneyite. Cu 88.4, arsenide, reddish-white, tarnishing. 5.0
- 160 Chalcocite, Copper Glance. Cu 79.8, sulphide. Redruthite, crystallized. 1.50
- 161+ compact, bright iron-black. 1.00
- 162 Covellite, Indigo Copper. Cu 66.4, sulphide, crystallized, thin hexagonal tables, indigo-blue. 3.00
- 163+ foliated, crystalline, bright. 2.00
- 164+ platiniferous, enclosing sperrylite, porous, dull. 2.00
- 165 Bornite, Peacock Ore. Cu 55.5, Fe, sulphide, crystallized. 2.50
- 166+ compact massive, bluish-coppery-brown, iridescent. .75
- 167+ Chalcopyrite, Copper Pyrites. Cu 34.5 and Fe sulphide, small tetrahedrons on pearl spar. .50
- 168 twin crystals. 1.00
- 169 reniform. 1.25
- 170+ massive compact, brass-yellow. .35
- 171 massive granular. .35
- 172 Tetrahedrite, Fahlerz or Gray Copper. Cu 52·1, Sb 24·8, sulphantimonite, perfect tetrahedrons, iron-black. 1.00
- 173+ massive compact, grayish iron-black. 1.00

- 174 Enargite. Cu 48-3, sulpharsenite, crystallized, black. 2.00
- 175+ cleavable granular. 1.00
- 176+ Atacamite. Cu 59.4, chloride, crystallized, emerald-green. 1.50
- 177 granular massive. 1.00
- 178 Cuprite, Ruby Copper. Cu 88-8, oxide, octahedral crytals. 2.00
- 179+ Chalcotrichite, capillary, ruby-red. .75
- 180+ massive compact, dark red. 1.50
- 181 partly altered to malachite, crystal, green. 1.50
- 182+ Tenorite, Melaconite. Cu 78.8, oxide, massive, dull black. 1.00
- 183+ Malachite. Cu 58.4, carbonate, capillary, green. .75
- 184 tuberose, compact. 2.00
- 185+ massive, bright green. 1.25
- 186+ Azurite. Cu 56-, carbonate, crystallized, deep blue. 1.00
- 187 tuberose, concentric, azure-blue. 1.25
- 188 altered to malachite, crystallized, green. 1.00
- 189+ massive. .75
- 190+ Chrysocolla. Cu 36.6, silicate, compact, turquois-blue. .50
- 191 Olivenite. Cu 38.8, arsenate, crystallized, olive-green. 1.50
- 192 Pseudomalachite. Cu 53·3, phosphate, radio-fibrous, dark green. 1.50
- 193+ Tyrolite. Cu 40.6, As 17.8, arsenate, foliated, green. .75
- 194+ Conichalcite. Cu 24.2, As 26.6 and Ca, arsenate, globular, green. 1.00
- 195+ Brochantite. Cu 56.2, sulphate, crystallized, dark green. .75
- 196 fibrous, green. 1.00
- 197 altered to cuprite (red oxide). 1.50
- 198 Kröhnkite. Cu 18-3 and Na, sulphate, fine blue. 2.00
- 199+ Chalcanthite. Cu 25.8, sulphate, deep blue. .50
- 200 Natrochalcite. Cu 33.4 and Na sulphate, pyramidal crystals, brilliant emerald-green. 3.00

# Nos. 59 and 60. Lithium, Barium, Strontium, Sodium, Potassium, Magnesium, Calcium, Boron and Carbon Minerals

#### Lithium, Li

- 201 Spodumene. Li 2., Al, silicate, cleavage, whitish. .20
- 202+ Lepidolite, Lithia Mica. Li 1.2, K, fluo-silicate, crystalline granular, lilac. .20
- 203 Triphyllite. Li 2.2, Fe and Mn phosphate, massive, brown. .60
- 204+ Amblygonite. Li 2.35, Al, fluo-phosphate massive, white. .50

#### Barium, Ba

- 205+ Witherite. Ba 68.9, carbonate, grayish-white. .20
- 206 Barite, Barytes or Heavy Spar. Ba 58.9, sulphate, crystals. .50
- 207+ massive lamellar, white. .20

#### Strontium, Sr

- 208+ Strontianite. Sr 59.3, carbonate, columnar, whitish. .20
- 209 Celestite. Sr 47.3, sulphate, bright clear crystals. 1.00
- 210+ cleavage, pale bluish. .20

#### Sodium, Na and Potassium K

- 211 Halite, Rock Salt. Na chloride, cubo-octahedral crystals, clear colorless. .50
- 212+ cubic cleavage. .20
- 213 Trona. Na acid carbonate, fibrous. .40
- 214+ Soda Nitre, Chili Saltpetre. N 16.4, Na nitrate, crystalline granular, white. .40
- 215 Thenardite. Na sulphate, crystallized, yellowish. .40
- 216+ Sylvite. K 52.4, chloride, cleavage. .20
- 217 Orthoclase, Potash Feldspar, K 6.6 and Al, silicate, crystal, grayish. .50
- 218+ Microcline, Potash Feldspar. K 6.6, Al, silicate, cleavage. .20
- 219 Muscovite, Potash Mica. K and Al silicate, cleavage sheet, 20

#### Calcium, Ca

- 220 Fluorite, Fluor Spar. F 48-9, Ca 51-1, cubic crystals, transparent. .50
- 221+ granular cleavable, greenish. .20
- 222 Calcite, Iceland Spar. CaO 56., carbonate, transparent doubly refracting. 1.00
- 223+ Marble, crystalline, white. .20
- Mexican Onyx, variegated bands, translucent. .60
- 225+ Hydraulic Limestone, Cement Rock, shaly, blackish. .30
- 226 Apatite. P 23.4, Ca, phosphate, large crystal, brown. .50
- 227+ granular, green. .20
- 228 Phosphate Rock, fossiliferous. .20
- 229 Gypsum, Selenite. Ca sulphate, large crystal. .75
- 230 Selenite, transparent cleavage. .20
- 231+ Alabaster, fine granular, white. .20
- 232 Dolomite. CO<sub>2</sub> 47.8, MgO 21.7 and Ca, carbonate, granular, white. .20

#### Magnesium, Mg

- 233+ Magnesite. MgO 47.6, CO<sub>2</sub> 52.4, carbonate, compact porcelainlike, white. .20
- 234+ Serpentine, Asbestus. Mg silicate, silky fibrous. .40
- 235 massive, green. .20
- 236+ Talc, Soapstone or Steatite. Mg silicate, schistose, grayish. .20
- 237+ Kieserite. Mg 17.4, sulphate. .40
- 238 Blödite. Mg 7.2, and Na, sulphate, crystallized. .50

#### Boron, B

- 239+ Boracite, Stassfurtite. B 11-8, Mg 18-8, chloroborate, massive.
- 240 Colemanite. B 16-1, Ca, borate, crystallized, white. .50
- 241 Borax. B 5.7, Na, borate, crystal. .40

#### Carbon, C

- 242+ Diamond. Pure C, small crystal (in tube), with large specimen of matrix, Kimberly "blue earth." 2.50
- 243 Carbonado, granular, black, small. 5.00
- 244+ Graphite, Black Lead or Plumbago. Pure C, foliated mass. .40
- 245+ PETROLEUM, Mineral Oil. Hydrocarbon. .20
- 246 ASPHALTUM, Wurtzilite, Mineral Pitch or Bitumen. Hydrocarbon, velvety black. .20
- 247+ MINERAL COAL, Anthracite or Hard Coal. .20
- 248 COPALITE. Congo Gum. Clear light yellow. .40

#### Silicon, Si

- 249+ Quartz, Rock Crystal. Si dioxide, clear colorless. .50
- 250 Opal, Tripolite. Infusorial Earth, Si dioxide, white. .40

# Nos. 61 and 62. Nickel, Cobalt, Chromium, Manganese and Aluminium Minerals.

#### Nickel, Ni

- 251 Josephinite. Ni 30.45, alloyed with Fe, pebbles. 1.00
- 252+ Pentlandite. Ni 22. and Fe, sulphide, cleavages in pyrrhotite, bronze-yellow, tarnishing. 2.00
- 253+ Niccolite, Arsenical Nickel. Ni 43·1, As 56·1, arsenide, massive, reddish-gray. 1.00
- 254 Millerite. Ni 64.7, sulphide, acicular crystals. 2.00
- 255+ fibrous crust, brass-yellow. 1.00

- 256 Breithauptite. Ni 32.8, Sb 67.2, antimonide, massive, violet copper-red. 1.50
- 257+ Pyrrhotite, Magnetic Pyrites. Fe sulphide with Ni 1.8—4.6, compact, bronze-yellow, tarnishing. .20
- 258 Polydymite. Ni 59.4 and Fe, sulphide, cubic cleavage, steelgray. 3.00
- 259+ Gersdorffite, Nickel Glance. Ni 34.5, As 45.3, sulph-arsenide, massive granular. 1.50
- 260+ Ullmannite. Ni 28-8, Sb 57-, sulphantimonide, massive granular, steel-gray. 1.00
- 261 Rammelsbergite. Ni 28·1, As 71·9, arsenide, massive, reddish tin-white. 1.00
- 262 Zaratite, Emerald Nickel. Ni 46.7, carbonate, massive. .60
- 263 Genthite. Ni 22.4, silicate, massive, bright green. .50
- 264+ Garnierite. Ni 20.7, silicate, massive, bright apple-green. .50
- 265 Annabergite. Ni 24., arsenate, massive, apple-green. 1.00

#### Cobalt, Co

- 266+ Linnaeite. Co 75.9, sulphide, massive, steel-gray. 1.50
- 267 Smaltite. Co 28.2, As 71.8, arsenide, cubo-octahedrons. 1.50
- 268+ massive, steel-gray. 1.00
- 269 Chloanthite. Co 28.8, arsenide, massive, steel-gray. 1.25
- 270 Cobaltite, Cobalt Glance. Co 35.5, As 45.2, sulph-arsenide, crystals. 1.00
- 271+ massive, reddish-gray. .60
- 272 Glaucodot. Co 23.8, Fe, sulpharsenide, large crystal. 1.25
- 273+ massive crystalline, tin-white. 1.00
- 274+ Asbolite, Earthy Cobalt. Co 1.—23.5 and Mg, oxide. .30
- 275 Erythrite, Cobalt Bloom. Co 28.7, As 25., arsenate, red. 1.00

#### Chromium, Cr

276+ Chromite, Chromic Iron. Cr 46.5 and Fe oxide, massive ironblack. .20

#### Manganese Mn

- 277+ Alabandite. Mn 63·1, sulphide, crystalline, blackish. 1.00
- 278 Hausmannite. Mn 84-1, oxide, massive, iron-black. .75
- 279 Braunite. Mn 65.2, oxide and silicate, massive. 1.00
- 280+ Pyrolusite. Mn 63.4, dioxide, crystalline granular, black. .20
- 281 radio-fibrous. .75
- 282 Manganite. Mn 62.4, sesquioxide, crystallized. 1.50
- 283+ massive fibrous. 1.00

- 284+ Psilomelane. Mn 40.5, manganate, compact, black. .20
- 285+ Bog Manganese, Black Wad. Mn 1.—19., impure oxide, earthy. .20
- 286 Rhodochrosite, Dialogite. Mn 47.8, carbonate, cleavable granular, light pink. .75
- 287 Rhodonite, Fowlerite. Mn 47.4 and Zn, silicate, cleavage, pink. .35

#### Aluminium, Al

- 288 Cryolite. Al 12.8, Na 32.8, fluoride, cuboid prismatic crystals, clear colorless. 1.50
- 289+ massive, translucent white. .30
- 290 Corundum. Al oxide, gray, crystals altered on surface. .50
- 291+ broad cleavage, bronze-gray. .50
- 292 Ruby, gem-sand, clear red. 1.00
- 293 Sapphire, broken crystals, deep blue. 1.00
- 294+ Emery, black, granular. .20
- 295+ Bauxite. Al 39.6, hyd. oxide, yellowish, earthy. .20
- 296+ Garnet, Almandite. Al and Fe silicate, large dodecahedral crystal. .30
- 297+ Kaolinite. Al silicate, earthy, white. .20
- 298 Pyrophyllite. Al silicate, radiated. .75
- 299 Alunogen. Al 3.97, sulphate, silky fibrous, whitish. .75
- 300+ Alunite. Al 9.8, K, sulphate. .30

# Nos. 63 and 64. Radio-active and Other Rare Element Minerals, including Uranium, Thorium, Yttrium, Cerium metals, Zirconium, Beryllium, Germanium and Caesium.

#### Radium, Ra and Uranium, U

- 301+ Uranophane. U 58., with Ra, He, etc., hydrous silicate. 2.00
  302+ Fergusonite. U 3.4, Y 18.2, Cb 28.3, Ta 8.6, U and Y columbate and tantalate with Ra, etc., pyramidal crystals, dull grayish-brown. 1.50
- 303 massive, brilliant vitreous brownish-black. 1.50
- 304 Sipylite. U 3. Cb 28.8, columbate with Ra, etc., brownish-black. 3.00
- 305+ Samarskite. U 11.7, Y 8.8, Cb 35., Ta 13.1, U and Y columbate and tantalate, with Ra, etc., massive, splendent velvet-black. 2.50

- 306 Annerodite. U 14.5, Cb 16.5, U, Y, etc., pyrocolumbate, massive, black. 4.00
- 307+ Euxenite. U 7.2, Y 14, Cb 24., Ti 12.9, U and Y columbate and titanate, with Ra, etc., massive, vitreous black. 1.50
- 308+ Torbernite, Copper-uranite. U 53·1, Cu 8·4, phosphate with Ra, green. 1.50
- 309+ Autunite, Lime-uranite. U 53.6 and Ca, phosphate with Ra, yellow. 1.25
- 310+ Uraninite, Pitchblende. U 71., Uranate of Uranyl with Ra, etc., massive. 3.00
- 311+ Bröggerite. U 70., cubo-octahedral crystals, dull black. 3.00
- 312 Cleveite. U 55., with Ra, He, etc., black. 3.00
- 313 Thorogummite. U 17., Th 39.5, silicate, rough prisms, yellowish-brown. 2.00
- 314 Gummite. U 55.7, resinous yellow. 2.00
- 315 Carnotite. U 54.8, V 10.2, Ra, etc., compact, yellow. 4.00
- 316+ disseminated in sandstone. 1.00

#### Thorium, Th

- 317 Tritomite. Th 7.4, Ce metals 47.9, massive, resinous dark brown. 5.00
- 318 Thorite. Th 65.2, etc., silicate, crystal, brownish-black. 2.50
- 319+ massive. 2.50
- 320+ Orangite, massive, brownish-yellow. 3.00
- 321 Yttrialite. Th 10.5, Y 36.8, silicate, massive, vitreous greenish-black. 8.00
- 322 Pyrochlore. Th 7., columbate of Ce metals, octahedral crystal, brown. 1.25
- 323+ Æschynite. Th 12.5, Ce 14.2, Cb 12.5, thorate niobate and titanate of Ce metals, massive, brownish-black. 1.50
- 324 Polymignite. Th 3., columbate and titanate (zirconate) of Ce metals, crystallized, black. 6.00
- 325+ Monazite. Sand, Ce 24·1, phosphate of Ce metals with 1·—6· Th, yellowish-brown. .40
- 326+ Thorianite. Th 60-9, Ce 6-2, U 10-2, with He, etc., oxide, cubic crystals, iron-black. 2.50

#### Yttrium, Y and Cerium, Ce Metals

- 327 Yttrocerite. Y 14.5, Ce 4.26, Ca, fluoride, violet-blue. .75
- 328+ Gadolinite. Y 40.4, silicate of Ce and Y metals, large coarse crystal. 5.00
- 329 massive, vitreous black. 2.50

- 330+ Thalenite. Y 51.6, silicate, massive, flesh-red. 3.00
- 331 Yttrotantalite. Y 18.5, Ta 18.7, tantalate and columbate, crystallized. 3.00
- 332 Hielmite. Y 31.7, Cb 6.6, Ta 51.3, Y, etc., stanno-tantalate and columbate, crystallized, black. 2.00
- 333+ Xenotime. Y 47.8, Ce metals, phosphate, pyramids. 2.00
- 334 massive, dull brown. 1.50
- 335 **Tysonite**. Ce 40·1, fluoride of Ce metals, massive, yellowish. 3.00
- 336+ Fluocerite. Ce 39.53, fluoride of Ce metals, yellowish. 1.50
- 337 Parisite. Ce 37.7, fluocarbonate of Ce metals, crystallized, brownish-yellow. 6.00
- 338 Bastnäsite. Ce 28-9, fluo-carbonate of Ce metals, massive, brown. 4.00
- 339+ Allanite. Ce 13.8, Ce metals, etc., silicate, massive, black. .50
- 340+ Cerite. Ce 30.8, silicate of Ce metals, etc., massive, purplishgray. 1.50
- 341+ Monazite. Ce 22·1, phosphate of Ce metals, etc., broken crystals, dull brown. 1.00

#### Zirconium, Zr and Beryllium, Be

- 342 Baddeleyite. Zr 70.4, oxide, fibrous globular, greenish. 4.00
- 343+ Zircon. Zr 49.7, silicate, crystals, brown. .40
- 344 Hyacinth, water-worn crystals, transparent red. .40
- 345 CYRTOLITE. Zr 35.5, Ce metals, silicate, crystals, brown. 1.00
- 346+ Beryl. Be 38.4, Al, silicate, massive, brownish-yellow. .35
- 347 Phenacite. Be 16.1, silicate, broken crystals, white. 4.00
- 348 Beryllonite. Be 72.7, Na, phosphate, transparent. 1.00

#### Germanium, Ge and Caesium, Cs

- 349+ Argyrodite. Ge 6.9, Ag 74.7, sulphide, massive, gray. 2.50 350+ Pollucite. Cs 28.5 Al, silicate, massive, glassy white. 3.00
- No. 43. Tin, Tungsten, Titanium, Molybdenum, Vanadium,

# Tantalum, Columbium, Arsenic, Mercury, Bismuth,

# Selenium, Tellurium and Sulphur Minerals Tin, Sn

- 351 Franckeite. Sn 12·3, Pb 50·5, sulphostannide, radio-foliate, blackish-gray. 1.50
- 352+ Cylindrite. Sn 26.3, Pb 35.4, sulphostannide, cylindrically foliated, lead-gray. 1.25

- 353+ Cassiterite, Tin Stone. Sn 78.6, oxide, prismatic crystals. 1.50
- 354 twin crystals, splendent brown. 2.50
- 355+ massive, dark brown. 1.00
- 356+ Stream Tin, water-worn grains. .50
- 357 disseminated in gangue. .30
- 358+ Stannite, Tin Pyrites. Sn 27.5, Cu 29.5, sulphide, massive greenish-iron-black. .75

#### Tungsten, W

- 359+ Wolframite. W 67-1, Fe tungstate with Mn, crystallized. 1.00
- 360 bladed crystalline, bright iron-black. .75
- 361+ Hubnerite. W 60.7, Mn, tungstate with Fe, bladed crystals, brown. 1.00
- 362 Scheelite. W 72., Ca tungstate, crystallized. 1.50
- 363+ massive, whitish. .75

#### Titanium, Ti

- 364+ Ilmenite, Menaccanite. Ti 36·1, Fe 36·8, oxide, black. .25
- 365+ Rutile. Ti 60-, oxide. Prismatic crystals, red. .50
- Nigrine, with Fe as impurity, crystallized, black. .50

#### Molybdenum, Mo

- 367+ Molybdenite. Mo 60-, sulphide, crystallized, lead-gray. .40
- 368 cleavages, loose. .75
- 369 Molybdite. Mo 65.6, Fe, oxide, pulverulent, yellow. 1.00
- 370 Wulfenite. Mo 25.9, Pb 56.2, molybdate, crystal aggregate vellow. 1.00
- 371+ tabular crystals, bright orange-red. 1.00

#### Vanadium, V

- 372 Roscoelite. V 14., silicate, small scales, dark brown. 3.00
- 373 Descloizite. V 12.7, Pb 51.3, vanadate, crystalline, brownish-red. 1.00
- 374 Endlichite. V 9.9, Pb 67.4, chloro-vanadate, massive, orange. 1.50
- 375+ Vanadinite. V 9.9, Pb 67.4, chloro-vanadate, crystallized, red.

#### Tantalum, Ta and Columbium, Cb

- 376 Columbite. Cb 59.9, Fe, columbate (and tantalate), crystallized. 2.00
- 377+ massive, iron-black. 1.00
- 378 Tantalite. Ta 69.9, Fe and Mn tantalate (and columbate), iron-black. 1.50

379+ Manganotantalite, massive, brownish-black. 1.50

380 Stibiotantalite. Ta 21·1, Sb 16·7, tantalate, water-worn pebbles, yellowish. 2.50

#### Arsenic, As

381 Arsenic. Native, spherical crystal aggregates. .75

382+ massive, fine granular, tin-white, tarnishing. .75

383 Realgar. As 70-1, monosulphide, red, crystallized. 1.25

384+ massive compact, light red. 1.00

385+ Orpiment. As 61., trisulphide, foliated mass, yellow. 1.00

386 Arsenopyrite, Mispickel. As 46., Fe 34.4, sulph-arsenide crystallized. 1.00

387+ massive, silver-white. .25

388 Löllingite, Leucopyrite. As 59.9, Fe, sulpharsenide, massive, tin-white. .35

#### Mercury, Hg

389+ Mercury. Native, minute tin-white globules in gangue. 1.00

390 Metacinnabarite. Hg 86.2, sulphide, disseminated masses, black. 1.00

391+ Cinnabar. Hg 86.2, sulphide, crystallized. 1.25

392 massive, fine granular, cochineal-red. 2.50

393 Livingstonite. Hg 24.8, Sb 53.1, sulphantimonite, columnar massive, blackish lead-gray. 2.50

Bismuth, Bi; Tellurium, Te; Selenium, Se and Sulphur, S

- 394+ Bismuth. Native, crystalline disseminated, reddish-silverwhite, tarnishing. 1.00
- 395 Bismuthinite, Bismuth Glance. Bi 81.2, sulphide, crystalline, lead-gray. 1.00
- 396 Emplectite. Bi 62., Cu 18.9, sulphobismuthite, crystallized, grayish. 1.25
- 397+ Tetradymite. Te 33.—49., Bi 67.—51., foliated, steel-gray. 1.50

398+ Guanajuatite. Se 36·3, Bi 63·7, selenide, bluish-gray. 2.00

399 Clausthalite. Se 27.7, Pb 72.3, selenide, massive, gray. 2.00

400+ Sulphur. Native, crystallized, yellow. .75

# PART VI

Crystallography

Crystals for Measurement and Study

# Crystallography

## Loose Crystals for Measurement and Study

#### Advanced Collections

In the five years since the first publication of the Complete Crystal List, our advanced collections of crystals have met with a wider acceptance than was anticipated. A number of prominent teachers of crystallography, well known as writers on the subject, after examining in detail the Complete Crystal Collection, expressed surprise at finding such a unique and excellent series on sale.

While reduced in price, the advanced collections are superior to those originally distributed by us, both in the planning of the list and in the quality of material furnished. The arrangement and definitions in Dana's "Text-book of Mineralogy" have been carefully followed, making the sets especially valuable to those using this work or Penfield's "Determinative Mineralogy," most of the crystal forms described therein being · included in the collection. An arrangement according to any other author, will, on request, be prepared without extra cost. The aim has been to accurately represent as large a number of forms as possible. A duplication of any combination has been avoided, even though occurring in different minerals. Variety of form is the primary object, while as many species and crystal groups have been introduced as was practicable. thirty-two possible groups in the six systems, only twenty-three are known in nature. Of these, every one is represented. revising the list, the Miller symbols have been added.

The individual crystals selected are the best our extensive facilities afford, and have been measured where necessary. They range generally from I to 4 cm. in length, and nearly all are sufficiently sharp and bright for the reflecting goniometer. The majority are large enough for contact measurement.

The mahogany cabinets holding the crystals are made according to our own designs, especially for these collections. The

4 x 3 cm. white glazed pasteboard trays display the crystals in an excellent manner. The crystals in each collection are numbered to correspond to the following list, besides having on each tray one of our small printed labels, giving name, composition and locality, as shown in Plate IV.

## No. 73A. Complete Crystal Collection

Three hundred measurable crystals. As described above and in the Complete Crystal list, this set evenly covers the whole field of crystallography. Many of these collections have been sold at the former price of \$150.00. The total value of the crystals is \$153.85, and the present "collection price," delivered to any address, with trays in cabinet, is \$120.00.

Without cabinet, 10 per cent. less.

Purchase in Parts. Delivered to any address, with trays. PART I. 150 Crystals marked \* (No. 75A), with 300 trays, in drawer cabinet, \$57.00.

(Part I, without cabinet, \$45.00).

PART II. 150 remaining crystals without cabinet, \$63.00. Purchase in Sections. Delivered to any address:

Section A. 200 Simple Crystals numbered 1—200, totaling

\$110.30, with 300 trays, in drawer cabinet, \$90.00. Without cabinet, \$78.00.

Section B. 50 Twin Crystals numbered 201—250, totaling \$26.95, without cabinet, \$18.00. Cabinet \$5.00 extra.

Section C. 50 specimens Illustrating Irregularities of Crystals and Pseudomorphs, numbered 251-300, totaling \$16.60, without cabinet, \$12.00. Cabinet \$1.50 extra.

# No. 75A. Abridged Crystal Collection

One hundred and fifty measurable crystals, marked \*, comprising the Abridged Crystal List. A careful elimination of rare and less important forms is here effected. With trays, in mahogany cabinet similar to that in Plate VIII. The total value of the crystals is \$63.75 and the "collection price," delivered to any address, is \$50.00. Without cabinet, 10 per cent. less.

# Complete Crystal Collection

No. 73A. 300 Numbers Comprising Entire List

# Abridged Crystal Collection

No. 75A. 150 Numbers Marked \*
The Figures Mentioned Are in Part II.

## I. Isometric System

The forms in this system can be referred to three axes, which are at right angles to one another and of equal lengths.

	Normal Group—Galena Type
I*	Cube a (100)
2*	Cube a (100)       Galena .30         Octahedron o (111)       Magnetite .20
2*	Dodecahedron $d$ (110)
4*	Tetrahexahedron $e$ (210) modifying cube $a$ (100),
•	fig. 523
5*	fig. 523
6	Hexoctahedron $t$ (421) modifying cube $a$ (100), fig.
	521
7*	Cube a (100) modified by octahedron o (111)Galena .30
8*	Cube $a$ (100) modified by trapezohedron $m$ (311) Fluorite 1.50
9	Octahedron $o$ (111) modified by cube $a$ (100)Galena .50
10*	Octahedron o (111) modified by dodecahedron d
	(110)Franklinite .75 Octahedron o (111) modified by dodecahedron d
11*	Octahedron $o$ (111) modified by dodecahedron $d$
	(110) and trapezohedron $m$ (311)Microlite 1.00
12	Octahedron o (111) modified by dodecahedron d
	(110), trapezohedron $m$ (311) and cube $a$ (100), similar to fig. 2141
	similar to fig. 2141
13*	Dodecahedron $d$ (110) modified by cube $a$ (100) Fluorite .75
14	Dodecahedron $d$ (110) modified by octahedron $o$
	(111) Cuprite 1.25
15*	Dodecahedron d (110) modified by trapezohedron
	n (211), fig. 1578
16	Trapezohedron n (211) modified by dodecahedron
	d (110), fig. 1580
	Pyritohedral Group—Pyrite Type
1/4	Pyritohedron e (210), fig. 289
	Cube a (100), fig. 290
19*	Octahedron o (111)
20	Pyritohedron e (210) modified by cube a (100), fig.
	295Pyrite .30

21*	Pyritohedron e (210) modified by octahedron e (111),
22	fig. 297
	octahedron o (111)
23	(111) and diploid s (321) Purity 75
24*	(111) and diploid s (321)
25*	Cube a (100) modified by pyritohedron e (210) and
-6	octahedron o (111)
26 27	Octahedron o (111) modified by pyritohedron e
-,	(210), fig. 294
28	(210), fig. 294
	(221)
29*	
	Tetrahedral Group—Tetrahedrite Type
30*	Tetrahedron o (111) modified by dodecahedron d
	(110) and tristetrahedron $n$ (211), fig. 434 Tetrahedrite
31	Tetrahedron $o$ (111) modified by cube $a$ (100) Boracite .50
32*	Cube $a$ (100) modified by tetrahedron $o$ (111) and
4	dodecahedron d (110), fig. 2387
33₹	Tetrahedron plus o (111) and minus o <sub>1</sub> , (111), tetrahedral symmetryZunyite .20
	Gyroidal or Plagihedral Group—Cuprite Type
a 4 th	Trapezohedral symmetrySal-ammoniac
34*	.40
	Tetartohedral Group—Ullmannite Type
35*	Cubic SymmetryUllmannite
	I.00
264	Groups Unidentified Cubic symmetryBoléite .75
30±	Octahedral symmetry
38*	Trapezohedral symmetry, fig. 1356Leucite .30
39	Cubo-octahedral symmetry, fig. 2135 Dysanalyte .25
	II. Tetragonal System
ang inte	The forms in this system are referred to three axes, all at right les to one another. The two lateral axes $a$ and $b$ are equal and exchangeable, while the vertical axis $c$ differs from these in length in character.
	Normal Group—Zircon Type
40 <b>*</b> 41	Unit pyramid $p$ (111), similar to fig. 1680Zircon .40 Unit pyramid $p$ (111) and base $c$ (001)Octahedrite .75

42*	Unit prism $m$ (110) and unit pyramid $p$ (111), fig.  1681
43*	Unit prism m (110) and two unit pyramids p (111)
44	and $u$ (331), fig. 1682
	two unit pyramids $p$ (111) and $u$ (331), similar to fig. 1682Zircon .40
45*	fig. 1682
46	Unit and diametral prisms $m$ (110) and $a$ (100),
	unit and diametral pyramids $p$ (111) and $e$ (101) and base $e$ (001), similar to fig. 1667
47	Diametral prism a (100) and unit pyramid p (111), similar to fig. 1687
48	Unit and diametral prisms $m$ (110) and $a$ (100) and unit pyramid $p$ (111), fig. 1690
49*	Unit and diametral prisms m (110) and a (100) and
50	diametral pyramid e (101), fig. 934Rutile .50 Unit, diametral and ditetragonal prisms m (110),
	a (100) and $l$ (310), unit and diametral pyramids $e$ (101) and $s$ (111)
51	e (101) and s (111)
	or zirconoid $x$ (311)
52	.50
53*	Diametral prism a (100) and unit pyramid p (111), fig. 1874
	.50
54₹	Diametral prism $a$ (100), unit pyramid $p$ (111) and base $c$ (001), fig. 1871
55	Diametral prism $a$ (100), two unit pyramids $p$ (111)
33	and s (113) and diametral pyramid e (101), simi-
	lar to fig. 955Octahedrite
	Pyramidal Group—Scheelite Type
56	Unit pyramid $p$ (111)
57	Unit and diametral pyramids $n$ (111) and $e$ (101) and base $e$ (001)
58*	Unit and diametral prisms m (110) and a (100) and
	unit pyramid r (111), fig. 1649
59	Unit prism $m$ (110) rounded, and base $c$ (001) Wulfenite .40
00 <b>*</b>	Unit pyramid $u$ (102) and base $c$ (001)
/1 <b>T</b>	and base $c$ (001), similar to fig. 2604

Sphenoidal Group—Chalcopyrite Type	
62* Sphenoid of first order p (111)	
63* Two sphenoids, plus p (111) and minus p <sub>1</sub> (111), octahedral symmetry, similar to fig. 274	
64 Acute sphenoid $\varphi$ (772) and scalenohedron $x$ (122), fig. 278	
III. Hexagonal System	
The forms in this system are referred to four axes. The three lateral axes, $a_1$ , $a_2$ and $a_3$ are equal and interchangeable and cross at angles of 60° and 120°, while the vertical axis $c$ is of different length and at right angles to them.	
Normal Group—Beryl Type	
65* Unit prism m (1010) and base c (0001), fig. 1508. Beryl .30 66* Unit prism m (1010), unit pyramid o (1011) and base c (0001), fig. 2480	
mids $s$ (1121) and $p$ (1122) and base $c$ (0001), similar to fig. 1505	
Hemimorphic Group—Iodyrite Type	
68* Unit prism m (1010), pyramid i (2021) and base c (0001), fig. 507	
Pyramidal Group—Apatite Type	
69* Unit prism <i>m</i> (1010) and base <i>c</i> (0001), similar to fig. 2219	
70* Unit prism $m$ (1010) and unit pyramid $x$ (1011),	
70* Unit prism m (10T0) and unit pyramid x (10T1), fig. 2193	
y (2021) and base $c$ (0001), similar to fig. 2194 Apatite .40 72* Unit and diametral prisms $m$ (1010) and $a$ (1120), two unit pyramids $x$ (1011) and $r$ (1012), diametral pyramid $s$ (1121) and base $c$ (0001), simi-	
lar to fig. 2197Apatite .75	
Pyramidal-Hemimorphic Group—Nephelite Type	
73* Unit prism m (1010) and base c (0001), similar to fig. 1537	
Rhombohedral Division	
Normal Group—Calcite Type	
74* Rhombohedron r (10T1), 74° 55′, fig. 1035 Calcite .40	

75	Rhombohedron $r$ (10 $\overline{1}$ 1), $73^{\circ}$
76	Rhombohedron $r(10T1)$ , 73°
77*	Obtuse rhombohedron e (0112), fig. 1037Calcite .30
78*	Acute rhombohedron f (0221), fig. 1070
70.	Acute rhombohedron $d$ (0881) and base $c$ (0001) Siderite .50
19	Positive and negative rhombohedrons
00+	Colored and negative monipoliculous
	Scalenohedron v (2131), fig. 1049
82	
83*	Scalenohedron $v$ (2131) and one rhombohedron $r$
_	(10T1), fig. 1051
84	Scalenohedron $v$ (2131) and prism $m$ (4041)Calcite .20
85	Unit prism $m$ (1010) and base $c$ (0001), fig. 1045 Calcite .30
86*	Unit prism $m$ (1010) and rhombohedron $e$ (0112),
	fig. 1039
87*	Unit prism $m$ (10 $\overline{10}$ ), rhombohedron $e$ (01 $\overline{12}$ ) and
•	scalenohedron v (2131), similar to fig. 1053 Calcite .20
88	Three scalenohedrons and two rhombohedrons,
•	similar to fig. 1057
89	Unit prism m (1010), three rhombohedrons and
09	two scalenohedrons
~~*	Acute pyramid $\gamma$ (8.8.16.3), second order Calcite .20
90*	Demonid of second and any (2272) and been a (0001). Committee 120
91	Pyramid of second order $n$ (2243) and base $c$ (0001), Corundum .40
92	Prism of second order a (1120) and pyramid v (4483)
93	$(44\overline{83})$
)3	base c (0001) Hematite .75
04*	base $c$ (0001)
74.	r (1011) and curved rhombohedron $u$ (1014),
	fig. 822Hematite .30
	Hemimorphic Group—Tourmaline Type
05*	Unit and second order prisms m (1010) and a
90	(1170) and rhombohedron $r$ (1071), fig. 1839 Tourmaline
	.40
06 <b>*</b>	Unit and second order prisms $m$ (10 $\overline{10}$ ) and $a$
90*	(1120) and two rhombohedrons $r$ (1011) and $o$
	(0221), fig. 1848Tourmaline
	.40
97	Striated rounded prisms $m$ (1010) and $a$ (1120)
	and steep rhombohedron y (4041)
	1.00
	Tri-rhombohedral Group—Phenacite Type
98	Unit and second order prisms m (10T0) and a
90	(1120) and third order rhombohedron $x$ (2132),
	fig. 1639Phenacite .40
00	Unit and second order prisms m (1010) and a
99	(1170) and two showball drops a (1071) and a
	(1170) and two rhombohedrons $r$ (1011) and $e$
	(01T2), similar to fig. 1637 Willemite .50

100 <b>*</b>	Rhombohedron $r$ (10 $\overline{1}$ 1), 73° 45′ Dolomite .20 Acute rhombohedron $M$ (40 $\overline{4}$ 1) Dolomite .30
	Trapezohedral Group-Quartz Type
102*	Two rhombohedrons $r$ (10I1) and $z$ (01I1), fig.
103*	589
	Unit prism $m$ (1010), rhombohedrons $r$ (1011) and $z$ (01T1) and trigonal pyramid $s$ (1121), fig.
105	593Quartz .40 Unit prism $m$ (1010), rhombohedrons $r$ (1011) and $z$ (0111) and acute rhombohedron $M$ (3031),
	fig. 594
107*	Unit prism $m$ (1010), rhombohedrons $r$ (1011) and $z$ (0111), trigonal pyramid $s$ (1121) and trigonal trapezohedron $x$ (5151). Left-handed
108	crystal, fig. 596
	IV. Orthorhombic System
	IV. Orthorhombic System in this system the forms are referred to three axes $a$ , $b$ , and $c$ at angles to one another and of unequal lengths.
	n this system the forms are referred to three axes $a$ , $b$ , and $c$ at
right	n this system the forms are referred to three axes a, b, and c at angles to one another and of unequal lengths.  Normal Group—Barite Type  Unit prism m (110) and base c (001), fig. 2418 Barite .30
right 109*	n this system the forms are referred to three axes a, b, and c at angles to one another and of unequal lengths.  Normal Group—Barite Type  Unit prism m (110) and base c (001), fig. 2418 Barite .30  Macrodome d (102) and brachydome o (011) Barite .30
right 109*	n this system the forms are referred to three axes a, b, and c at angles to one another and of unequal lengths.  Normal Group—Barite Type  Unit prism m (110) and base c (001), fig. 2418 Barite .30  Macrodome d (102) and brachydome o (011) Barite .30  Unit prism m (110), macrodome d (102) and base
right 109*	n this system the forms are referred to three axes a, b, and c at angles to one another and of unequal lengths.  Normal Group—Barite Type  Unit prism m (110) and base c (001), fig. 2418Barite .30  Macrodome d (102) and brachydome o (011)Barite .30  Unit prism m (110), macrodome d (102) and base c (001), similar to fig. 2420Barite .30  Unit prism m (110), macrodome d (102), brachypinacoid b (010), pyramid z (111) and base c
right 109* 110	n this system the forms are referred to three axes a, b, and c at angles to one another and of unequal lengths.  Normal Group—Barite Type  Unit prism m (110) and base c (001), fig. 2418Barite .30  Macrodome d (102) and brachydome o (011)Barite .30  Unit prism m (110), macrodome d (102) and base c (001), similar to fig. 2420Barite .30  Unit prism m (110), macrodome d (102), brachypinacoid b (010), pyramid z (111) and base c (001), similar to fig. 2425Barite .50  Macrodome d (102), brachydome o (011), macro-
right 109* 110 111 112	n this system the forms are referred to three axes a, b, and c at angles to one another and of unequal lengths.  **Normal Group—Barite Type**  Unit prism m (110) and base c (001), fig. 2418 Barite .30 Macrodome d (102) and brachydome o (011) Barite .30 Unit prism m (110), macrodome d (102) and base c (001), similar to fig. 2420 Barite .30 Unit prism m (110), macrodome d (102), brachypinacoid b (010), pyramid z (111) and base c (001), similar to fig. 2425 Barite .50 Macrodome d (102), brachydome o (011), macropinacoid a (100) and base c (001) Barite .30 Unit prism m (110), macrodome d (102), brachydome o (011), brachypinacoid b (010) and
right 109* 110 111 112	n this system the forms are referred to three axes a, b, and c at angles to one another and of unequal lengths.  *Normal Group—Barite Type*  Unit prism m (110) and base c (001), fig. 2418 Barite .30  Macrodome d (102) and brachydome o (011) Barite .30  Unit prism m (110), macrodome d (102) and base c (001), similar to fig. 2420

117*	Unit and obtuse pyramids p (111) and s (113) and
0	base $c$ (001), sphenoidal type, similar to fig. 20. Sulphur .60
118	Unit and obtuse pyramids $p$ (111) and $s$ (113) and brachydome $n$ (011), similar to fig. 17Sulphur .60
119*	Unit and obtuse pyramids $p$ (111) and $s$ (113),
	brachydome $n$ (011) and base $c$ (001), fig. 15. Sulphur .40
120	Unit prism m (110), brachypinacoid b (010) and
	three pyramids $p$ (111), $s$ (113) and $r$ (343), similar to fig. 113
121	Unit prism m (110) and brachydome u (014), fig.
	360Arsenopyrite
	.75
122*	Unit prism $m$ (110), pyramid $o$ (111) and brachy-
	pinacoid $b$ (010)
123*	Unit prism $m$ (110), macrodome $t$ (106) and base
	c (001) striated, similar to fig. 2412 Thenardite .20
124*	Unit prism m (110), brachydome s (011) and base
125	c (001), fig. 1717 And alusite .50 Unit prism $m$ (110), macrodome $d$ (102), macro-
123	pinacoid $a$ (100) and base $c$ (001)
126	Unit prism $m$ (110), macrodomes $d$ (102) and $l$
	(104), brachydome $o$ (011), macropinacoid $a$
	(100), pyramids and base $c$ (001)
127	(100), pyramids and base $c$ (001)
	.50
128*	Unit prism m (110), pyramids e (122) and z (112), similar to fig. 958
	similar to fig. 958
129	and breehydome t (021)  Received to A co
120	and brachydome $t(021)$
130	striated, similar to fig. 2458Anhydrite .40
131	Unit and brachyprisms $m$ (110) and $l$ (120) and
-0-	two brachydomes $f(021)$ and $y(041)$
132	Unit and brachyprisms $m$ (110) and $l$ (120),
	brachydome $y(041)$ and base $c(001)$
133*	Unit and brachyprisms m (110) and l (120), unit
	and obtuse pyramids $u$ (111) and $i$ (221),
	brachydome s (041), pyramid x (243) and base $x = x^{(001)} \text{ similar to for } x^{(001)}$
134	c (001), similar to fig. 1706
134	brachydomes $f$ (021) and $y$ (041), pyramids $o$
	(221), $u$ (111), $i$ (223) and base $c$ (001)
125*	Unit and brachyprisms $m$ (110) and $l$ (120), unit
-30	and obtuse pyramids $u$ (111) and $o$ (221),
	brachydome $y$ (041), macrodome $d$ (201) and
	base $c$ (001), similar to fig. 1703
136	Unit and brachyprisms $m$ (110) and $l$ (120), unit
-	and obtuse pyramids $u$ (111), $o$ (221) and $i$ (223),

	brachydome $y$ (041), brachypinacoid $b$ (010), macrodome $d$ (201) and base $c$ (001), similar to
137*	fig. 1709
138*	base $c$ (001), similar to fig. 1857 Staurolite .40 Unit prism $m$ (110), brachypinacoid $b$ (010), one
139	sets of macrodomes $r$ (101) and base $c$ (001).
140	fig. 1857
141	to fig. 1218
142	Unit prism $m$ (110) and pyramid $r$ (131) striated. :Childrenite
143	Macropinacoid a (100), brachypinacoid b (010), macrodome k (103), pyramids o (111) and u
144	(133) and base $c$ (001), similar to fig. 2146 Columbite .75 Unit prism $m$ (110), prism $g$ (130), macropinacoid $a$ (100), macrodomes $h$ (201), $k$ (103) and $l$ (106),
145*	pyramids and base c (001), similar to fig. 2147. Columbite 2.00 Unit prism m (110), prism s (120), brachypinacoid b (010), brachydome k (103) and macrodome d (101), similar to fig. 1612
	Hemimorphic Group—Calamine Type
146*	Unit prism $m$ (110), macropinacoid $a$ (100), brachypinacoid $b$ (010), two macrodomes and brachypinacoid $b$ (010).
147	dome $i$ (031)
148*	Macrodome $s$ (101), brachypinacoid $b$ (010) and base $c$ (001), similar to fig. 2287 Struvite .30
149	Unit prism m (110), macrodome s (101) and base c (001)
	Sphenoidal Group—Epsomite Type
150	Unit prism $m$ (110), sphenoid $z$ , plus and minus (111)Epsomite .50

### V. Monoclinic System

In this system the forms are referred to three axes, a, b and c, of unequal lengths, with a and c intersecting at an acute angle behind, while b is at right angles to a and c.

### Normal Group—Gypsum Type

151*	Unit prism $m$ (110), clinopinacoid $b$ (010) and unit
	pyramid $l$ (111), fig. 2501
152*	Unit prism $m$ (110), clinopinacoid $b$ (010), unit
	pyramid l (111) and orthodome e (103), fig.
	2504
153	Unit and clinoprisms $m$ (110) and $k$ (130), clino-
	pinacoid $b$ (010), unit pyramid $l$ (111) and
	orthodome $e$ (103)
154	Unit prism $m$ (110), clinopinacoid $b$ (010) and
	base c (001), similar to fig. 1291
155*	Unit prism m (110), clinopinacoid b (010), ortho-
(	dome $y$ (201) and base $c$ (001), fig. 1297 Orthoclase .20
150*	Unit prism m (110), prism z (130), clinopinacoid
	b (010), orthodome $y$ (201) and base $c$ (001) Orthoclase .20
157	Unit prism m (110), clinopinacoid b (010), ortho-
158	dome y (701), pyramid $\sigma$ (111) and base $\sigma$ (001). Orthoclase .40
150	Unit prism $m$ (110), prism $s$ (130), clinopinacoid $b$ (010), orthodome $y$ (201), pyramid $o$ (111) and
	base $c$ (001), similar to fig. 1301 Orthoclase .40
150*	Unit prism $m$ (110), orthodome $x$ (101) and base
-39.	c (001)
160	Unit prism m (110), prism z (130), clinopinacoid b
	(010), orthodome $x$ (101) and base $c$ (001) Adularia .30
161	Unit prism m (110), orthopinacoid a (100), ortho-
	dome $w$ (101) and pyramid $r$ (111) Monazite .50
162*	Unit prism m (110), orthopinacoid a (100), clino-
	pinacoid b (010), orthodome $\phi$ (T01), pyramids
	$u$ (111), $s$ (111), $\lambda$ (331) and base $c$ (001),
	similar to fig. 1325
163*	Unit prism m (110), orthopinacoid a (100), clino-
_	pinacoid b (010) and pyramid s (T11), fig. 1400. Augite .30
164*	Unit prism m (110), orthopinacoid a (100), clino-
_	pinacoid $b$ (010), pyramids $u$ (111) and $o$ (221). Augite .30
165	Unit prism $m$ (110), brachypinacoid $b$ (010),
	brachydome t (032), macrodome e (302), pyra-
	mids $n$ (331) and $q$ (332) and base $c$ (001) Herderite 4.00
166*	Unit prism m (110), clinopinacoid b (010), clino-
	dome $r$ (011) and orthodome $p$ (101), similar
	to fig. 1485
167*	Unit prism $m$ (110) and pyramid $q$ (T11), fig.
	2555Kröhnkite 1.00
168*	Unit prism m (110), pyramid n (111) and base c
_	(001), fig. 2113Titanite .40
169	Unit prism $m$ (110), orthodome $x$ (102) and base
	c (001)
170*	Unit pyramids p. (111) and e (111)Lazulite .40

171	Unit prism $m$ (110), macrodomes $v$ (101) and $k$ (101), brachydome $r$ (011) and base $c$ (001), similar to $6a$ 746.
172*	similar to fig. 546
	and base $c$ (001)
173	
174*	Prism $M$ (221), clinopinacoid $b$ (010) and base $c$ (001), similar to fig. 1957
175	Unit prism $m$ (110), pyramid $h$ (221), orthodomes $\sigma$ (101) and $\theta$ (101), clinodomes $l$ (023) and $p$ (021) and base $c$ (001)
176*	Unit prism $m$ (110), orthopinacoid $a$ (100), pyramid $h$ (221) and two orthodomes
177	Unit prism $m$ (110), pyramid, orthodome $\sigma$ (101) and base $c$ (001), similar to fig. 1252
178	Unit prism $m$ (110), clinopinacoid $b$ (010), orthodomes $s$ (201) and $t$ (201) and base $c$ (001), fig. 1881
179	Unit prism $m$ (110), orthopinacoid $a$ (100), orthodome $x$ (102), clinodomes $m_x$ (011), $g$ (012) and $t$ (013), pyramids $n$ (111) and $\epsilon$ (112) and base $c$ (001), similar to fig. 1733
180	Unit prism m (110), clinodome e (011) and pyramid r (112)
181	Unit prism $m$ (110), ortho- and clinopinacoids $a$ (100) and $b$ (010), pyramids $z$ (221) and $o$ (111) and base $c$ (001), fig. 2402
182	Unit prism $m$ (110), prism $f$ (120) and pyramid $t$ (111), similar to fig. 2465
183	Unit prism $m$ (110), clinodomes $z$ (011) and $w$ (012), orthodome $k$ (101) and base $c$ (001), similar to fig. 2468
184	Long prism $m$ (110) striated, and clinodome $z$ (011) Crocoite .75
185*	Long unit prism $m$ (110) striated, clinodomes $z$ (011) and $w$ (012), pyramid $t$ (111), orthodome $k$ (101) and base $c$ (001)
186*	Unit prism $m$ (110), prism $t$ (210), orthopinacoid $a$ (100), clinodomes $\kappa$ (011) and $a$ (021), orthodome and pyramids $\beta$ (111) and $\omega$ (131). Colemanite .50
187	Unit prism $m$ (110), prism $t$ (210), orthopinacoid $a$ (100), clinopinacoid $b$ (010), orthodomes $h$ (201) and $i$ (101), clinodomes $k$ (311) and $a$ (100), pyramids $\beta$ (111), $v$ (221), $k$ (311), $v$ (111) and base $c$ (001), similar to fig. 2395

### Clinohedral Group—Clinohedrite Type

### VI. Triclinic System

In this system the forms are referred to three axes, a, b, and c, of unequal lengths, and which intersect at oblique angles.

### Normal Group—Axinite Type

	2
189*	Unit prisms $M$ (110) and $m$ (110), macropinacoid
	a (100), macrodome $s$ (201) and pyramids $r$
	(111) and $x$ (111), fig. 1774
100	(111) and $x$ (111), fig. 1774
190	one prisms 22 (110) and 36 (170), macropinacoid
	a (100), brachyprism w (130), brachydome y
	(021), pyramids $r$ (111) and $n$ (131) and base $c$
	(001) similar to fig. 1775 Axinite .75
1014	(001), similar to fig. 1775
1917	1 (10) is in and m (110), brachyphiacold
	b (010), brachydome $o$ (II1), macrodome $x$ (I01)
	and base c (001), fig. 1338 Pericline .40
<b>IQ2</b> *	Unit prisms $M$ (110) and $m$ (110), macrodome $x$
	(701) and base $c(001)$ Albite 20
104	(101) and base $c$ (001)
193	onit prisms M (110) and M (110), brachypmacoid
	b (010), brachydome $e$ (021), macrodome $y$ (201),
	pyramid $p$ (I11) and base $c$ (001)
104*	Unit prisms $M$ (110) and $m$ (110), macrodome $x$ (101), brachypinacoid $b$ (010) and base $c$ (001). Amazonstone
-74	(TO1) brachypinacoid h (O10) and base c (O01) Amazonstone
	.20
195	Unit prisms $M$ (110) and $m$ (110), prisms $z$ (130)
	and $f$ (130), brachypinacoid $b$ (010), macro-
	dome $x$ (101), pyramid $o$ (111) and base $c$ (001). Amazonstone
	, , , , , , , , , , , , , , , , , , ,
	.30
190	Unit prisms M (110) and m (110), prisms z (130)
	and f (130), brachypinacoid b (010), macro-
	domes $\hat{x}$ (T01) and $\hat{y}$ (201) and base $\hat{c}$ (001) Amazonstone
	.40
T07#	Prisms W (170) and w (110) macroningsoid a
19/+	Prisms $M$ (110) and $m$ (110), macropinacoid $a$ (100), brachypinacoid $b$ (010) and pyramid $q$
	(100), brachypinacold $b$ (010) and pyramid $q$
	(011)Cyanite 1.00
<b>198</b> *	Prisms $M$ (110) and $m$ (110), brachypinacoid $b$ (010), pyramid $q$ (221) and base $c$ (001) Rhodonite .75
- / -	(010) pyramid $a$ (221) and hase $c$ (001) Rhodonite .75
***	Driema M (170) and m (110) breehypingsoid h
199	Prisms $M$ (110) and $m$ (110), brachypinacoid $b$ (010) and pyramid $k$ (221), fig. 1434
	(U1U) and pyramid $k$ (221), hg. 1434
200	Prisms $M$ (110) and $m$ (110), pyramids $h$ (221), $g$ (111), $f$ (443), $d$ (221) and base $c$ (001) Babingtonite
	$g(\overline{111}), f(\overline{443}), d(\overline{221})$ and base $g(\overline{001}), \dots, \overline{Babingtonite}$
	1.50
	1.50

### Twins

I. Isometric System	tem
---------------------	-----

	1. Isomew System
201	Octahedrons o (111), contact, tw. pl. parallel to octahedral face, fig. 859
202*	octahedral face, fig. 859
203*	Pyritohedrons e (210) penetration, tw. axis normal to dodecahedral face, fig. 303
204	Tetrahedrons o (111), contact tw. pl. parallel to octahedral face, fig. 199Sphalerite .30
	II. Tetragonal System
205	
205	Prismatic, tw. pl. parallel to pyramid e (101), fig. 1686Zircon .50
	Prismatic, tw. pl. parallel to pyramid e (101) Rutile .50
207	Prismatic, tw. pl. parallel to pyramid e (101), repeated twinning
208*	repeated twinning
	repeated twinning, eightling, fig. 947Rutile .30
209	
-	.75
210	Tw. pl. parallel to a pyramid face (trilling)Cumengéite  -75
211	Tw. pl. parallel to a pyramid face (truncated trilling)
	111. 110.000000000000000000000000000000
212	Contact, tw. pl. pyramid e (3034), fig. 508 Iodyrite .40
212	Contact, tw. pl. pyramid e (3034), fig. 508 Iodyrite .40 Acute rhombohedrons, penetration. Vertical or
	c axis, tw. axis
	c axis, tw. axis
214*	c axis, tw. axis
214*	c axis, tw. axis
214* 215*	c axis, tw. axis
214* 215*	c axis, tw. axis
214* 215* 216*	c axis, tw. axis
214* 215*	c axis, tw. axis
214* 215* 216*	c axis, tw. axis
214* 215* 216* 217 218*	c axis, tw. axis
214* 215* 216* 217 218* 219	c axis, tw. axis
214* 215* 216* 217 218* 219	c axis, tw. axis
214* 215* 216* 217 218* 219 220*	c axis, tw. axis
214* 215* 216* 217 218* 219 220*	c axis, tw. axis
214* 215* 216* 217 218* 219 220*	c axis, tw. axis
214* 215* 216* 217 218* 219 220* 221 222*	Caxis, tw. axis
214* 215* 216* 217 218* 219 220* 221 222*	Caxis, tw. axis

225	Prismatic, contact, tw. pl. prism m (110)
228	Repeated twinning, tw. pl. m (110), "Wheel Ore," fig. 415
229	Penetration, tw. pl. parallel to macrodome e (101), Arsenopyrite
231* 232 233* 234	Cruciform, tw. pl. brachydome x (032), fig. 1859. Staurolite .75 Cruciform, tw. pl. pyramid z (232), fig. 1860 Staurolite .50 Cruciform, tw. pl. brachydome e (011), fig. 2414. Thenardite .20 Contact, tw. pl. prism m (110), "Spear head" twin Cerussite .50 Contact, tw. pl. prism m (110), stellate twin, fig. 1223 Cerussite .75 Contact, tw. pl. prism m (110), reticulated twinning Cerussite .50
	V. Monoclinic System
237 238 239* 240* 241 242* 243 244* 245*	Contact, tw. pl. orthopinacoid a (100), fig. 1402. Augite .30 Contact, tw. axis c, similar to fig. 2280. Lazulite .50 Contact, tw. axis c, Carlsbad twin. Orthoclase .50 Penetration, tw. axis c, Carlsbad twin, fig. 1302. Orthoclase .30 Contact, tw. pl. clinodome n (021), Baveno twin, fig. 1305. Orthoclase .40 Penetration, tw. pl. orthopinacoid a (100). Gypsum .30 Contact, tw. pl. orthopinacoid a (100), "Swallow-tail" twin, fig. 2511. Gypsum .20 Contact, tw. pl. orthopinacoid a (100). Titanite .75 Contact, tw. pl. base c (001), fig. 2558. Kröhnkite 1.00 Cruciform-penetration, tw. pl. base c (001), compounded on e (011), then twinned on m (110), similar to fig. 1888. Harmotome
240+	Cruciform-penetration, tw. pl. base c (001) simple form
	VI. Triclinic System
247*	Albite Law, tw. pl. brachypinacoid b (010), polysynthetic, cleavageLabradorite
248 249	Pericline law, tw. pl. parallel to b axis

250	Manebach law, tw. pl. base c (001), similar to fig. 1306	
	Regular Groupings of Crystals	
251*	Parallel growth of crystals of one species, arborescent, fig. 70	
252	Parallel growth of crystals of one species, rosette, fig. 827	
253* 254*	Parallel growth of crystals of one species, capped. Amethyst .50 Parallel growth of crystals of two species  Cyanite within Staurolite .50	
255*	Parallel growth of crystals of two species Chalcopyrite on Sphalerite .30	
	Irregularities of Crystals	
Disto	rtion:	
256	Elongated cube a (100) Fluorite .50	
257*	Twisted or saddle-shaped cube a (100)	
258	Flattened dodecahedron d (110)	
259*	Elongated dodecahedron $d$ (110)	
26t	Florgated cubo-octahedron a (100), a (111) Dysanalyte .20	
262	Elongated cubo-octahedron $a$ (100), $o$ (111) Dysanalyte .20 Elongated trapezohedron $n$ (211)	
263*	Flattened rhombohedron	
264*	Abnormal development of one rhombohedronQuartz .30	
265	Abnormal development of opposite rhombohedral	
266	facesQuartz .30 Elongated rhombohedron and flattened prismQuartz .30	
Impe	rfections on the Surfaces of Crystals:	
267	Striations due to oscillatory combination, on cube,	
~6Q+	fig. 290	
200*	nrism Ouartz 20	
269		
	rhomb	
270	Striations due to repeated twinning, cleavage Microcline . 20	
	Markings from erosion, etc., on cube	
272	Markings from erosion, etc., on pyramid	
274	Pseudo-octahedral symmetry, parallel arrange-	
-/4	ment of minute cubo-tetrahexahedronsFluorite .40	
275*	Pseudo-octahedral symmetry, parallel arrange-	
	ment of small dodecahedronsFluorite .40	
276*	Curved surfaces due to oscillatory combinations,	
	prism and scalenohedron	

277* Curved surfaces due to independent molecular conditions, rounded, saddle-shaped, fig. 1133 Dolomite .20 278* Curved surfaces due to independent molecular conditions, sheaf, fig. 1896
Internal Imperfections and Inclusions
283* Enclosing liquid with moving bubble
16. 1/20
Pseudomorphs
By Substitution:
291* Quartz replacing
By Deposition:
292 Incrustation of Quartz on
By Alteration:
294* Paramorph of Rutile after
forming
forming
ing
300 Partial exchange of constituents of Muscovite, forming

# Index to Complete Crystal List Price List of Loose Crystals

As the same form or combination may sometimes be found in many species, the collection which does not duplicate forms, necessarily omits some important minerals. The following can generally be furnished as individual crystals when desired.

The Roman numeral before each name indicates the system of crystallization: I. Isometric; II. Tetragonal; III. Hexagonal or Rhombohedral; IV. Orthorhombic; V. Monoclinic; VI. Triclinic.

The number or numbers after the names, indicate their position in the preceding Descriptive List of the Complete Crystal Collection.

V.	Adularia, 159, 160\$ .30	I. Boléite, 36\$ .75
VI.	Albite, 192	I. Boracite, 31, 32
VI.	Albite, twin, 24860	V. Borax, 181
	Alexandrite, twin 2.50	IV. Bournonite, twin, 228 1.50
	Amazonstone, 194-196 .20— .40	IV. Brochantite50
	Amazonstone, twin, 25075	IV. Bromlite50
III.	Amethyst, 25350	IV. Brookite, 128, 12940 -1.00
	Amphibole, 163, 16430	IV. Brookite, alt., 12750
I.	Analcite40	IV. Calamine, 14650
II.	Anatase, Octahedrite, 41, 55	III. Calcite, 74, 77, 78, 80-90, 269,
	75- 1.∞	276, 291
IV.	Andalusite, 12450	III. Calcite twin, 215-217 .40- 1.00
	Anglesite, 125, 12675- 1.00	II. Cassiterite 1.00
IV.	Anhydrite, 130	II. Cassiterite, twin50- 1.00
VI.	Anorthite, 19350	IV. Celestite, 115, 1164075
	Anorthite, twin 1.00	IV. Cerussite, 140, 29340— .50
	Apatite, 70-724075	IV. Cerussite, twin, 233-235
II.	Apophyllite, 52-544050	.50— .75
	Aragonite twin, 223, 224.30— .50	III. Chabazite, 7620
	Arsenopyrite, 121	III. Chabazite, twin, 21420
	Arsenopyrite, twin, 22925	IV. Chalcocite, twin 1.00
V.	Augite, 163, 16430	II. Chalcopyrite, 62-6430- 1.00
	Augite, twin, 236 30	II. Chalcopyrite, twin, 209
	Axinite, 189, 19050— .75	VI. Chesterlite
	Azurite, 175-1775075	IV. Chiastolite, 29040
	Babingtonite, 200 1.50	IV. Childrenite, 142 1.00
	Barite, 109-1143050	V. Chondrodite
	Beryl, 65, 67, 27930— 1.00	IV. Chrysoberyl, twin, 22750
	Beryllonite, 28450	IV. Chrysolite, 145
V.	Blödite 1.50	III. Cinnabar, 10850
	· · · · · · · · · · · · · · · · · · ·	,

III.	Cinnabar, twin, 213\$ .20	IV. Glaucodot\$1.∞
V.	Clinohedrite, 188 4.00	III. Gmelinite, twin
	Cobaltite50	V. Gypsum, 151–153, 28720
V.	Colemanite, 186, 187 .50— .75	V. Gypsum, twin, 241, 242
IV.	Columbite, 143, 14475— 2.00	.20— .30
	Copper, 251, 25930— .50	I. Halite, 260, 28120— .25
Ш.	Corundum, 91, 92, 272 .30— .50	III. Hanksite, 66
	Covellite 3.00	V. Harmotome, twin, 245 40
V.	Crocoite, 182-185	I. Hauerite 1.00
V.	Cryolite, 171	III. Hematite, 93, 94, 252,
II.	Cuméngéite	263, 299
II.	Cuméngéite, twin, 210, 211	V. Herderite, 165 4.00
	.50— .75	V. Heulandite, 178
· I.	Cuprite	V. Hornblende, 166
I.	Cuprite, alt., 14 1.25	II. Hyacinth, 47
VI.	Cyanite, 197 1.00	II. Idocrase, Vesuvianite,
	Cyanite, twin, 249	45, 46
IV.	Danburite	III. Iodyrite, 68
V.	Datolite, 17950	III. Iodyrite, twin, 21240
	Diamond 1.00	III. Jarosite 1.00
IV.	Diaspore50	V. Kröhnkite, 167 1.00
	Diopside, 162	V. Kröhnkite, twin, 244 1.50
	Dioptase 1.00	VI. Labradorite, twinned cleavage
Ш.	Dolomite, 100, 101, 277 .20— .30	247 1.50
	Durangite25	IV. Laurionite
	Dysanalyte, 37, 39, 261 .20— .25	V. Lazulite, 170
I.	Embolite	V. Lazulite, twin, 23750
	Enargite50	V. Leadhillite, twin2.50
	Endlichite	I. Leucite, 3830
	Enstatite40	IV. Libethenite, 141 1.00
	Epididymite50	I. Magnetite, 220
	Epidote, 172	II. Malacon, 48
٧.	Epistilbite75	IV. Manganite50
	Epsomite, 150	IV. Marcasite
	Eudialyte 1.00	IV. Marcasite, twin, 22630
	Eudidymite50	I. Martite, 299
	Eudidymite, twin	II. Matlockite 2.00
1.	Fluorite, 4, 6, 8, 13, 256, 271,	II. Meionite 1.50
_	273-275, 292,20— 1.50	II. Melilite
	Fluorite, twin, 202	IV. Meneghinite40
_	Fowlerite, Rhodonite, 19875	VI. Microcline, 194–19620— 40
_	Franklinite, 10	VI. Microcline, twin, 250
	Gahnite50	I. Microlite, 11, 12 1.00— 1.50
	Galena, 1, 7, 930— .50	III. Mimetite
1.	Garnet, 3, 5, 15, 16,	III. Molybdenite40
	258, 262	V. Monazite, 16150
	Gay-Lussite, 180	IV. Monticellite
V.	Glauberite	V. Muscovite, 173, 17420

V.	Natrochalcite\$3.00	II. Scheelite, 56\$1.00
IV.	Natrolite, 122	IV. Scorodite 1.00
	Nephelite, 73	I. Senarmontite
	Neptunite 1.00	III. Siderite, 75, 793050
	Newberyite	I. Smaltite
	Northupite	III. Smithsonite40
II.	Octahedrite, 41, 5575— 1.00	I. Sphalerite, 25530
VI.	Oligoclase, 288	I. Sphalerite, twin, 20430
	Olivenite 1.00	V. Sphene, 168, 16940— .75
	Orthoclase, 154-158 .2040	V. Sphene, Titanite, twin, 24375
V.	Orthoclase, twin, 238-240	I. Spinel
	20 50	I. Spinel, twin,20150
V.	Pachnolite,	II. Stannite3.00
VI.	Paisbergite, Rhodonite, 199. 1.00	IV. Staurolite, 137-1393050
V.	Penninite50	IV. Staurolite, twin, 230, 231
	Pericline, 191	.5075
VI.	Pericline, twin, 24830	IV. Stephanite, 14750
Ш.	Phacolite, twin, 218, 219	IV. Stibnite, 120
	.75— 1.00	V. Stilbite, 278
I.	Pharmacosiderite 1.00	V. Stilbite, twin
Ш.	Phenacite, 98	II. Stolzite, 57
V.	Phillipsite, twin, 246	IV. Strontianite, twin
	Phosgenite75	IV. Struvite, 148, 149 30—
V.	Pinite, 300	IV. Sulphur, 117–11940— .60
	Pirssonite 2.00	I. Tetrahedrite, 3050
Ш.	Proustite 1.25	IV. Thenardite, 123
III.	Pyrargyrite 1.00	IV. Thenardite, twin, 23220
I.	Pyrite, 17-29, 257, 267	II, Thorite 2.50
	.20— .75	V. Titanite, 168, 169 40— .75
	Pyrite, twin, 203	V. Titanite, twin, 243
	Pyrolusite40	IV. Topaz, 131-1362075
	Pyromorphite, 6930	III. Tourmaline, 95–97,280
III.	Pyrosmalite	.30— 1.00
	Pyroxene, 162–164	III. Troostite, Willemite, 9950
	Pyroxene, twin, 23630	I. Ullmannite, 35 1.00
	Pyrrhotite	I. Uraninite 1.00
Ш.	Quartz, 102–107, 264-266,	III. Vanadinite
	268, 282, 283, 285, 286,	II. Vesuvianite, 45, 4650— .75
	289	V. Vivianite
	Quartz, twin, 220-222 .50- 1.50	П. Wernerite, 5850
	Raspite, twin 1.00	III. Willemite, Troostite, 9950
<u>v</u> .	Realgar75	IV. Witherite, twin, 225
	Rhodochrosite50	V. Wolframite40
	Rhodonite, 198, 19975— 1.00	II. Wulfenite, 59, 60, 61 .40— 1.25
	Rutile, 49, 5050	II. Xenotime40
	Rutile, twin, 206-208 . 3075	Π. Zircon, 40, 42-44, 5120— .40
	Sal-ammoniac, 3440	II. Zircon, twin, 205
П.	Scapolite, 5850	I. Zunyite, 33

### Elementary

### No. 77A. School Crystal Set

The crystals selected for this set, while essentially the same as those in the advanced collections, are generally over I cm. in length, many reaching 3 or 4 cm. Being intended for elementary work, they are sufficiently sharp for contact measurement, while many are bright enough for the reflecting goniometer. As far as practicable, the commoner forms and habits, of symmetrical and model-like aspect, have been used in planning the list and selecting the actual specimens.

On the back of the 3 x 4 cm. tray accompanying each crystal, is pasted our miniature label giving name, composition and locality. The light and neat mahogany cabinet holding

the collection, is shown in Plate VIII.

Single crystals are sold at the prices in the School Crystal List. They total \$18.35. The "collection price" for the fifty crystals, with trays, in mahogany cabinet, delivered to any address, is \$15.00. Without cabinet, 10 per cent. less.

### No. 77A. School Crystal List

I. Isometric System		
I	Cube a (100), fig. 512	Fluorite .20
2	Octahedron o (111)	Magnetite .20
3	Dodecahedron $d$ (110)	Garnet .30
4	Trapezohedron $n$ (211), fig. 1356	Leucite .30
5	Cube $a$ (100) modified by octahedron $o$ (111)	Galena .30
6	Octahedron $o$ (111) and dodecahedron $d$ (110)	Franklinite .75
7	Dodecahedron $d$ (110) modified by trapezohedron $n$	,
•	(211), fig. 1578	Garnet .40
8	Dodecahedron $d$ (110) modified by octahedron $d$	•
	(111)	Cuprite, .75
9	Pyritohedron e (210), fig. 289	Pyrite .20
10	Cube a (100), modified by pyritohedron $\epsilon$ (210), fig.	1
	295Limonite pseudomorph after	Pyrite .20
II	Twin, penetration, pyritohedrons $e$ (210), tw. axis	3
	normal to dodecahedral face, fig. 303	Pyrite .30
	II. Tetragonal System	
12	Unit prism $m$ (110) and unit pyramid $b$ (111), fig.	
	Unit prism m (110) and unit pyramid p (111), fig. 1681	Zircon .20
13	Unit and diametral prisms m (110) and a (100) and	
Ū	diametral pyramid e (101), fig. 934	Rutile .50
14	Unit and diametral prisms m (110) and a (100) and	
•	base c (001)	Vesuvianite .75

15 Diametral prism a (100), unit pyramid p (111) and base c (001), fig. 1871
.40 Unit and diametral pyramids u (102) and s (113) and base c (001), similar to fig. 2604
17 Sphenoid of first order $p$ (111)
III. Hexagonal System
19 Unit prism m (10T0) and base c (0001), fig. 1508 Beryl .30
20 Unit prism $m$ (1010) and unit pyramid $x$ (1011),
fig. 2193Apatite .40 21 Unit prism m (1010) and unit pyramids x (1011)
and $y(20\overline{21})$
22 Rhombohedron $r$ (10 $\overline{1}$ 1), $73^{\circ}$
23 Rhombohedron $r$ (1011), $73^{\circ}45'$
24 Scalenohedron v (2131), fig. 1049
(1011) and curved rhombohedron $u$ (1014), fig.
822
and two rhombohedrons $r$ (10T1) and $\rho$ (02Z1).
fig. 1848
and g (0111), ng. 584
28 Twin, scalenohedrons v (2131), contact, tw. pl. base
29 Twin, penetration, normal rhombohedron r (1011),
tw. axis c, fig. 1912
IV. Orthorhombic System
30 Macrodome $d$ (102), brachydome $o$ (011), macropinacoid $a$ (100) and base $c$ (001)
31 Unit prism m (110), pyramid o (111) and brachy-
pinacoid $a$ (100) and base $c$ (001)
brachydome $n$ (011) and base $c$ (001), fig. 15 Sulphur .40 33 Unit prism $m$ (110), brachydome $s$ (011) and base $c$
(001), fig. 1717

#### V. Monoclinic System

	· · ·
39	Unit prism m (110), clinopinacoid b (010) and unit
	pyramid <i>l</i> (111), fig. 2501
40	Unit prism m (110), clinopinacoid b (010), ortho-
4 T	dome $y$ (201) and base $c$ (001), fig. 1297Orthoclase .20 Unit prism $m$ (110), orthopinacoid $a$ (100), clino-
4+	pinacoid $b$ (010) and pyramid $s$ (111), fig. 1400Augite .30
42	Unit prism m (110), clinopinacoid b (010), clino-
	dome $r$ (011) and orthodome $\phi$ (101), similar to
	fig. 1485
43	Unit prism $m$ (110), pyramid $n$ (111) and base $c$ (001),
	fig. 2113
44	Unit prism m (110), ortho- and clinopinacoids a
	(100) and b (010), pyramids z (221) and e (111)
4 ~	and base $c$ (001), fig. 2402
45	(001), similar to fig. 1957
46	Unit prism $m$ (110), clinodomes $z$ (011) and $w$ (012),
Ψ-	orthodome $k$ (101) and base $c$ (001), fig. 2468 Crocoite .75
47	orthodome $k$ (101) and base $c$ (001), fig. 2468 Crocoite .75 Twin, penetration, tw. axis $c$ , Carlsbad twin Orthoclase .30
	VI. Triclinic System
ΔQ	Unit prisms M (110) and m (110), macropinacoid
40	a (100), macrodome s (201) and pyramids $r$ (111)
	and x (111), fig. 1774
40	Unit prisms M (110) and m (110), prisms z (130)
"	and $f(130)$ , brachypinacoid $b(010)$ , macrodome $x$
	(101), pyramid $o$ (111) and base $c$ (001)
50	Unit prisms $M$ (110) and $m$ (110), macrodome $x$
	(10T) and base $c$ (001)

### No. 80. Lecture Table Crystals

Twenty-five crystals, about 10 x 7 cm. (4 x 23/4 in.).

The number of these is limited by nature's supply, as few minerals occur in crystals large enough to be recognized across a room, or even when passed from hand to hand among the class. We have, however, arranged this incomplete series, embracing merely representative examples of the simpler forms. Some are a little rough in outline, but all are sufficiently well defined to illustrate the form, and are eminently adapted to this purpose. A number of these minerals are rarely found in such very large crystals, hence the total of the individual values exceeds \$45.00 The "collection price," delivered to any address, in trays and No. I chest, is \$40.00. Without chest 10 per cent. less. List sent on application.

## PART VII

Physical Mineralogy

Hardness, Structure,
Color, Effect of Radium, Etc.

## Physical Mineralogy

Series Illustrating Hardness, Structure, Specific Gravity,

Color, Effect of Radium, etc.

With the exception of crystals, there are no collections prepared by us upon which are bestowed a greater amount of expert labor than in the selection of just the right specimens to illustrate the various physical characters of minerals. Thus, in the hardness sets, crystals or cleavages are selected; under structure, color and luster, a particular specimen from among many of its kind is carefully chosen with a view to exactly illustrating the required characteristic; the specimens selected for specific gravity tests are as pure and compact as they are found in nature, thus approximating the theoretical ratio; the cleavage series has been extended and illustrations of parting added. In general the definitions of Dana have been followed.

It should be borne in mind that the mere names of minerals opposite the different terms, do not indicate the nicety of variation shown by the individual specimens chosen. The same species often well represents different characters. As far as practicable, however, the duplication of species has been avoided.

The entire physical series, and notably the color section, makes one of our most showy and attractive collections.

FREE TRANSPORTATION to any address.

PRICES include pasteboard trays, or blocks with museum specimens if requested; also oak chests (or without chests 10 per cent. less), where minerals total \$20.00.

The following are according to the Complete Physical Series List.

#### No. 92A. Hardness Series

Ten museum specimens averaging 12 x 9 cm.  $(434 \times 3\frac{1}{2} \text{ in.})$ , \$8.00.

#### No. 92. Student's Hardness Series

Ten hand specimens averaging 10 x 7 cm.  $(4 \times 2\frac{3}{4} \text{ in.})$ , \$4.00.

### No. 93A. Fusibility Series

Six museum specimens averaging 12 x 9 cm.  $(4\frac{3}{4} \times 3\frac{1}{2} \text{ in.})$ , \$4.00.

#### No. 93. Student's Fusibility Series

Six hand specimens averaging 10 x 7 cm. (4 x 23/4 in.), \$2.00.

#### No. 94A. Structure Series

Twenty-five museum specimens averaging 12 x 9 cm.  $(4\frac{3}{4} \times 3\frac{1}{2} \text{ in.})$ , \$25.00.

#### No. 94. Student's Structure Series

Twenty-five hand specimens averaging 10 x 7 cm.  $(4 \times 2)$ 4 in.), \$12.50.

### No. 95A. Cleavage, Fracture and Tenacity Series

Twenty-five museum specimens averaging 12 x 9 cm.  $(4\frac{3}{4}$  x  $3\frac{1}{2}$  in.), \$18.00.

### No. 95. Student's Cleavage, Fracture and Tenacity Series

Twenty-five hand specimens averaging 10 x 7 cm. (4 x  $2\frac{3}{4}$  in.), \$9.00.

### No. 96A. Taste, Odor and Feel Series

Nine museum specimens averaging 12 x 9 cm.  $(4\frac{3}{4} \times 3\frac{1}{2} \text{ in.})$ , \$5.00.

#### No. 96. Student's Taste, Odor and Feel Series

Nine hand specimens averaging 10 x 7 cm.  $(4 \times 2\frac{3}{4} \text{ in.})$ , \$2.50.

### No. 97A. Specific Gravity Series

Twenty-five museum specimens averaging 12 x 9 cm.  $(4\frac{3}{4} \times 3\frac{1}{4} \text{ in.})$ , \$40.00.

### No. 97. Student's Specific Gravity Series

Twenty-five hand specimens averaging 10 x 7 cm. (4 x  $2\frac{3}{4}$  in.), \$20.00.

#### No. 101A. Color Series

Fifty museum specimens averaging 12 x 9 cm.  $(4\frac{3}{4} \times 3\frac{1}{2} \times 3)$  in.), \$50.00.

#### No. 101. Student's Color Series

Fifty hand specimens averaging 10 x 7 cm.  $(4 \times 2\frac{3}{4} \text{ in.})$ , \$25.00.

#### No. 102A. Luster Series

Twenty-five museum specimens averaging 12 x 9 cm. (43/4 x  $3\frac{1}{2}$  in.), \$25.00.

#### No. 102. Student's Luster Series

Twenty-five hand specimens averaging 10 x 7 cm.  $(4 \times 2)$ 4 in.), \$12.50.

### No. 104A. Series Illustrating Effect of Radium, Etc.

Twenty-five museum specimens averaging 12 x 9 cm. (43/4 x  $3\frac{1}{2}$  in.), \$25.00.

No. 104. Student's Series Illustrating Effect of Radium, Etc.

Twenty-five hand specimens averaging 10 x 7 cm.  $(4 \times 2)$ 4 in.), \$12.50.

### No. 111A. Complete Physical Series

Includes all of the foregoing "A" series. Two hundred museum size specimens averaging 12 x 9 cm. (4¾ x 3½ in.). The total of the individual museum specimen values in the Physical Series List exceeds \$230.00. Delivered to any address, with trays (or blocks if requested), in four No. 3 chests, \$200.00. Without chests 10 per cent. less.

### No. 111. Student's Complete Physical Series

Includes all of the foregoing "Student's" Series. Two hundred hand size specimens averaging 10 x 7 cm. (4 x 2¾ in.). The total of the individual hand specimen values in the Physical Series List exceeds \$115.00. Delivered to any address, with trays, in two No. 3 chests, \$100.00. Without chests, 10 per cent. less.

## Physical Series List

### Entire List Constitutes Nos. 111A and 111.

### Hardness, Nos. 92A and 92

1 Hardness 1Talc	6 Hardness 6Feldspar		
2 Hardness 2 Gypsum	7 Hardness 7Quartz		
3 Hardness 3 Calcite	8 Hardness 8Topaz		
4 Hardness 4 Fluorite	9 Hardness 9 Corundum		
5 Hardness 5Apatite	10 Hardness 10 Diamond		
Fusibility, No	Fusibility, Nos. 93A and 93		
11 Fusibility 1 Stibnite	14 Fusibility 4 Actinolite		
12 Fusibility 2 Natrolite	15 Fusibility 5Orthoclase		
13 Fusibility 3 Almandite	16 Fusibility 6Bronzite		
Structure, Nos. 94A and 94			
17 BladedCyanite	30 Mammillary Chalcedony		
18 Columnar Tremolite	31 Globular Pisolite		
19 Fibrous Asbestus	32 Nodular Menilite		
20 Reticulated Cerussite	33 Amygdaloidal Laumontite		
21 Stellated Wollastonite	34 Coralloidal Flos Ferri		
22 Radiated Tourmaline	35 Dendritic Wad		
23 Curved Folia Talc	36 Mossy Calc Tufa		
24 Straight Folia Biotite	37 Capillary Chalcotrichite		
25 Coarse GranularSandstone	38 Acicular Aragonite		
26 Fine Granular Marble	39 Drusy Quartz		
27 Compact Magnesite	40 Stalactitic Stalactite		
28 Friable Bauxite	41 Amorphous Deweylite		
29 Velvety Aurichalcite			

### Cleavage, Parting, Fracture and Tenacity, Nos. 95A and 95

Cleavage, Parting, Fracture and Tenacity, Nos. 95A and 95		
Cleavage	Fracture	
42 Cubic	54 Conchoidal Smoky Quartz	
43 Octahedral Fluorite	55 Even Lithographic Stone	
44 Dodecahedral Sphalerite	56 Uneven Rhodonite	
45 Basal Apophyllite	57 Hackly Franklinite	
46 Prismatic Amphibole	58 Earthy Turgite	
47 Clinodiagonal Orthoclase	59 Splintery Pectolite	
48 Rhombohedral Calcite	TENACITY	
49 Pinacoidal Gypsum	60 Brittle Siderite	
D	61 Tough Emery	
PARTING	62 Imperfectly Sectile. Alabaster	
50 Basal Pyroxene	63 Highly Sectile Embolite	
51 Pyramidal Corundum	64 Malleable Copper	
52 Octahedral Magnetite	65 Flexible Itacolumite	
53 Hemi-orthodome Adularia	66 Elastic Muscovite	
Taste, Odor and Feel, Nos. 96A and 96		
TASTE	71 Sulphurous Pyrite	
67 Saline Halite	72 Bituminous Asphaltum	
68 Alkaline Natron	73 Argillaceous Kaolinite	
69 Bitter Carnallite	74 Fetid Anthraconite	
ODOR	FEEL	
70 Alliaceous Mispickel	75 Greasy Graphite	
Specific Gravity.	Nos. 97A and 97	
	are approximate.)	
Unmetallic Luster	89 Sp. Gr. 4.7Zircon	
76 Sp. Gr. 1.0Copalite	69 5p. 67. 4.7	
77 Sp. Gr. 1.6 Anthracite	Metallic Luster	
78 Sp. Gr. 1.9 Thaumasite	90 Sp. Gr. 5.0	
79 Sp. Gr. 2.1Opal	91 Sp. Gr. 5.7Arsenic	
80 Sp. Gr. 2.3Gypsum	92 Sp. Gr. 6.0 Arsenopyrite	
81 Sp. Gr. 2.6Albite	93 Sp. Gr. 6.2Smaltite	
82 Sp. Gr. 2.8 Prochlorite	94 Sp. Gr. 6.7Cassiterite	
83 Sp. Gr. 3.0Cryolite	95 Sp. Gr. 7.5	
84 Sp. Gr. 3.2 Apatite	96 Sp. Gr. 8.0 Cinnabar	
85 Sp. Gr. 3.5 Titanite	97 <i>Sp. Gr. 8.9</i> Copper	
86 Sp. Gr. 3.8Limonite	98 <i>Sp. Gr. 9.8</i> Bismuth	
87 Sp. Gr. 4.0 Sphalerite	99 Sp. Gr. 13.6 Mercury	
88 <i>Sp. Gr.</i> 4.3Witherite	100 Sp. Gr. 18.0	

### Color, Nos. 101A and 101

Color, Nos. 101A and 101		
RED  101 Flesh-Red Chabazite 102 Rose-Red Rose Quartz 103 Scarlet-Red Crocoite 104 Orange-Red Wulfenite 105 Purplish-Red Cinnabar 106 Garnet-Red Almandite 107 Brick-Red Jasper 108 Blood-Red Zincite YELLOW 109 Sulphur-Yellow Sulphur 110 Orange-Yellow Orpiment 111 Ochre-Yellow Ochre 112 Resin-Yellow Calcite 114 Brownish-Yellow Dolomite GREEN 115 Olive-Green Olivine 116 Sage-Green Serpentine 117 Verdigris-Green Amazonstone 118 Sea-Green Fluorite 119 Emerald-Green Brochantite 120 Apple-Green Garnierite 121 Grass-Green Malachite 122 Leek-Green Williamsite VIOLET 123 Reddish-Violet Amethyst	127 Azure-Blue	
124 Bluish-Violet Sodalite BLUE 125 Indigo-Blue Covellite 126 Prussian-Blue Chalcanthite	147 Silver-White Arsenopyrite 148 Bronze-Yellow Pyrrhotite 149 Copper-Red Copper 150 Brass-Yellow Chalcopyrite	
Luster, Nos. 102A and 102.		

KINDS OF LUSTER	155 Greasy Elæolite
151 MetallicJamesonite	156 PearlyDolomite
152 Adamantine Endlichite	157 SilkySatin Spar
153 Vitreous	DEGREES OF LUSTER
154 ResinousSphalerite	158 Splendent Hematite

306 PHYSICAL	MINERALOGY	
159 ShiningDolomite 160 GlisteningPapierspath 161 GlimmeringFlint LUSTER PHENOMENA 162 Play of ColorsOpal 163 Change of Colors Labradorite 164 OpalescenceMoonstone 165 ChatoyancyTiger Eye 166 IridescenceCoal 167 TarnishBornite	168 Dichroism Epidote 169 Asterism Phlogopite 170 Schiller Sunstone DIAPHANEITY 171 Transparent Quartz 172 Semi-Transparent Fluorite 173 Translucent Alabaster 174 Semi-Translucent Onyx 175 Double Refraction Iceland Spar	
Effect of Radium, Röntgen, and Ultra-Violet Rays, Heat, Friction and Magnetism. Nos. 104A and 104.		
RADIUM  176 Phosphorescent Diamond  177 Fluorescent Willemite  RÖNTGEN RAYS  178 Fluorescent, Blue Fluorite  179 Phosphorescent, White Aragonite	188 Pyro-Electric with Lateral PolarityQuartz, hexagonal 189 Thermo-ElectricPyrite 190 Phosphorescent, Blue. Chlorophane 191 Phosphorescent, RedLepi-	
180 Opaque	dolite FRICTION  192 — Electricity Amber 193 + Electricity Quartz 194 Triboluminescent, Red . Hexagonite 195 Triboluminescent, Yellow Dolomite MAGNETISM	

enite

186 Opaque......Mica

HEAT

187 Pyro-Electric with Terminal

Polarity.... Tourmaline,

rhombohedral

196 Polarity.....Lodestone

197 Strongly Magnetic . . . . Pyrr-

198 Weakly Magnetic . . . . Garnet 199 Paramagnetic......Siderite

200 Diamagnetic . . . . Wulfenite

hotite

## Part VIII

Chemical Mineralogy

Collections of Specimens for Analysis

Laboratory List

of Minerals sold by Weight

## Chemical Mineralogy

### Specimens for Blowpipe and Wet Analysis

The material selected for these collections is as near chemically pure as the minerals usually occur in nature. All are clean, typical examples of distinct species. The list embraces those commonly covered in an elementary course, and includes most of the minerals recommended by Penfield, Dana, Toula, Plattner and other writers.

If a more extended collection is desired, it may be selected from the alphabetical Price List in Part III. or the School of Mines List in Part V. If the price there given is in any instance for impure material, a smaller but pure specimen will be supplied, provided the order mentions "for analysis." If instead of trimmed specimens, a more extensive set of pure lumps and fragments is preferred, they may be purchased by weight from the Laboratory List, pages 311 to 319.

### No. 119A. Series for Chemical Analysis.

One hundred museum size specimens of pure minerals, averaging 12 x 9 cm. (43/4 x 31/2 in.). The average weight is about 1000 grams (2.2 pounds avd.), generally affording sufficient of each kind for 400 to 500 analyses. Individual museum specimens may be purchased at double the listed hand size prices. The sum of such individual values in the museum size is \$129.20. The "collection price" for all the specimens is \$100.00, delivered to any address. This price includes pasteboard trays, or blocks if requested, and two No. 3 Oak Chests. Without chests, 10 per cent. less. If preferred, an equal weight of fragments in boxes, instead of trays or blocks, is sold in chests at \$100.00.

PURCHASE IN PARTS. Free delivery with trays and No. 3 Chest. Without chest, 10 per cent. less.

PART I. 50 names marked with \*, totaling \$59.70 ...\$50.00 PART II. 50 remaining names, totaling \$69.50 ..... 50.00

### No. 119. Student's Series for Chemical Analysis

One hundred hand size specimens, averaging 10 x 7 cm. (4 x 23/4 in.). Like the preceding, but smaller. The average weight is about 450 grams (1 pound avd.), generally affording sufficient of each kind for 200 to 250 analyses. Individual specimens sold at listed prices. These total \$64.60. The "collection price" for all the specimens is \$50.00, delivered to any address. This includes pasteboard trays with one No. 3 Oak Chest. Without chest, 10 per cent. less. If preferred, an equal weight of fragments in boxes, instead of trays, is sold in chest at \$50.00. According to the Chemical List.

PURCHASE IN PARTS. Free delivery with trays and No. 2 Chest, shown in Plate II. Without chest, 10 per cent. less. PART I. 50 starred names \*, totaling \$29.85.....\$25.00 PART II. 50 remaining names, totaling \$34.75..... 25.00

#### Chemical List

	0-0	
	Aluminium, Al	Chromium, Cr
I	Corundum\$ .50	18* CHROMITE20
2*	BAUXITE	Cobalt, Co
3*	CRYOLITE	19 SMALTITE\$1.00
4	<b>WAVELLITE</b>	20* COBALTITE60
5	ALUNITE	21* GLAUCODOT 1.00
	Antimony, Sb	Copper, Cu
6*	STIBNITE	22 CHALCOCITE 1.00
	Arsenic, As	23 BORNITE
7*	REALGAR 1.00	24* CHALCOPYRITE 35
	ARSENOPYRITE25	25 TETRAHEDRITE 1.00
0.	Barium, Ba	26 ENARGITE 1.00
^		27* CUPRITE 1.50
-9		28* MALACHITE 1.25
10*	BARITE	
	_ Beryllium, Be	29 CHRYSOCOLLA50
II	Beryl	30* CHALCANTHITE 50
	Bismuth, Bi	Gold and Tellurium, Au, Te
12*	BISMUTITE 1.00	31* SYLVANITE 2.00
		_ Iron, Fe
	Boron, B	32* Pyrite
13	BORAX	33* Hematite
14	COLEMANITE50	34 MAGNETITE
	Calcium, Ca	35 FRANKLINITE40
I 5*	FLUORITE	36 LIMONITE20
16*	CALCITE20	37* SIDERITE20
17	GYPSUM20	
-/	GIISUM	38 DUFRENITE

	Lead, Pb	Tantalum and Columbium,
39*	GALENA \$ .40	Ta, Cb
	JAMESONITE 1.00	72 Columbite\$ .50
	CERUSSITE 1.25	Thorium, Th
42	Pyromorphite75	73 THORIANITE 2.50
43	WULFENITE 1.00	Tin, Sn
	Anglesite 1.50	· · · · · · · · · · · · · · · · · · ·
45	CROCOITE 1 . 00	74 STANNITE
	Lithium, Li	75* Cassiterite50
46*	LEPIDOLITE	Titanium, Ti
47	Amblygonite50	76* RUTILE
	Magnesium, Mg	77 ILMENITE
48*	MAGNESITE	Tungsten, W
49*	DOLOMITE	78* Wolframite
	Kieserite	79 SCHEELITE
•	Manganese, Mn	Radium and Uranium, Ra, U
51	ALABANDITE 1.00	80* URANINITE 3.00
52*	Pyrolusite	Vanadium, V
53	MANGANITE 1.00	
	RHODOCHROSITE75	81* VANADINITE 1.00
55		Yttrium, Y (with Er, La, Di)
	Mercury, Hg	82 GADOLINITE 2.50
56*	CINNABAR 1.25	Zinc, Zn
	Molybdenum, Mo	83* Sphalerite20
57*	MOLYBDENITE40	84 ZINCITE
	Nickel, Ni	85* CALAMINE
	MILLERITE 1.00	86* Smithsonite40
	NICCOLITE 1.00	87 WILLEMITE60
	Pyrrhotite	Zirconium, Z
61*	GARNIERITE	88 ZIRCON
	Phosphorus, P	
62*	APATITE	Silicates, Insoluble
	Potassium, K	89 ALBITE
	SYLVITE	90* PYROXENE
64*	CARNALLITE30	91* AMPHIBOLE, Actino-
	Selenium, Se	lite,
65	GUANAJUATITE 2.00	92* GARNET, Almandite30
	Silver, Ag	93 EPIDOTE
66	ARGENTITE 1.25	94 SERPENTINE20
67*	Pyrargyrite 1.00	95* KAOLINITE
	Sodium, Na	96 TOURMALINE40
	HALITE	Silicates, Soluble
69		97 WOLLASTONITE 75
	Strontium, Sr	98 DATOLITE
70	STRONTIANITE20	99* NATROLITE
71*	CELESTITE	100 STILBITE

### Laboratory List

### Minerals Sold by Weight

Ton Lots of many rare minerals supplied to experimenters and manufacturers. Prices on request if quantity desired is stated. Correspondence solicited with producers and consumers of rare ores, also colored semi-precious stones, such as Azurite, Turquois Matrix, etc., etc.

QUALITY. Pieces usually consist of irregular lumps or fragments of about 3 to 9 cm (1½ to 3½ in.) length, more or less. Specimens trimmed to uniform sizes cost more. The material furnished is about as pure as found in Nature. Where more than 5 per cent. of gangue rock or matrix is attached, the per cent. of pure mineral is noted.

FREE TRANSPORTATION to any address, with privilege of returning any unsatisfactory item at our expense.

EXTRA STRONG CARTONS (double-thick cylindrical cardboard boxes) hold each mineral conveniently and permanently.

A MINIMUM PRICE of \$0.20 is charged for even the smallest quantity of any mineral sold by weight.

10 TO 50 KG. samples of one mineral cost proportionately less than listed. Thus 10 kg. or over, 10 per cent. less. 50 kg., 20 per cent. less.

LESS THAN THE LISTED QUANTITY is charged at a rate 25 per cent. higher proportionately than the list price. Thus Manganotantalite listed at \$3.00 per kilo, costs \$0.37 for one-tenth kilo; Glaucodot at \$4.00 per kilo costs \$1.25 for one-quarter kilo; Argyrodite at \$2.50 D. costs \$0.31 G.; Beryl at \$0.40 K. costs \$0.20 for ¼ K. (minimum charge), etc.

A METAL CLASSIFICATION of the economic minerals in this list, showing the minerals carrying each metal, will be found in Part V.

Foreign Money is, for convenience, accepted as follows: \$1.00=4/-=M. 4.=Fcs. 5=L. 5.

#### COMPARISON OF WEIGHTS

```
1000 grams = 1 kilogram (K.) = about 2½ pounds avoirdupois.
100 " = 1 hectogram (H.) = " 3½ ounces "
10 " = 1 dekagram (D.) = " ½ ounce "
1 gram (G.) = " 15½ grains
```

## Laboratory List

Prices per kilo (2.2 lbs.)

Achroite, crystals, D., \$1.50	Anhydrite \$ .20
Actinolite, crystalline\$ .40	Anhydrite, vein in halite40
Adularia 2.00	Annabergite 4.00
Adularia, Moonstone, pre-	Ånnerödite, H., \$2.00
cious, H., \$1.50	Anorthite, xls., H., \$1.50
Aegyrite 2.00	Anthophyllite, radio-fibrous .40
Aeschynite, H., \$1.00	Anthracite Coal20
Agalmatolite 1.00	Anthraconite, Stinkstone 20
Agate, banded or moss 50	Antimony, H., \$1.25
Alabandite, 75 per cent 2.00	Apatite, granular, brown 20
Alabaster	granular, green20
Albertite	compact, whitish20
Albite, lamellar, white 20	See Phosphate Rock.
Albite cleavage, striated. 20	Apophyllite, H., \$0.75
Algodonite, H., \$1.00	Aquamarine, D., \$0.40
Allanite	Aragonite, banded 30
Allemontite, H., \$1.00	Argentite, D., \$0.75
Allophane, cupriferous 1.00	Argyrodite, D., \$2.50
Almandite, large crystals40	Arkansite, xls., D., \$1.00
Aluminite, H., \$1.50	Arkansite, paramorphosed
Alunite	to rutile, crystals 1.00
Amazonstone, crystallized .40	Arsenic I.25
Amber, H., \$0.75	Arsenopyrite
Amblygonite, cleavable, 70	Asbestus, Amphibole, gray .40
Amethyst, deep colored 3.00	Asbestus, Chrysotile, (Ser-
Amethyst, light colored I .00	pentine), green 1.00
Amethyst, light with milky	Asbolite, Earthy Cobalt 50
quartz 50	Asphaltum
Amphibole. See following:	See also: Elaterite, Wurtzilite, Alber- tite, Gilsonite.
Actinolite, Asbestus, Byssolite, Hex- agonite, Hornblende, Tremolite.	Atacamite 4.00
Analcite, H., \$0.75	Augite, crystals 2.00
Anatase, crystals, G., \$1.25	Aventurine, Oligoclase 2.00
Andalusite 3.00	Averturine, Perthite40
Andorite, D., \$0.60	Awaruite, grains in magnetite
Andradite, granular, pink,	sand, D., \$2.50
50 per cent	Axinite, yellow or brown I .00
Anglesite	Azurite 4.00
A LIEUTE STATE OF THE PARTY OF	,

Barite, lamellar \$ .20	Calcite, xl., Nail Head Spar \$1.00
Basanite	See also: Calc Tufa, Chalk, Hydraulic
Bauxite, pisolitic, yellowish .20	Limestone, Iceland Spar, Limestone, Lithographic Stone, Marble, Onyx,
Bauxite, nodules in clay20	Travertine.
Berthierite, 50 per cent 4.00	Cancrinite, H., \$0.60
Beryl, green or yellow 40	Cancrinite, 5 per cent in
(Ton Lots at Market Prices.)	nephelite-syenite40
Beryl, Aquamarine, D., \$0.40	Cannel Coal
Beryllonite, D., \$3.00	Carnallite
Biotite60	Carnotite, 10 per cent 2.00
Bismuth, H., \$0.60	Cassiterite, massive 2.00
Bismuthinite, H., \$0.90	Cassiterite, in feldspar30
Bismutite, H., \$2.50	Cassiterite, Stream Tin 1 . 00
Bituminous Coal, irides-	Celestite, cleavage40
cent	Celestite, fibrous
F	Cerargyrite, D., \$1.00
To:	Cerite 1 . 50
Blende, cleavable40 Blödite, crystals 1.00	Cerussite, massive, 75 per
Bog Iron Ore	cent
Boleite, crystals, D., \$1.00	Cerussite, cryst'd, white 1.60
D . C. C	Cervantite
Borax	Chabazite, H., \$0.75
TD	Chalcanthite 2.00
Bornite, argentiferous 1.50 Bort, carat, \$4.00	Chalcedony, nodules 50
Boulangerite, 50 per cent. 1.50	Chalcedony, mammillary,
Bournonite, H., \$2.00	with coral impressions 50
D	Chalcocite 1.50
Brochantite, massive 4.00	Chalcophanite 1.00
Brochantite, fibrous, 50	Chalcopyrite
per cent 3.00	Chalcopyrite, 33 per cent 20
	Chalk
Bröggerite, D., \$1.00	Chert
Bronzite, sublamellar, gray .40 Brookite, xls., D., \$1.00	Chlorastrolite, H., \$1.00
	Chlorite, Prochlorite 40
Brookite, paramorphosed	Chloritoid, Masonite50
to rutile, crystals 1.00	Chlorophyllite
Brown Coal	Chondrodite 2.00
Brucite, H., \$0.75	Chromite
Byssolite	Chrysocolla50
Calamine	Chrysolite, Dunite30
Calcite, cleavages 20	Chrysolite, gem, D., \$0.75
Calcite, rhombic cleavages .40	Chrysotile, Asbestus 1.00
Calcite, crystals, scalenohe-	Cinnabar 4.00
drons	Cinnabar, 10 per cent 1.00

Cinnana Stand	Callindrida do on
Cinnamon Stone\$ .60	Cylindrite\$2.00
Citrine 1.50	Datolite 2.00
Clinochlore 2.00	Datolite, 25 per cent 1.25
Coal. See following:	Descloizite, H., \$1.50
Anthracite, Bituminous, Lignite, Cannel	Deweylite 1.00
Cobaltite, pure, compact . 2.50	Diallage with saussurite50
Cobaltite, 10 per cent50	Diaspore, lamellar 4.00
Coccolite	Diopside 1 . 00
Colemanite 1.00	Dolomite, fine, white20
Colophonite 1.50	Dolomite, coarse, yellow 20
Columbite 2.00	Dolomite, compact20
Copiapite 2.50	Dolomite, Pearl Spar 1.00
Copper, native 2.00	Domeykite 2.00
Copper, native in con-	Domeykite, 40 per cent 1.00
glomerate, 5 per cent40	Dysanalyte cryst'ls, H. \$1.50
Copper Glance 1.50	Dyscrasite, D., \$1.50
Copper Pyrites 50	Elæolite 1.00
Copper Pyrites, 33 per cent20	Elaterite 1.00
Cordierite 3.00	Embolite, D., \$1.00
Corundum, cleavages 1.00	
Corundum, crystals 1.00	Emery, granular
Corundum, Emery 20	Emplectite, H., \$1.00
Corundum, Ruby, D., \$2.50	Enargite, cleavable 2.00
Corundum, Sapphire, crys-	Endlichite with wulfenite 4.00
tals, D., \$0.50	Enstatite, sublamellar gray 40
Corundum, Sapphire, aste-	Epidote, nodules, compact 50
riated, water-worn crys-	Epidote, crystallized 50
tals, H., \$2.00	Epidote, gray 50
Covellite, bright foliated 3.00	Erythrite, H., \$0.60
Covellite, with pyrite 2.00	Eudialyte 4.00
Covellite, dull, platinifer-	Euxenite, H., \$0.75
ous (sperrylite) 3.00	Feldspar, Calcium — see
Crocidolite, altered to	Anorthite
Quartz, Tiger-Eye, cha-	Feldspar, Plagioclase—see
toyant	Albite, Oligoclase, Lab-
Crocidolite, unaltered 1.00	radorite.
Crocoite, crystals 2.00	Ell D. I. O.
Cryolite	thoclase and Microcline.
Cryolite with siderite30	
Cuprite 2.00	Feldspar, Soda—see Albite
Cuprite, 2 per cent	Fergusonite, H., \$1.50
Cyanite, bladed, blue50	Fibrolite
Cyanite, bladed, green75	Fire Opal, H., \$1.00
Cyanite, clear blue crystals	Flexible Sandstone40
in paragonite 1.50	Fluorite, greenish20

Fluorite, cubes, clear emer-	Gypsum, Satin Spar \$ .50
ald-green, H.,\$0.50	Gypsum, Selenite, clear
Fluorite, cleavages, trans-	colorless cleavage30
lucent, pink\$ .60	Halite, clear cleavage40
Fowlerite, crystalline60	Halite, granular
Franckeite 2.00	Halloysite 2.00
Franklinite, granular40	Halotrichite 2.00
Franklinite, granular, with	Hardystonite60
zincite and willemite 60	Hardystonite with wille-
Freibergite 3.00	mite and franklinite75
Fuchsite 2.00	Hausmannite 1.00
Gadolinite 4.00	Heavy Spar, lamellar20
Galena, argentiferous	Heliotrope 1.50
Galena, argentif., 40 per cent 50	Hematite, compact20
Galena, cleavable50	Hematite, crystallized 1.00
Garnet, Almandite, large	Hematite with jasper, "Jas-
crystals	pilite"
Garnet, Andradite, pink,	Hematite, micaceous
granular, 50 per cent 1.00	Hematite, oölitic20
Garnet, Grossularite60	Hematite, Pencil Ore 1.00
Garnet, Polyadelphite60	Hercynite 1.50
Garnierite, 75 per cent	Hessite, D., \$1.50
Garnierite, 5 per cent 20	Heulandite, H., \$0.75
Gersdorffite, H., \$0.60	Hexagonite 1.00
Gibbsite	Hielmite, H., \$2.00
Gilsonite	Hornblende
Glaucodot 4.00	Horn Silver, D., \$1.00
Glauconite	Hübnerite 2.00
Gold, Rand Conglomerate 50	Hyacinth, D., \$0.75
Gold, native, G., \$1.25	Hyalite, H., \$1.50
Gold Pyrites	Hydrotalcite, Houghite75
Goslarite 2.00	Hypersthene, cleavage 3.00
Göthite 1.50	Iceland Spar, colorless 4.00
Graphite, lumps	
Graphite, powdered 40	
Gray Copper 2.00	Idocrase
Gray Copper, argentif 3.00	Idrialite, H., \$1.00
Grossularite60	Ilmenite
Guanajuatite, D., \$2.50	Infusorial Earth20
Guano	Iolite, Chlorophyllite75
Gummite, H., \$2.00	Iolite, Cordierite 3.00
Gypsum, Alabaster 20	Iridosmine, G., \$2.00
Gypsum, coarsely fibrous 20	Iron, Meteoric, Aerolite
Gypsum, granular 20	(stone), D., \$2.00

Iron, Meteoric, altered to	Limonite, Bog Iron Ore\$ .20
limonite-magnetite shale\$1 .00	Limonite, fibrous
Iron, Meteoric, Siderite,	Limonite, iridescent40
plates, H., \$4.00	Limonite, Yellow Ochre 20
Iron, Meteoric, Siderolite	Linnæite, H., \$1.25
iron and stone, H., \$4.00	Lodestone, extra strong75
Iron, Terrestrial, H., \$3.00	Lodestone, strong25
Iron Pyrites, crystallized40	Löllingite 1.00
Iron Pyrites, massive20	Ludwigite 3.00
Itacolumite	Magnesite, compact20
Jade (Jadeite) 2.00	Magnesite, Pinolite, cleav 20
Jamesonite 3.00	Magnetite, granular20
Jasper, red50	Magnetite, crystallized
Jasper, variegated 50	Magnetite, see Lodestone.
Jasper, yellow	Malachite 2.00
Jasperized Wood50	Manganite 1.00
Jefferisite60	Manganotantalite 3.00
Jeffersonite	Marble, fine, white20
Josephinite, D., \$0.50	Marble, coarse, red20
Kainite	Marcasite 1.00
Kaolinite	Margarite, lamellar, pink . 1.25
Kaolinite containing piso-	Margarite, schistose, green 1.00
lites of bauxite20	Margarodite 1.00
Keilhauite 2.50	Margarodite, with brown
Kieserite	tourmaline, dravite 1.50
Kjerulfine, H., \$0.75	Martite, crystallized 50
Kröhnkite, broken crystals	Masonite50
H., \$0.60	Massicot, D., \$0.30
Labradorite, chatoyant60	Meerschaum 3.00
Labradorite, ordinary 30	Melaconite 2.50
Lapis Lazuli, azure-blue 2.00	Melanterite 2.00
Laumontite 4.00	Meliphanite, H., \$3.00
Laumontite amygdules in	Mellite, D., \$1.25
diabase, 5 per cent40	Meneghinite, xls., D., \$2.00
Lazulite, H., \$1.00	Menilite
Lazurite 2.00	Meteorites (see Iron)
Lepidolite, fine granular,	Mexican Onyx30
pale lilac	Mica: See Muscovite, Bi-
Lepidolite, coarse, scaly,	otite, Phlogopite, etc.
deep bluish-violet 20	Microcline, cleavage20
Leucite, H., \$1.00	Microcline, Amazonstone,
Lignite	crystal
Limestone, compact, gray,	Microlite, xls., D., \$3.00
blue, buff, white, etc 20	Milky Quartz, massive 20

Pyroxene. See following: Augite, Coccolite, Diopside, Heden-	Sapphire, crystals, D., \$0.50
bergite, Jeffersonite.	Satin Spar, fibrous, white. \$ .50
Pyrrhotite, nickeliferous \$ .20	Scapolite50
Pyrrhotite with pyrite20	Scheelite 2.00
Quartz. See following:	Scheelite, 10 per cent 50
Agate, Amethyst, Basanite, Chalced-	Schorlomite 2.00
ony, Chert, Citrine, Flint, Helio- trope, Itacolumite, Jasper, Jasper-	Scolecite, H., \$0.75
ized Wood, Milky, Moss Agate,	Selenite, clear cleavages 30
Rock Crystal, Rose, Smoky.	Semi-opal, see Opal
Rammelsbergite 3.00	Sepiolite, Meerschaum 3.00
Realgar	Serpentine, granular
Realgar, 10 per cent	Serpentine, Precious 60
Rhodochrosite 1.00	See also Chrysotile (Asbestus), Ophi- calcite, Verd-Antique, Williamsite.
Rhodonite60	Siderite
Rhodonite, Fowlerite with	Sillimanite, 50 per cent40
franklinite	Silver, native, D., \$0.60
Ripidolite 2.00	l ou
Rock Crystal, transparent 1.00	Silver, sulphide ore
Roemerite 4.00	A 1.1 1.
Rose Quartz, deep pink 1.00	l
Rose Quartz, pale pink 40	·
Rubellite, crystals, H., \$1.50	~ ~
Rubellite in lepidolite, 10	Smoky Quartz, enclosing
per cent 50	tourmaline 1.00
Ruby, D., \$2.50	Soapstone, talc
Ruby Silver, Dark, D., \$0.60	Sodalite
Ruby Silver, Light, D., \$1.00	Soda Niter
Ruby Spinel, water-worn	Soda Niter, stained with
crystals, H., \$1.50	chromic acid, high per- centage of iodine 1.00
Rutile, ordinary red, brown	
or black, containing iron .75	Sphalerite, cleavable
(Ton Lots at Market Prices.)	Sphalerite, granular
Rutile.—We keep in stock, for regular delivery to manufacturers, the	Sphalerite, 20 per cent20
best iron-free red grades, ground	Spinel, Ruby, water-worn
to fine yellows.	crystals, H., \$1.50
Sal-ammoniac, H., \$0.75	Spodumene, cleavable
Salt, Rock—see Halite.	Staffelite 1.50
Samarskite, H., \$1.00	Stannite 1.50
Sanidine 1.50	Stannite with pyrite
Sandstone, red, blue, gray,	Staurolite 4.00
white, etc	Steatite
Sandstone, banded60	Stephanite, D., \$1.00
Sandstone, flexible 40	Stibiotantalite, D., \$1.00
Sapphire, asteriated, water-	Stibnite
worn crystals, H., \$2.00	Stilbite 1.50

Stinkstone, Anthraconite . \$ .20	Turgite, with limonite \$ .20
Stream Tin 1 .00	Turgite, ocherous, red, loose 20
Strontianite	Turquois, H., \$0.75
Succinite 4.00	Turquois, 10 per cent 1.00
Sulphur60	Ulexite
Sulphur, 30 per cent40	Ullmannite 2.50
Sulvanite	Uraninite, D., \$0.60
Sunstone, Oligoclase 2.00	Uraninite, Bröggerite, cry-
Sunstone, Perthite 40	stals, D., \$1.00
Sylvanite, G., \$1.00	Uvarovite, green 1.50
Sylvite	Vanadinite, H., \$0.75
Tachhydrite 50	Verd-Antique40
Talc, foliated30	Vesuvianite 1.00
Talc, Steatite20	Vivianite, H., \$2.00
Tantalite 4.00	Wad
Tantalite, manganotantal-	Wad, Asbolite 1.00
ite 3.00	Wagnerite, H., \$0.75
(Ton Lots at Market Prices.)	Wavellite, 50 per cent 1.00
Tantalite with cassiterite 2.00	Wernerite50
Tetrahedrite 2.00	Willemite 1.00
Tetrahedrite, argentiferous 3.00	Willemite with franklinite
Thaumasite 1.00	and zincite60
Thomsenolite, H., \$1.00	Williamsite
Thorianite, xls., H., \$3.00	Witherite
Thorite, H., \$2.50	Wolframite 2.00
Thorite, Orangite, H., \$4.00	Wollastonite, stellated 2.00
Thulite, fine pink50	Wulfenite, crystallized 4.00
Tiger Eye, yellow 50	Wulfenite with endlichite . 4.00
Titanite, crystals, brown 80	Wurtzilite
Titanium oxide—see Rutile.	Wurtzite 1.50
(Ton Lots at Market Prices.)	Xanthosiderite 1.00
Topaz, broken crystals 4.00	Xenotime, H., \$0.75
Topaz, massive 2.00	Yellow Ochre
Tourmaline, black	Zincite 2.50
Tourmaline, brown60	Zincite, 50 per cent 1.25
Tourmaline, Rubellite,	Zincite with franklinite,
crystals, H., \$1.50	and willemite 60
Tourmaline, Rubellite, 10	Zinc Blende, see Sphalerite.
per cent. in lepidolite 50	Zinkenite, H., \$1 . 50
Tremolite	Zinnwaldite
Tremolite, 50 per cent 40	Zircon 1.25
Triphylite 2.00	Zircon, Hyacinth, D., \$0.75
Triplite, H., \$0.75	Zoisite, columnar, gray 50
Tripolite	Zoisite, Thulite fine pink 50

### Index

Alphabetical Price List	212	Industrial Minerals and Ores	253
Aluminium Minerals	270	Iron Minerals	262
Antimony Minerals	264	Kilo Prices	311
Approval System	7	Labels	12
Arsenic Minerals	274	Laboratory List, prices by weight	311
Barium Minerals	267	Lead Minerals	263
Beryllium Minerals	272	Lithium Minerals	266
Bismuth Minerals	274	Luster Series	300
Block-Mounts	15	Magnesium Minerals	268
Boron Minerals	268	Manganese Minerals	269
Cabinet for Crystals	20	Mercury Minerals	274
Cabinets, Drawer	18	Mining Collection	255
Cadmium Minerals	265	Molybdenum Minerals	273
Caesium Minerals	272	Mounts, Block	15
Calcium Minerals	267	Museum Size Specimens	15
Carbon Minerals	268	Nickel Minerals	268
Cerium (Metals) Minerals	27 I	Odor Series	300
Chemical Mineralogy	307	Ores	253
Chests, Oak	8	Pasteboard Blocks	15
Chromium Minerals	260	Pasteboard Trays	11
Cleavage Series	300	Payment	7
Cobalt Minerals	269	Physical Mineralogy	300
Collections:	-0,	Platinum Minerals	261
Advanced Crystal	276	Potassium Minerals	267
Advanced Systematic	23	Price List, Alphabetical	211
Chemical	308	Price List of Crystals	292
College	27	Prospector's Mining Collection	256
Complete Crystal	277	Quality	7
Complete Type	24	Radio-active Minerals	270
Crystal, Advanced	276	Radium Minerals	270
Crystal, Elementary	296	Rare Element Minerals	270
Crystal, Lecture Table	298	Rare Minerals for Manufacturers	6
High School	242	Selenium Minerals	274
Mining	255	Silicon Minerals	268
Mining	242	Silver Minerals	260
Physical, Hardness, Color, etc.	300	Sizes of Specimens	15
Prospector's	255	Small Size Specimens	17
School	241	Sodium Minerals	267
School of Mines	254	Specific Gravity Series	300
Systematic, Advanced	23	Strontium Minerals	267
Systematic, Elementary	242	Structure Series	300
University	26	Sulphur Minerals	274
Varietal	25	Synopsis of Dana's System	29
Color Series	300	Tantalum Minerals	273
Columbium Minerals	273	Taste Series	300
Copper Minerals	265	Tellurium Minerals	274
Crystal Cabinet	20	Tenacity Series	300
Crystallography	275	Terms	7
Crystallography	31	Thorium Minerals	271
Dana's System, Synopsis of	29	Tin Minerals	272
Economic Mineralogy	253	Titanium Minerals	273
Effect of Radium, etc. Series	300	Transportation Free	7
Elementary Collections	241	Traveling Exhibit	,
Elementary Crystal Collection	295	Trays, Pasteboard	11
Fracture Series	300	Tungsten Minerals	273
Fusibility Series	300	Uranium Minerals	270
Germanium Minerals	272	Vanadium Minerals	273
Gold Minerals	260	Weight, Prices by	311
Guarantee of Quality	7	Yttrium Minerals	271
Hand Size Specimens	17	Zinc Minerals	264
Hardness Series	300	Zirconium Minerals	272
	300		-,-

				]
•				
`				
	•			
•			,	
		,		







CZ53#2

Foote, Warren M.

Complete mineral catalogy

BORROWER'S NAME DATE DUE

### DATE DUE

NOT TO LEA

GAYLORD

ED IN U.S.A.